

IE 521 Convex Optimization

Lecture 3: Separation Theorems

Niao He

2nd May 2019

Outline

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation
Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of Separation, Strict and Strong Separation
Separation Hyperplane Theorem
Strong Separation Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

Quick Review

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation

Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

- ▶ Radon's Theorem
 - ▶ Any set of $d + 2$ points in \mathbb{R}^d can be partitioned into two disjoint sets whose convex hulls intersect.
- ▶ Helley's Theorem
 - ▶ If every $(d + 1)$ of the sets from a collection of n sets in \mathbb{R}^d intersect ($n > d$), then the whole collection of sets intersect.

Question

What's in common and what's different?




Figure: Separation of sets

Warm-up

Quick Review

Questions

Separation
Theorems

Definitions of
Separation, Strict and
Strong Separation

Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of
Alternatives

Farkas' Lemma
LP Duality

Separation of Sets

Definition. Let S and T be two nonempty convex sets in \mathbb{R}^n . A hyperplane $H = \{x \in \mathbb{R}^n : a^T x = b\}$ with $a \neq 0$ is said to separate S and T if $S \cup T \not\subset H$ and

$$S \subset H^- = \{x \in \mathbb{R}^n : a^T x \leq b\}$$

$$T \subset H^+ = \{x \in \mathbb{R}^n : a^T x \geq b\}$$




Figure: Separation of two sets

- Separation is equivalent to say

$$\sup_{x \in S} a^T x \leq \inf_{x \in T} a^T x \text{ and } \inf_{x \in S} a^T x < \sup_{x \in T} a^T x.$$

Strict Separation of Sets

Definition. Let S and T be two nonempty convex sets in \mathbb{R}^n . A hyperplane $H = \{x \in \mathbb{R}^n : a^T x = b\}$ with $a \neq 0$ is said to strictly separate S and T if

$$S \subset H^{--} = \{x \in \mathbb{R}^n : a^T x < b\}$$

$$T \subset H^{++} = \{x \in \mathbb{R}^n : a^T x > b\}$$




Figure: Strict Separation of two sets

Strong Separation of Sets

Definition. Let S and T be two nonempty convex sets in \mathbb{R}^n . A hyperplane $H = \{x \in \mathbb{R}^n : a^T x = b\}$ with $a \neq 0$ is said to strongly separate S and T if there exists $b' < b < b''$ such that


$$S \subset \{x \in \mathbb{R}^n : a^T x \leq b'\}$$
$$T \subset \{x \in \mathbb{R}^n : a^T x \geq b''\}$$

Remark.

- ▶ Strong separation \implies strict separation.
- ▶ Strict separation $\not\Rightarrow$ strong separation.
- ▶ Strong separation is equivalent to say

$$\sup_{x \in S} a^T x < \inf_{x \in T} a^T x.$$

Example



In all examples, S and T are two closed and convex sets

- (A) S and T are separated, but not strictly;
- (B) S and T are strictly separated, but not strongly;
- (C) S and T are strongly separated.

Separation Hyperplane Theorem

Theorem. Let S and T be two nonempty convex sets. Then S and T can be **separated** if and only if

$$\text{rint}(S) \cap \text{rint}(T) = \emptyset.$$

Corollary. Let S be a nonempty convex set and $x_0 \in \partial S$. There exists a supporting hyperplane $H = \{x : a^T x = a^T x_0\}$ with $a \neq 0$ such that

$$S \subset \{x : a^T x \leq a^T x_0\}, \text{ and } x_0 \in H.$$

Warm-up

Quick Review

Questions

Separation
TheoremsDefinitions of
Separation, Strict and
Strong SeparationSeparation
Hyperplane TheoremStrong Separation
Hyperplane TheoremTheorems of
Alternatives

Farkas' Lemma

LP Duality

Proof of Separation Theorem

S and T can be separated iff $\text{rint}(S) \cap \text{rint}(T) = \emptyset$.

► **Necessity.**

- S and T are separated implies that for some $a \neq 0$,

$$\sup_{x \in S} a^T x \leq \inf_{x \in T} a^T x.$$

- If $z \in \text{rint}(S) \cap \text{rint}(T)$, then

$$z = \operatorname{argmax}_{x \in S} \{a^T x\} = \operatorname{argmin}_{x \in T} \{a^T x\}.$$

- The linear function $f(x) = a^T x$ has to be constant on both S and T , i.e., $S \cap T \subset H$. (why?)

Warm-up

Quick Review

Questions

Separation
TheoremsDefinitions of
Separation, Strict and
Strong SeparationSeparation
Hyperplane TheoremStrong Separation
Hyperplane TheoremTheorems of
Alternatives

Farkas' Lemma

LP Duality

Proof of Separation Theorem

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation

Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

S and T can be separated iff $\text{rint}(S) \cap \text{rint}(T) = \emptyset$.

► **Sufficiency.** Based on constructive steps:

1. Separation of a convex set S and $x_0 \notin \text{cl}(S)$ (**key step**);
2. Separation of a convex set S and $x_0 \notin \text{rint}(S)$;
3. Separation of 0 and $\text{rint}(S) - \text{rint}(T)$;
4. Separation of S and T .

Separation of Convex Set and A Point Outside

Proposition. Let S be convex and closed, $x_0 \notin S$. Then x_0 and S can be separated.

Proof. Define

$$d(\{x_0\}, S) := \inf \{\|x_0 - x\|_2 : x \in S\}$$

$$\text{proj}(x_0) := \arg\min_{x \in S} \{\|x_0 - x\|_2\}$$

Then $d(\{x_0\}, S) > 0$ and $\text{proj}(x_0)$ exists and is unique (why?).
The hyperplane

$$H := \{x : a^T x = b\}, \quad a = x_0 - \text{proj}(x_0), \quad b = a^T x_0 - \frac{\|a\|_2^2}{2}$$

separates x_0 and S , i.e. $a^T x < b, \forall x \in S, a^T x_0 > b$. (why?)




Figure: Separation of a convex set and a point

Strong Separation Hyperplane Theorem

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation

Separation
Hyperplane Theorem

Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

Theorem. Let S and T be two nonempty convex sets. Then S and T can be **strongly separated** if and only if

$$\text{dist}(S, T) := \inf \{\|s - t\|_2 : s \in S, t \in T\} > 0.$$

In particular, if $S - T$ is closed and $S \cap T = \emptyset$, then S and T can be strongly separated.

Remarks

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation
Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

- ▶ “ $S - T$ is closed” is only a sufficient condition for strong (strict) separation, not a necessary condition.
- ▶ Even if both S and T are closed convex, $S - T$ might not be closed, and they might not even be strictly separated.
- ▶ When both S and T are closed convex, $S \cap T = \emptyset$ and at least one of them is bounded, then $S - T$ is closed, and S and T can be strongly separated.

Proof of Strong Separation Theorem

S and T can be strongly separated iff $\text{dist}(S, T) > 0$.

- **Necessity.** If S and T are strongly separated, then

$$\exists a \neq 0 : \alpha := \sup_{x \in S} a^T x < \inf_{y \in T} a^T y := \beta.$$

Hence, $\forall x \in S, y \in T$:

$$\|a\|_2 \|y - x\|_2 \geq a^T(y - x) \geq \beta - \alpha \Rightarrow \|y - x\|_2 \geq \frac{\beta - \alpha}{\|a\|_2}.$$

- **Sufficiency.** Suppose $r := \text{dist}(S, T) > 0$, then $(S - T)$ and $B(0, r)$ are two disjoint convex sets. By Separation Theorem, $\exists a \neq 0$,

$$\sup_{z \in S - T} a^T z = \sup_{x \in S, y \in T} a^T(x - y) \leq \inf_{z \in B(0, r)} a^T z < 0.$$

Warm-up

Quick Review

Questions

Separation

Theorems

Definitions of
Separation, Strict and
Strong SeparationSeparation
Hyperplane TheoremStrong Separation
Hyperplane Theorem

Theorems of

Alternatives

Farkas' Lemma

LP Duality

Feasibility of Linear System

Example . Show that the following system have no solution.

$$\begin{cases} x_1 - x_2 + 2x_3 & \leq 0 & \cdots \times 3 \\ -x_1 + x_2 - x_3 & \leq 0 & \cdots \times 5 \\ 2x_1 - x_2 + 3x_3 & \leq 0 & \cdots \times 3 \\ 4x_1 - x_2 + 10x_3 & > 0 \end{cases}$$

$$3 \times Eq.(1) + 5 \times Eq.(2) + 3 \times Eq.(3) \Rightarrow 4x_1 - x_2 + 10x_3 \leq 0.$$

Note the system

$$\begin{cases} y_1 - y_2 + 2y_3 & = 4 \\ -y_1 + y_2 - y_3 & = -1 \\ 2y_1 - y_2 + 3y_3 & = 10 \\ y_1, y_2, y_3 & \geq 0 \end{cases} \text{ has a solution } y = (3, 5, 3).$$

Warm-up

Quick Review

Questions

Separation

Theorems

Definitions of
Separation, Strict and
Strong Separation

Separation
Hyperplane Theorem

Strong Separation
Hyperplane Theorem

Theorems of
Alternatives

Farkas' Lemma

LP Duality

The Celebrated Farkas' Lemma

Theorem. (Farkas' Lemma) Let $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$. Exactly one of the following sets must be empty:

- (i) $\{x \in \mathbb{R}^n : Ax = b, x \geq 0\};$
- (ii) $\{y \in \mathbb{R}^m : A^T y \leq 0, b^T y > 0\}.$

- ▶ System (i) and (ii) are often called strong alternatives, i.e. exactly one of them must be feasible.
- ▶ This is an example of “theorem on alternatives”.



Figure: Gyula Farkas
(1847–1930)


Geometric View of Farkas' Lemma

- Let $A = [a_1 | a_2 | \dots | a_n]$, define the cone
 $\text{Cone} \{a_1, \dots, a_n\} = \{\sum_{i=1}^n x_i a_i : x_i \geq 0, i = 1, \dots, n\}$

$\{Ax = b, x \geq 0\}$ is infeasible

$\iff b \notin \text{Cone} \{a_1, \dots, a_n\}$

$\implies \exists y, y^T a_i \leq 0, \forall i, y^T b > 0$



- Farkas' lemma can be regarded as a special case of the separation theorem.

Warm-up

Quick Review

Questions

Separation

Theorems

Definitions of
Separation, Strict and
Strong SeparationSeparation
Hyperplane TheoremStrong Separation
Hyperplane Theorem

Theorems of

Alternatives

Farkas' Lemma

LP Duality

Proof of Farkas' Lemma

- ▶ System (i) feasible \Rightarrow system (ii) infeasible.
Otherwise, $0 < b^T y = (Ax)^T y = x^T (A^T y) \leq 0$.

- ▶ System (i) infeasible \Rightarrow system (ii) feasible.

Let $C = \text{Cone}\{a_1, \dots, a_n\}$.

Then C is convex and **closed**. And $b \notin C$.

- ▶ By the separation theorem, b and C can be strongly separated, i.e. $\exists y \neq 0 \in \mathbb{R}^m, \gamma \in \mathbb{R}$, s.t.

$$y^T z \leq \gamma, \forall z \in C, y^T b > \gamma.$$

- ▶ Since $0 \in C$, we have $\gamma \geq 0$.
- ▶ Show that $\gamma = 0$. Suppose $\gamma > 0$, and $\exists z_0 \in C$ such that $y^T z_0 > 0$, then $y^T(\alpha z_0) > \gamma$ for α large enough.
- ▶ Since $a_1, \dots, a_n \in C$, we have $y^T a_i \leq 0, \forall i = 1, \dots, m$, i.e., $A^T y \leq 0$.

Remarks

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation
Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

- ▶ The closedness of the cone $\text{Cone}\{a_1, \dots, a_n\}$ is crucial here. Note that in general, when S is not a finite set, $\text{Cone}(S)$ is not always closed.
- ▶ Farkas' Lemma can also be proved by Fourier-Motzkin elimination.
- ▶ Result can be generalized to convex inequalities.

Variant of Farkas' Lemma

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation
Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

Theorem. Exactly one of the following two sets must be empty:

- (i) $\{x \in \mathbb{R}^n : Ax \leq b\}$
- (ii) $\{y \geq 0 : A^T y = 0, b^T y < 0\}$

Theorem. Exactly one of the following two sets must be empty:

- (i) $\{x \in \mathbb{R}^n : Ax = b\}$
- (ii) $\{y \in \mathbb{R}^m : A^T y = 0, b^T y \neq 0\}$

Duality of Linear Program

Warm-up

Quick Review
Questions

Separation
Theorems

Definitions of
Separation, Strict and
Strong Separation
Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of
Alternatives

Farkas' Lemma
LP Duality

Consider the primal and dual pair of linear programs

$$\begin{array}{ll} \min & c^T x \\ (P) \quad & \text{s.t. } Ax = b \\ & x \geq 0 \end{array} \qquad \begin{array}{ll} \max & b^T y \\ (D) \quad & \text{s.t. } A^T y \leq c \end{array}$$

Theorem. (LP Duality) If (P) has a finite optimal value, then so does (D) and the two values equal each other.

Proof: Homework Exercise.

Who introduced LP duality?

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation
Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality



Figure: Leonid
Kantorovich
(1912–1986)




Figure: George
Dantzig (1914–2005)



Figure: John von
Neumann
(1903–1957)

References

Warm-up

Quick Review
Questions

Separation Theorems

Definitions of
Separation, Strict and
Strong Separation
Separation
Hyperplane Theorem
Strong Separation
Hyperplane Theorem

Theorems of Alternatives

Farkas' Lemma
LP Duality

- ▶ Boyd & Vandenberghe, Chapter 2.5
- ▶ Ben-Tal & Nemirovski, Chapter 1.2.5-1.2.6