

IE 521 Convex Optimization

Lecture 17: Interior Point Method

Newton's Method

Niao He

16th April 2019

Outline

Path Following Scheme

Newton's Method

Classical Newton's Method

Affine Invariance

Newton vs GD

Classical Analysis

Newton's Decrement

Newton's Method for Self-Concordant Functions

Damped Newton Method

Path Following Scheme

$$\begin{aligned} \min_x \quad & f(x) \\ \text{s.t.} \quad & x \in X := \{x : g_i(x) \leq 0, i = 1, \dots, m\} \end{aligned} \quad (P)$$

Barrier Method: solve a series of unconstrained problems

$$x^*(t) := \operatorname{argmin}_x \{t \cdot f(x) + F(x)\} \quad (t > 0) \quad (P_t)$$

Barrier Function:

- ▶ $F : \operatorname{int}(X) \rightarrow \mathbb{R}$ and $F(x) \rightarrow +\infty$ as $x \rightarrow \partial(X)$
- ▶ F is twice continuously differentiable and convex
- ▶ F is *non-degenerate*, i.e. $\nabla^2 F(x) \succ 0, \forall x \in \operatorname{int}(X)$

Central Path:

$$x^*(t) \in \operatorname{int}(X) \longrightarrow x^*, \text{ as } t \longrightarrow \infty$$

Path Following Scheme

Question: Need to specify

1. the barrier function $F(x)$?

▶ **Self-concordant barriers**, e.g..

$$F(x) = -\sum_{i=1}^m \log(-g_i(x))$$

2. the method to solve unconstrained problems (P_t)?

▶ **Newton's method**

3. the policy to update the penalty parameter t ?

Classical Newton's Method

Assume $f(x)$ is twice continuously differentiable on \mathbb{R}^n .

$$\min_{x \in \mathbb{R}^n} f(x)$$

Newton's Method:

$$x_{k+1} = x_k - [\nabla^2 f(x_k)]^{-1} \nabla f(x_k), k = 0, 1, 2, \dots$$

- ▶ Newton's method can break down if $f(x)$ is degenerate.
- ▶ $d_k = -\nabla^2 f(x_k)^{-1} \nabla f(x_k)$ is called Newton's's direction.
- ▶ Newton's direction is not necessarily a descent direction.

Classical Newton's Method: Interpretation

$$x_{k+1} = x_k - [\nabla^2 f(x_k)]^{-1} \nabla f(x_k), k = 0, 1, 2, \dots \quad (\star)$$

► $(\star) \iff$ *minimizing quadratic approximation of f*

► Recall Taylor expansion of $f(x)$:

$$f(x + h) = f(x) + \nabla f(x)^T h + \frac{1}{2} h^T \nabla^2 f(x) h + o(\|h\|^2)$$

► (\star) is the solution to the quadratic approximation

$$x_{k+1} = \min_x \left\{ f(x_k) + \nabla f(x_k)^T (x - x_k) + \frac{1}{2} (x - x_k)^T \nabla^2 f(x_k) (x - x_k) \right\}$$

Remark. When f is quadratic and non-degenerate, Newton's method converges in one step.

Classical Newton's Method: Interpretation

Path Following Scheme

Newton's Method

Classical Newton's Method

Affine Invariance

Newton vs GD

Classical Analysis

Newton's Decrement

Newton's Method for Self-Concordant

Functions

Damped Newton Method

$$x_{k+1} = x_k - [\nabla^2 f(x_k)]^{-1} \nabla f(x_k), k = 0, 1, 2, \dots \quad (\star)$$

► $(\star) \iff$ *solving linearized optimality condition*

- From first-order optimality condition: $\nabla f(x) = 0$
- Taylor expansion:

$$\nabla f(x + h) \approx \nabla f(x) + \nabla^2 f(x)h$$

- (\star) is the solution to the linear system:

$$\nabla f(x_k) + \nabla^2 f(x_k)(x - x_k) = 0$$

Affine Invariance of Newton's Method

Newton's method is invariant w.r.t. affine transformation.

- ▶ Let A be non-singular and consider the function

$$\hat{f}(y) = f(Ay).$$

- ▶ The Newton steps for f and \hat{f} are

$$x_{k+1} = x_k - [\nabla^2 f(x_k)]^{-1} \nabla f(x_k)$$

$$\begin{aligned} y_{k+1} &= y_k - [\nabla^2 \hat{f}(y_k)]^{-1} \nabla \hat{f}(y_k) \\ &= y_k - A^{-1} [\nabla^2 f(Ay_k)]^{-1} \nabla f(Ay_k) \end{aligned}$$


- ▶ If $y_0 = A^{-1}x_0$, then $y_k = A^{-1}x_k$.
- ▶ Newton's method follows the same trajectory in the 'x-space' and 'y-space'.

Newton's Method vs Gradient Descent

$$x_{k+1} = x_k - \nabla^2 f(x_k)^{-1} \nabla f(x_k) \quad (\text{Newton})$$

$$x_{k+1} = x_k - \gamma_k \nabla f(x_k) \quad (\text{GD})$$

- ▶ Affine vs non-affine invariant
- ▶ Second-order vs. first-order
- ▶ Expensive vs. cheap iteration
- ▶ Local vs. global convergence



Newton vs. GD

Figure from Tibshirani lecture notes

Illustration: Convergence of Newton's Method

Consider the nonconvex function:

$$f(x) = x^3$$

Q. Does Newton's method converge? How fast?

$$f'(x) = 3x^2, \quad f''(x) = 6x$$

$$x_{k+1} = x_k - (6x_k)^{-1} \cdot 3x_k^2 = \frac{1}{2} \cdot x_k \quad (\text{Newton})$$

- Converges in a linear rate to a stationary point

Illustration: Convergence of Newton's Method

Consider the strictly convex function:

$$f(x) = \sqrt{1 + x^2}$$

Q. Does Newton's method converge? How fast?

$$f'(x) = \frac{x}{\sqrt{1 + x^2}}, \quad f''(x) = \frac{1}{(1 + x^2)^{3/2}}$$

$$x_{k+1} = x_k - (1 + x_k^2)^{3/2} \frac{x_k}{\sqrt{1 + x_k^2}} = -x_k^3 \quad (\text{Newton})$$

- ▶ if $|x_0| < 1$, converges in a cubic rate (extremely fast)
- ▶ if $|x_0| = 1$, oscillates between 1 and -1
- ▶ if $|x_0| > 1$, diverges

Local Quadratic Convergence

Theorem. Assume that

- ▶ f has a Lipschitz Hessian: for some $M > 0$,

$$\|\nabla^2 f(x) - \nabla^2 f(y)\|_2 \leq M\|x - y\|_2$$

- ▶ f has a strict local minimum x^* : for some $\mu > 0$,

$$\nabla^2 f(x^*) \succeq \mu I$$

- ▶ The initial point x_0 is close enough to x^* :

$$\|x_0 - x^*\|_2 \leq \frac{\mu}{2M}$$

Then Newton's method is well-defined and converges to x^* at a quadratic rate

$$\|x_{k+1} - x^*\|_2 \leq \frac{M}{\mu} \|x_k - x^*\|_2^2.$$

Local Quadratic Convergence

Corollary. It follows that

$$\begin{aligned}\frac{M}{\mu} \|x_k - x^*\|_2 &\leq \left[\frac{M}{\mu} \|x_{k-1} - x^*\|_2 \right]^2 \\ &\leq \dots \\ &\leq \left[\frac{M}{\mu} \|x_0 - x^*\|_2 \right]^{2^k} \\ &\leq \left(\frac{1}{2} \right)^{2^k}\end{aligned}$$

- The number of iterations to achieve an accuracy ϵ , i.e. $\|x_k - x^*\|_2 \leq \epsilon$, is at most

$$k \geq \log_2 \log_2 \left(\frac{M}{\mu \epsilon} \right)$$

- The above results hold true for any unconstrained minimization regardless of convexity.

Lemma on Hessian Lipschitzness

Lemma. Assume f has a Lipschitz Hessian with constant M , then for any x, y ,

$$\|\nabla f(y) - \nabla f(x) - \nabla^2 f(x)(y - x)\|_2 \leq \frac{M}{2} \|y - x\|_2^2.$$

Proof.

$$\begin{aligned} & \|\nabla f(y) - \nabla f(x) - \nabla^2 f(x)(y - x)\|_2 \\ &= \left\| \int_0^1 \nabla^2 f(x + t(y - x))(y - x) dt - \nabla^2 f(x)(y - x) \right\|_2 \\ &= \left\| \int_0^1 \left[\nabla^2 f(x + t(y - x)) - \nabla^2 f(x) \right] (y - x) dt \right\|_2 \\ &\leq \int_0^1 M \cdot t \|y - x\|_2^2 dt \\ &= \frac{M}{2} \|y - x\|_2^2 \end{aligned}$$

Proof of Local Convergence

First, we have

$$\begin{aligned}x_{k+1} - x^* &= x_k - x^* - [\nabla^2 f(x_k)]^{-1} \nabla f(x_k) \\&= [\nabla^2 f(x_k)]^{-1} [\nabla^2 f(x_k)(x_k - x^*) - \nabla f(x_k)] \\&= [\nabla^2 f(x_k)]^{-1} [\nabla f(x^*) - \nabla f(x_k) - \nabla^2 f(x_k)(x^* - x_k)]\end{aligned}$$

$$\Rightarrow \|x_{k+1} - x^*\|_2 \leq \|[\nabla^2 f(x_k)]^{-1}\|_2 \cdot \frac{M}{2} \|x_k - x^*\|_2^2$$

We can show by induction that

$$\begin{aligned}\|x_k - x^*\|_2 &\leq \frac{\mu}{2M} \\ \nabla^2 f(x_k) &\succeq \frac{\mu}{2} I\end{aligned}$$

This concludes the proof.

Local Convergence for Strongly Convex Functions

Theorem. Assume that

- ▶ f has a Lipschitz Hessian with constant $M > 0$;
- ▶ f is μ -strongly convex: $\nabla^2 f(x) \succeq \mu I, \forall x$;
- ▶ The initial point x_0 satisfies $\|\nabla f(x_0)\|_2 \leq \frac{2\mu^2}{M}$.

Then the gradient converges to zero quadratically

$$\|\nabla f(x_{k+1})\|_2 \leq \frac{M}{2\mu^2} \|\nabla f(x_k)\|_2^2.$$

Proof. This is because

$$\begin{aligned} \|\nabla f(x_{k+1})\|_2 &= \|\nabla f(x_{k+1}) - \nabla f(x_k) - \nabla^2 f(x_k)(x_{k+1} - x_k)\|_2 \\ &\leq \frac{M}{2} \left\| \left[\nabla^2 f(x_k) \right]^{-1} \nabla f(x_k) \right\|_2^2 \\ &\leq \frac{M}{2} \left\| \left[\nabla^2 f(x_k) \right]^{-1} \right\|_2^2 \cdot \|\nabla f(x_k)\|_2^2 \\ &\leq \frac{M}{2\mu^2} \|\nabla f(x_k)\|_2^2 \end{aligned}$$

Issue with Affine Invariance

- ▶ Recall that Newton's method is invariant w.r.t. affine transformations.
- ▶ The region of quadratic convergence should not depend on the Euclidean metric.
- ▶ However, in the classical analysis, the assumption and the measure of error, e.g. the Lipschitz continuity of Hessian, depend heavily on the Euclidean metric and is not affine invariant.
- ▶ A natural remedy is to assume self-concordance.
- ▶ Self concordant function are especially well suited for Newton method.

Newton's Decrement

Definition. Newton decrement is defined as :

$$\lambda_f(x) = \sqrt{\nabla f(x) [\nabla^2 f(x)]^{-1} \nabla f(x)}$$

- Relates to the decrease of the second order Taylor expansion after a Newton step:

$$f(x) - \min_h \left\{ f(x) + h^T \nabla f(x) + \frac{1}{2} h^T \nabla^2 f(x) h \right\} = \frac{1}{2} \lambda_f^2(x)$$

- Can be viewed as an approximate bound of the suboptimality gap $f(x) - f^*$.
- Newton decrement is also *affine-invariant*.

Newton's Decrement vs Local Norms

$$\lambda_f(x) = \sqrt{\nabla f(x)[\nabla^2 f(x)]^{-1}\nabla f(x)}$$

- Equals to the local norm of Newton's direction $d(x)$:

$$\|d(x)\|_x = \| -\nabla^2 f(x)^{-1}\nabla f(x) \|_x = \lambda_f(x)$$

- Equals to the conjugate local norm of $\nabla f(x)$:

$$\|\nabla f(x)\|_{x,*} = \|[\nabla^2 f(x)]^{-1/2}\nabla f(x)\|_2 = \lambda_f(x)$$

Newton's Decrement and Self-concordance

Recall that standard self-concordant functions f has nice properties inside the Dikin ellipsoid: $\forall y : \|y - x\|_x = \gamma < 1$,

$$(1) \quad y \in \text{dom}(f)$$

$$(2) \quad (1 - r)^2 \nabla^2 f(x) \preceq \nabla^2 f(y) \preceq \frac{1}{(1-r)^2} \nabla^2 f(x)$$

$$(3) \quad \frac{\gamma^2}{1+\gamma} \leq \langle \nabla f(y) - \nabla f(x), y - x \rangle \leq \frac{\gamma^2}{1-\gamma}$$

$$(4) \quad \omega(\gamma) \leq f(y) - f(x) - \langle \nabla f(x), y - x \rangle \leq \omega_*(\gamma), \text{ where } \omega(\gamma) = \gamma - \ln(1 + \gamma), \omega_*(\gamma) = -\gamma - \ln(1 - \gamma).$$

Proposition.

- ▶ If $\lambda_f(x) < 1$, the point $x_+ = x - d(x) \in \text{dom}(f)$
- ▶ If x^* is a minimizer of f , then $\lambda_f(x^*) = 0$
- ▶ If $\lambda_f(x_0) < 1$ for some $x_0 \in \text{dom}(f)$, then f has a unique minimizer.

Newton's Decrement and Self-concordance

Proposition. If $\lambda_f(x_0) < 1$ for some $x_0 \in \text{dom}(f)$, then f has a unique minimizer.

Proof.

Suffice to show the level set $\{y : f(y) \leq f(x_0)\}$ is bounded.

$$\begin{aligned} f(y) &\geq f(x_0) + \langle \nabla f(x_0), y - x_0 \rangle + \omega(\|y - x_0\|_{x_0}) \\ &\geq f(x_0) - \|\nabla f(x_0)\|_{x_0, *} \cdot \|y - x_0\|_{x_0} + \omega(\|y - x_0\|_{x_0}) \\ &= f(x_0) - \lambda_f(x_0) \cdot \|y - x_0\|_{x_0} + \omega(\|y - x_0\|_{x_0}) \end{aligned}$$

$$\text{Hence, } f(y) \leq f(x_0) \implies \frac{\omega(\|y - x_0\|_{x_0})}{\|y - x_0\|_{x_0}} \leq \lambda_f(x_0) < 1$$

Note the function $\phi(t) = \frac{\omega(t)}{t} = 1 - \frac{1}{t} \ln(1+t)$ is strictly increasing in $t \geq 0$. Hence, $\|y - x_0\|_{x_0} \leq t^*$ for some t^* .

Example: Newton's Decrement

Consider the self-concordant function

$$f(x) = \epsilon x - \ln(x)$$

with $\text{dom}(f) := \{x : x > 0\}$.

$$\lambda_f(x) = \sqrt{\left(\epsilon - \frac{1}{x}\right) \left(\frac{1}{x^2}\right)^{-1} \left(\epsilon - \frac{1}{x}\right)} = |1 - \epsilon x|$$

- ▶ When $\epsilon \leq 0$, $\lambda_f(x) \geq 1$, and the function is unbounded below and there does not exist a minimizer.
- ▶ When $\epsilon > 0$, $\lambda_f(x) < 1$, for $x \in (0, \frac{2}{\epsilon})$, there exists a unique minimizer $x^* = \frac{1}{\epsilon}$.

Affine-invariant Metrics

Newton method: initialize $x_0 \in \text{dom}(f)$ and update via

$$x_{k+1} = x_k - [\nabla^2 f(x_k)]^{-1} \nabla f(x_k), k = 0, 1, 2, \dots$$

Accuracy Measure:

- ▶ Function gap: $f(x_k) - f(x^*)$
- ▶ Newton's decrement: $\lambda_f(x_k) = \|\nabla f(x_k)\|_{x_k, *}$
- ▶ Local distance to the minimizer: $\|x_k - x^*\|_{x_k}$
- ▶ Distance to the minimizer: $\|x_k - x^*\|_{x^*}$

Remark. Indeed, all of these measures are independent of Euclidean metric and equivalent locally.

Affine-invariant Metrics

Proposition. (Nesterov, 2004) When $\lambda_f(x) < 1$, we have

1. $f(x) - f(x^*) \leq \omega_*(\lambda_f(x)) \leq \frac{\lambda_f(x)^2}{2(1-\lambda_f(x))^2}$
2. $\|x - x^*\|_x \leq \frac{\lambda_f(x)}{1-\lambda_f(x)}$
3. $\|x - x^*\|_{x^*} \leq \frac{\lambda_f(x)}{1-\lambda_f(x)}$

We will focus mainly on the convergence in terms of $\lambda_f(x)$.

Local Convergence of Self-concordant Functions

Theorem. If $x_k \in \text{dom}(f)$ and $\lambda_k < 1$, then $x_{k+1} \in \text{dom}(f)$ and

$$\lambda_{k+1} \leq \left(\frac{\lambda_k}{1 - \lambda_k} \right)^2.$$

Remark. Let λ^* be such that $\frac{\lambda^*}{(1-\lambda^*)^2} = 1$.

- ▶ If $\lambda_k < \lambda^*$, $\lambda_{k+1} < \lambda_k$.
- ▶ Region of quadratic convergence is

$$\lambda_f(x) \leq \lambda^* = \frac{3 - \sqrt{5}}{2} \approx 0.38.$$

- ▶ Still might diverge if not started with a point with $\lambda_f(x)$ small enough.

Q. How to ensure global convergence?

Proof of Local Convergence

- Note $\|x_{k+1} - x_k\|_{x_k} = \lambda_f(x_k) = \lambda_k < 1$, so $x_{k+1} \in \text{dom}(f)$.
- It holds that

$$\nabla^2 f(x_{k+1}) \succeq (1 - \lambda_k)^2 \nabla^2 f(x_k)$$

$$\lambda_{k+1} \leq \frac{1}{1 - \lambda_k} \sqrt{\nabla f(x_{k+1})^T [\nabla^2 f(x_k)]^{-1} \nabla f(x_{k+1})}$$

- Note

$$\begin{aligned} \nabla f(x_{k+1}) &= \nabla f(x_{k+1}) - \nabla f(x_k) - [\nabla^2 f(x_k)](x_{k+1} - x_k) \\ &= \underbrace{\left[\int_0^1 \nabla^2 f(x_k + t(x_{k+1} - x_k)) - \nabla^2 f(x_k) dt \right]}_G (x_{k+1} - x_k) \end{aligned}$$

- Hence,

$$\lambda_{k+1} \leq \frac{1}{1 - \lambda_k} \sqrt{(x_{k+1} - x_k)^T G^T [\nabla^2 f(x_k)]^{-1} G (x_{k+1} - x_k)}$$

Proof of Local Convergence (Cont'd)

- Further,

$$\lambda_{k+1} \leq \frac{\lambda_k}{1 - \lambda_k} \underbrace{\|[\nabla^2 f(x_k)]^{-1/2} G [\nabla^2 f(x_k)]^{-1/2}\|_2}_H$$

- Note that

$$G \succeq \nabla^2 f(x_k) \int_0^1 \left[(1 - t\lambda_k)^2 - 1 \right] dt = \left(\frac{\lambda_k^2}{3} - \lambda_k \right) \nabla^2 f(x_k)$$

$$G \preceq \nabla^2 f(x_k) \int_0^1 \left[\frac{1}{(1 - t\lambda_k)^2} - 1 \right] dt = \frac{\lambda_k}{1 - \lambda_k} \nabla^2 f(x_k)$$

- This implies that

$$\|H\|_2 \leq \max \left\{ \lambda_k - \frac{\lambda_k^2}{3}, \frac{\lambda_k}{1 - \lambda_k} \right\} = \frac{\lambda_k}{1 - \lambda_k}.$$

Damped Newton Method

Damped Newton method: initialize $x_0 \in \text{dom}(f)$ and update via

$$x_{k+1} = x_k - \frac{1}{1 + \lambda_f(x_k)} [\nabla^2 f(x_k)]^{-1} \nabla f(x_k)$$

Remark. Damped Newton procedure is always well-defined:

$$\|x_{k+1} - x_k\|_{x_k} = \frac{\lambda_f(x_k)}{1 + \lambda_f(x_k)} < 1 \Rightarrow x_{k+1} \in W_1^0(x_k) \subseteq \text{dom}(f).$$

Global Convergence of Damped Newton

Theorem. The damped Newton method satisfies that

1. (Descent phase) $\forall k \geq 0$,

$$f(x_{k+1}) \leq f(x_k) - \omega(\lambda_f(x_k)).$$

2. (Quadratic convergence phase) If $\lambda_k(x_k) < \frac{1}{4}$, then

$$\lambda_f(x_{k+1}) \leq 2[\lambda_f(x_k)]^2.$$

Proof.

$$\begin{aligned} f(x_{k+1}) &\leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \omega_*(\|x_{k+1} - x_k\|_{x_k}) \\ &= f(x_k) - \frac{\lambda_f(x_k)}{1 + \lambda_f(x_k)} + \omega_*\left(\frac{\lambda_f(x_k)}{1 + \lambda_f(x_k)}\right) \end{aligned}$$

where $\omega_*(t) = -t - \ln(1 - t)$.

Iteration Complexity of Damped Newton Method

Remark.

- *Damped Newton stage*: when $\lambda_f(x_k) \geq \beta \in (0, 1/4)$

$$f(x_{k+1}) \leq f(x_k) - \omega(\beta) \Rightarrow N_1 \leq \frac{f(x_0) - f(x^*)}{\omega(\beta)}.$$

- *Damped/Basic Newton stage*: when $\lambda_f(x_k) < \beta$

$$\lambda_f(x_{k+1}) \leq 2 \left[\lambda_f(x_k) \right]^2 \Rightarrow N_2 \leq O(1) \log_2 \log_2 \left(\frac{1}{\epsilon} \right).$$

The total complexity to find a solution with $\lambda_f(x) \leq \epsilon$:

$$O(1) \left[f(x_0) - f^* + \log \log \left(\frac{1}{\epsilon} \right) \right]$$

References

- ▶ Nesterov (2004), Introductory Lectures on Convex Optimization, Chapter 4.1.4-5
- ▶ Nemirovski (2004), Interior Point Polynomial Time Methods in Convex Programming, Chapter 1