

IE 521 Convex Optimization

Lecture 12: Conic Programming

Niao He

5th March 2019

Outline

Generalized Inequality

Conic Programs

General Conic Program

LP

SOCP

SDP

Examples of Conic Programs

Norm Minimization

Sparse Group Lasso

Robust Linear Program

Extend Linear Programs to Convex Programs

IE 521 Convex
Optimization

Niao He

Generalized
Inequality

► Traditional approach:

$$\begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax - b \geq 0 \end{array} \quad \Rightarrow \quad \begin{array}{ll} \min_x & f_0(x) \\ \text{s.t.} & f_i(x) \geq 0, i = 1, \dots, m \end{array}$$

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

Extend Linear Programs to Convex Programs

IE 521 Convex
Optimization

Niao He

Generalized
Inequality

- Traditional approach:

$$\begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax - b \geq 0 \end{array} \quad \Rightarrow \quad \begin{array}{ll} \min_x & f_0(x) \\ \text{s.t.} & f_i(x) \geq 0, i = 1, \dots, m \end{array}$$

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

- Modern approach:

$$\begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax \geq b \end{array} \quad \Rightarrow \quad \begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax \succcurlyeq b \end{array}$$

Extend Linear Programs to Convex Programs

IE 521 Convex
Optimization

Niao He

Generalized
Inequality

- ▶ Traditional approach:

$$\begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax - b \geq 0 \end{array} \quad \Rightarrow \quad \begin{array}{ll} \min_x & f_0(x) \\ \text{s.t.} & f_i(x) \geq 0, i = 1, \dots, m \end{array}$$

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

- ▶ Modern approach:

$$\begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax \geq b \end{array} \quad \Rightarrow \quad \begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax \succcurlyeq b \end{array}$$

Q. What properties should the vector inequality “ \succcurlyeq ” satisfy?

“Good” Vector Inequalities

The partial ordering “ \succcurlyeq ” on \mathbb{R}^m should satisfy:

- (i) Reflexibility: $a \succcurlyeq a$
- (ii) Anti-symmetry: $a \succcurlyeq b$ and $b \succcurlyeq a$ implies $a = b$
- (iii) Transitivity: $a \succcurlyeq b$ and $b \succcurlyeq c$ implies $a \succcurlyeq c$
- (iv) Homogeneity: $a \succcurlyeq b$ and $\lambda \geq 0$ implies $\lambda a \succcurlyeq \lambda b$
- (v) Additivity: $a \succcurlyeq b$ and $c \succcurlyeq b$ implies $a + c \succcurlyeq b + d$

“Good” Vector Inequalities

The partial ordering “ \succcurlyeq ” on \mathbb{R}^m should satisfy:

- (i) Reflexibility: $a \succcurlyeq a$
- (ii) Anti-symmetry: $a \succcurlyeq b$ and $b \succcurlyeq a$ implies $a = b$
- (iii) Transitivity: $a \succcurlyeq b$ and $b \succcurlyeq c$ implies $a \succcurlyeq c$
- (iv) Homogeneity: $a \succcurlyeq b$ and $\lambda \geq 0$ implies $\lambda a \succcurlyeq \lambda b$
- (v) Additivity: $a \succcurlyeq b$ and $c \succcurlyeq b$ implies $a + c \succcurlyeq b + d$

Remark.

$$a \succcurlyeq b \iff a - b \succcurlyeq 0 \iff a - b \in \mathcal{K}$$

where $\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$.

“Good” Set \mathcal{K}

$$\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$$

“Good” Set \mathcal{K}

$$\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$$

1. \mathcal{K} is nonempty: $0 \in \mathcal{K}$

“Good” Set \mathcal{K}

$$\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$$

1. \mathcal{K} is nonempty: $0 \in \mathcal{K}$
2. \mathcal{K} is pointed:

$$a, -a \in \mathcal{K} \Rightarrow a = 0$$

“Good” Set \mathcal{K}

$$\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$$

1. \mathcal{K} is nonempty: $0 \in \mathcal{K}$
2. \mathcal{K} is pointed:

$$a, -a \in \mathcal{K} \Rightarrow a = 0$$

3. \mathcal{K} is closed w.r.t multiplication of nonnegative scalars:

$$a \in \mathcal{K}, \lambda \geq 0 \Rightarrow \lambda a \in \mathcal{K}$$

“Good” Set \mathcal{K}

$$\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$$

1. \mathcal{K} is nonempty: $0 \in \mathcal{K}$

2. \mathcal{K} is pointed:

$$a, -a \in \mathcal{K} \Rightarrow a = 0$$

3. \mathcal{K} is closed w.r.t multiplication of nonnegative scalars:

$$a \in \mathcal{K}, \lambda \geq 0 \Rightarrow \lambda a \in \mathcal{K}$$

4. \mathcal{K} is closed w.r.t addition:

$$a, b \in \mathcal{K} \Rightarrow a + b \in \mathcal{K}$$

“Good” Set \mathcal{K}

$$\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$$

1. \mathcal{K} is nonempty: $0 \in \mathcal{K}$

2. \mathcal{K} is pointed:

$$a, -a \in \mathcal{K} \Rightarrow a = 0$$

3. \mathcal{K} is closed w.r.t multiplication of nonnegative scalars:

$$a \in \mathcal{K}, \lambda \geq 0 \Rightarrow \lambda a \in \mathcal{K}$$

4. \mathcal{K} is closed w.r.t addition:

$$a, b \in \mathcal{K} \Rightarrow a + b \in \mathcal{K}$$

“Good” Set \mathcal{K}

$$\mathcal{K} = \{a \in \mathbb{R}^m : a \succcurlyeq 0\}$$

1. \mathcal{K} is nonempty: $0 \in \mathcal{K}$

2. \mathcal{K} is pointed:

$$a, -a \in \mathcal{K} \Rightarrow a = 0$$

3. \mathcal{K} is closed w.r.t multiplication of nonnegative scalars:

$$a \in \mathcal{K}, \lambda \geq 0 \Rightarrow \lambda a \in \mathcal{K}$$

4. \mathcal{K} is closed w.r.t addition:

$$a, b \in \mathcal{K} \Rightarrow a + b \in \mathcal{K}$$

\mathcal{K} must be a nonempty pointed convex cone!

Example: Coordinate-wise Vector Inequality

The usual coordinate-wise vector inequality:

$$a \geq b \iff a_i - b_i \geq 0, i = 1, \dots, m$$

Example: Coordinate-wise Vector Inequality

The usual coordinate-wise vector inequality:

$$a \geq b \iff a_i - b_i \geq 0, i = 1, \dots, m$$

Nonnegative orthant cone:

$$\mathcal{K} = \mathbb{R}_+^m := \{x = [x_1; \dots; x_m] : x_i \geq 0, i = 1, \dots, m\}$$




Figure: Nonnegative orthant

Example: Coordinate-wise Vector Inequality

The usual coordinate-wise vector inequality:

$$a \geq b \iff a_i - b_i \geq 0, i = 1, \dots, m$$

Nonnegative orthant cone:

$$\mathcal{K} = \mathbb{R}_+^m := \{x = [x_1; \dots; x_m] : x_i \geq 0, i = 1, \dots, m\}$$




Figure: Nonnegative orthant

Remark. $\mathcal{K} = \mathbb{R}_+^m$ is also closed and has non-empty interior.

Regular Cone and Generalized Inequality

Generalized Inequality

- $\mathcal{K} \subseteq \mathbb{R}^m$ is a regular cone if it is a closed pointed convex cone with nonempty interior.

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

Regular Cone and Generalized Inequality

Generalized Inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

- ▶ $\mathcal{K} \subseteq \mathbb{R}^m$ is a regular cone if it is a closed pointed convex cone with nonempty interior.
- ▶ Define a generalized inequality:

$$a \geq_{\mathcal{K}} b \iff a - b \in \mathcal{K}$$

Regular Cone and Generalized Inequality

Generalized Inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program


- ▶ $\mathcal{K} \subseteq \mathbb{R}^m$ is a regular cone if it is a closed pointed convex cone with nonempty interior.
- ▶ Define a generalized inequality:

$$a \geq_{\mathcal{K}} b \iff a - b \in \mathcal{K}$$

- ▶ Strict inequality:


$$a >_{\mathcal{K}} b \iff a - b \in \text{int}(\mathcal{K})$$

Examples



Ice cream cone

$$L = \{(x_1, x_2, y) \in \mathbb{R}^3 : \sqrt{x_1^2 + x_2^2} \leq y\}$$



Positive semidefinite cone

$$S = \{(x, y, z) \in \mathbb{R}^3 : \begin{bmatrix} x & y/\sqrt{2} \\ y/\sqrt{2} & z \end{bmatrix} \succeq 0\}$$

Figure from Vandenberghe lecture notes

Conic (Linear) Program

Generalized
inequality

Conic Programs

General Conic
Program

LP

SOCF

SDP

Examples of Conic
Programs

Norm Minimization

Sparse Group Lasso

Robust Linear
Program

$$\begin{array}{ll} \min_x & c^T x \\ \text{s.t.} & Ax \geq_{\mathcal{K}} b \end{array} \quad (\text{CP})$$

where

- ▶ \mathcal{K} is a regular cone,
- ▶ $Ax \geq_{\mathcal{K}} b$ iff $Ax - b \in \mathcal{K}$.

Linear Program (LP)

Generalized
Inequality

Conic Programs

General Conic
Program

LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

► Nonnegative Orthant:

$$\mathcal{K} = \mathbb{R}_+^m = \{x \in \mathbb{R}^m : x_i \geq 0, i = 1, \dots, m\}$$

► Linear Programs (LP):

$$\min_x \left\{ c^T x : a_i^T x - b_i \geq 0, i = 1, \dots, m \right\} \quad (LP)$$

Second Order Conic Program (SOCP)

- Lorentz Cone (second order/ice cream cone):

$$\mathcal{K} = L^{n_1} \times \dots \times L^{n_m}$$

$$L^n = \{x \in \mathbb{R}^n : x_n \geq \sqrt{\sum_{i=1}^{n-1} x_i^2}\}$$

- Second Order Conic Program:

$$\min_x \left\{ c^T x : \|D_i x - d_i\|_2 \leq e_i^T x - f_i, i = 1, \dots, m \right\}$$

(SOCP)

Second Order Conic Program (SOCP)

- Lorentz Cone (second order/ice cream cone):

$$\mathcal{K} = L^{n_1} \times \dots \times L^{n_m}$$

$$L^n = \{x \in \mathbb{R}^n : x_n \geq \sqrt{\sum_{i=1}^{n-1} x_i^2}\}$$

- Second Order Conic Program:

$$\min_x \left\{ c^T x : \| D_i x - d_i \|_2 \leq e_i^T x - f_i, i = 1, \dots, m \right\} \quad (SOCP)$$

- Here the linear map is

$$Ax - b := \left[\underbrace{D_1 x - d_1; e_1^T x - f_1}_{\mathbb{R}^{n_1}}; \dots; \underbrace{D_m x - d_m; e_m^T x - f_m}_{\mathbb{R}^{n_m}} \right]$$

Semidefinite Program (SDP)

► Positive Semidefinite Cone:

$$\mathcal{K} = S_+^m = \{X \in S^m : X \succeq 0\}$$

► Semidefinite Programs:

$$\min_x \left\{ c^T x : \sum_{i=1}^n x_i A_i - B \succeq 0 \right\} \quad (SDP)$$

Generalized
inequality

Conic Programs

General Conic
Program

LP

SOCP

SDP

Examples of Conic
Programs

Norm Minimization

Sparse Group Lasso

Robust Linear
Program

Semidefinite Program (SDP)

► Positive Semidefinite Cone:

$$\mathcal{K} = S_+^m = \{X \in S^m : X \succeq 0\}$$

► Semidefinite Programs:

$$\min_x \left\{ c^T x : \sum_{i=1}^n x_i A_i - B \succeq 0 \right\} \quad (SDP)$$

► Here the linear map is

$$\mathcal{A}(x) - b = \sum_{i=1}^n x_i A_i - B : \mathbb{R}^n \rightarrow S^m$$

with $A_1, \dots, A_n, B \in S^m$.

LP, SOCP, and SDP

$$\text{LP} \subseteq \text{SOCP} \subseteq \text{SDP}$$

Generalized
Inequality

Conic Programs

General Conic
Program

LP

SOCP

SDP

► $\text{LP} \subseteq \text{SOCP}$:

Examples of Conic
Programs

Norm Minimization

Sparse Group Lasso

Robust Linear
Program

► $\text{SOCP} \subseteq \text{SDP}$:

LP, SOCP, and SDP

$$\text{LP} \subseteq \text{SOCP} \subseteq \text{SDP}$$

- $\text{LP} \subseteq \text{SOCP}$:

$$a_i^T x - b_i \geq 0 \Leftrightarrow \begin{bmatrix} 0 \\ a_i^T x - b_i \end{bmatrix} \in L^2$$

- $\text{SOCP} \subseteq \text{SDP}$:

Generalized
Inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

LP, SOCP, and SDP

$$\text{LP} \subseteq \text{SOCP} \subseteq \text{SDP}$$

► $\text{LP} \subseteq \text{SOCP}$:

$$a_i^T x - b_i \geq 0 \Leftrightarrow \begin{bmatrix} 0 \\ a_i^T x - b_i \end{bmatrix} \in L^2$$

► $\text{SOCP} \subseteq \text{SDP}$:

$$\begin{bmatrix} x_1 \\ \vdots \\ x_{n-1} \\ x_n \end{bmatrix} \in L^n \Leftrightarrow \begin{bmatrix} x_n & x_1 & x_2 & \dots & x_{n-1} \\ x_1 & x_n & & & \\ x_2 & & x_n & & \\ \vdots & & & \ddots & \\ x_{n-1} & & & & x_n \end{bmatrix} \succeq 0$$

Schur Complement

Proposition. Let $S = \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix}$ be symmetric with $R \succ 0$.

$S \succeq 0$ if and only if $P - Q^T R^{-1} Q \succeq 0$.

Schur Complement

Proposition. Let $S = \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix}$ be symmetric with $R \succ 0$.

$S \succeq 0$ if and only if $P - Q^T R^{-1} Q \succeq 0$.

Proof.

$$S \succeq 0 \Leftrightarrow \forall u, v : \begin{bmatrix} u^T & v^T \end{bmatrix} \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} \geq 0$$

Schur Complement

Proposition. Let $S = \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix}$ be symmetric with $R \succ 0$.

$S \succeq 0$ if and only if $P - Q^T R^{-1} Q \succeq 0$.

Proof.

$$\begin{aligned} S \succeq 0 &\Leftrightarrow \forall u, v : \begin{bmatrix} u^T & v^T \end{bmatrix} \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} \geq 0 \\ &\Leftrightarrow \inf_{u, v} u^T P u + 2v^T Q u + v^T R v \geq 0 \end{aligned}$$

Schur Complement

Proposition. Let $S = \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix}$ be symmetric with $R \succ 0$.

$S \succeq 0$ if and only if $P - Q^T R^{-1} Q \succeq 0$.

Proof.

$$\begin{aligned} S \succeq 0 &\Leftrightarrow \forall u, v : \begin{bmatrix} u^T & v^T \end{bmatrix} \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} \geq 0 \\ &\Leftrightarrow \inf_{u, v} u^T P u + 2v^T Q u + v^T R v \geq 0 \\ &\Leftrightarrow \inf_u u^T (P - Q^T R^{-1} Q) u \geq 0 \end{aligned}$$

Schur Complement

Proposition. Let $S = \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix}$ be symmetric with $R \succ 0$.

$S \succeq 0$ if and only if $P - Q^T R^{-1} Q \succeq 0$.

Proof.

$$\begin{aligned} S \succeq 0 &\Leftrightarrow \forall u, v : \begin{bmatrix} u^T & v^T \end{bmatrix} \begin{bmatrix} P & Q^T \\ Q & R \end{bmatrix} \begin{bmatrix} u \\ v \end{bmatrix} \geq 0 \\ &\Leftrightarrow \inf_{u, v} u^T P u + 2v^T Q u + v^T R v \geq 0 \\ &\Leftrightarrow \inf_u u^T (P - Q^T R^{-1} Q) u \geq 0 \\ &\Leftrightarrow P - Q^T R^{-1} Q \succeq 0 \end{aligned}$$

Remarks

Generalized
inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

- ▶ In principle, any convex problem can be represented as a conic program.
- ▶ Other cones exist: exponential cone, power cone, copositive cone, etc.
- ▶ In practice, only the primitive cones (LP, SOCP, SDP) are used, limited by the available algorithms and solvers.
- ▶ Indeed, an extremely wide spectrum of convex problems can be converted into three standard conic programs.

Example: ℓ_2 -norm minimizationGeneralized
inequality

Conic Programs

General Conic
Program
LP
SOCP
SDPExamples of Conic
Programs

Norm Minimization

Sparse Group Lasso
Robust Linear
Program

$$\begin{aligned} \min_{x \in \mathbb{R}^n} \|x\|_2 &\iff \min_{x,t} t &\iff \min_{x,t} t \\ \text{s.t. } t &\geq \|x\|_2 &\text{s.t. } \begin{bmatrix} x \\ t \end{bmatrix} \succeq_{L^{n+1}} 0 \end{aligned}$$

$$\begin{aligned} \min_{x \in \mathbb{R}^n} x^T x &\iff \min_{x,t} t &\iff \min_{x,t} t \\ \text{s.t. } t &\geq x^T x &\text{s.t. } \begin{bmatrix} 2x \\ t-1 \\ t+1 \end{bmatrix} \succeq_{L^{n+2}} 0 \end{aligned}$$

Example: Convex Quadratic Programming

Let $Q = LL^T \succeq 0$.

$$\min_{x \in \mathbb{R}^n} x^T Q x + q^T x$$

Generalized
inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

Example: Convex Quadratic Programming

IE 521 Convex
Optimization

Niao He

Let $Q = LL^T \succeq 0$.

$$\min_{x \in \mathbb{R}^n} \quad x^T Q x + q^T x$$

$$\begin{aligned} &\iff \min_{x, t} \quad t \\ &\text{s.t.} \quad \begin{bmatrix} 2L^T x \\ t - q^T x - 1 \\ t - q^T x + 1 \end{bmatrix} \succeq_{L^{n+2}} 0 \end{aligned}$$

Generalized
inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization
Sparse Group Lasso
Robust Linear
Program

Example: Spectral Norm Minimization

Let A_1, \dots, A_n be symmetric matrix in $\mathbb{R}^{p \times p}$.

$$\min_{x \in \mathbb{R}^n} \lambda_{\max} \left(\sum_{i=1}^n x_i A_i \right)$$

Generalized
inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization

Sparse Group Lasso
Robust Linear
Program

Example: Spectral Norm Minimization

Let A_1, \dots, A_n be symmetric matrix in $\mathbb{R}^{p \times p}$.

$$\min_{x \in \mathbb{R}^n} \lambda_{\max} \left(\sum_{i=1}^n x_i A_i \right)$$

$$\begin{aligned} \iff \min_{x, t} \quad & t \\ \text{s.t.} \quad & t \cdot I_p - \sum_{i=1}^n x_i A_i \succeq 0 \end{aligned}$$

Example: Spectral Norm Minimization

Let A_1, \dots, A_n be general matrix in $\mathbb{R}^{p \times q}$.

$$\min_{x \in \mathbb{R}^n} \sigma_{\max} \left(\sum_{i=1}^n x_i A_i \right)$$

Generalized
inequality

Conic Programs

General Conic
Program
LP
SOCP
SDP

Examples of Conic
Programs

Norm Minimization

Sparse Group Lasso
Robust Linear
Program

Example: Spectral Norm Minimization

Let A_1, \dots, A_n be general matrix in $\mathbb{R}^{p \times q}$.

$$\min_{x \in \mathbb{R}^n} \sigma_{\max} \left(\sum_{i=1}^n x_i A_i \right)$$

$$\begin{aligned} &\iff \min_{x, t} \quad t \\ &\text{s.t.} \quad \begin{bmatrix} t \cdot I_p & \sum_{i=1}^n x_i A_i \\ \sum_{i=1}^n x_i A_i^T & t \cdot I_q \end{bmatrix} \succeq 0 \end{aligned}$$

Example: Sparse Group Lasso

$$\min_w \left\{ \|Xw - y\|_2^2 + \lambda \sum_{i=1}^p \|w_i\|_2 \right\}$$

- ▶ $X \in \mathbb{R}^{m \times n}, y \in \mathbb{R}^m$ are the input data
- ▶ $w = [w_1; \dots; w_p] \in \prod_{i=1}^p \mathbb{R}^{n_i}$ is the decision variable.

Example: Sparse Group Lasso

$$\min_w \left\{ \|Xw - y\|_2^2 + \lambda \sum_{i=1}^p \|w_i\|_2 \right\}$$

- ▶ $X \in \mathbb{R}^{m \times n}, y \in \mathbb{R}^m$ are the input data
- ▶ $w = [w_1; \dots; w_p] \in \prod_{i=1}^p \mathbb{R}^{n_i}$ is the decision variable.

$$\begin{aligned} \min_{w, t_0, t_1, \dots, t_p} \quad & t_0 + \lambda \sum_{i=1}^p t_i \\ \text{s.t.} \quad & \begin{bmatrix} 2(Xw - y) \\ t_0 - 1 \\ t_0 + 1 \end{bmatrix} \succeq_{L^{m+2}} 0 \quad (SOCP) \\ & \begin{bmatrix} w_i \\ t_i \end{bmatrix} \succeq_{L^{n_i+1}} 0, \quad \forall i = 1, \dots, p. \end{aligned}$$

Example: Robust Linear Program

- Recall the robust linear program

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & a_i^T x \leq b_i, \forall a_i \in \mathcal{E}_i, i = 1, \dots, m \end{aligned}$$

where $\mathcal{E}_i = \{\bar{a}_i + P_i u : \|u\|_2 \leq 1\}$.

Example: Robust Linear Program

- Recall the robust linear program

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & a_i^T x \leq b_i, \forall a_i \in \mathcal{E}_i, i = 1, \dots, m \end{aligned}$$

where $\mathcal{E}_i = \{\bar{a}_i + P_i u : \|u\|_2 \leq 1\}$.

- This is equivalent to the convex program

$$\begin{aligned} \min_x \quad & c^T x \\ \text{s.t.} \quad & \bar{a}_i^T x + \|P_i^T x\|_2 \leq b_i, i = 1, \dots, m \end{aligned}$$

which is an SOCP.

Example: Robust Portfolio Selection

- Assume that returns $r_i, i = 1, \dots, n$ are exactly known.
To maximize the return leads to

$$\max_x \quad r^T x, \text{ s.t. } x \geq 0, \mathbf{1}^T x = 1$$

Example: Robust Portfolio Selection

- Assume that returns $r_i, i = 1, \dots, n$ are exactly known.
To maximize the return leads to

$$\max_x r^T x, \text{ s.t. } x \geq 0, \mathbf{1}^T x = 1$$

- Now assume the returns are known within an ellipsoid:

$$\mathcal{U} = \left\{ \hat{r} + \rho \hat{\Sigma}^{1/2} u : \|u\|_2 \leq 1 \right\}$$

where \hat{r} and $\hat{\Sigma}$ are the empirical mean and covariance.
The robust portfolio problem is

$$\max_x \min_{r \in \mathcal{U}} r^T x, \text{ s.t. } x \geq 0, \mathbf{1}^T x = 1$$

Example: Robust Portfolio Selection

- Assume that returns $r_i, i = 1, \dots, n$ are exactly known.
To maximize the return leads to

$$\max_x r^T x, \text{ s.t. } x \geq 0, \mathbf{1}^T x = 1$$

- Now assume the returns are known within an ellipsoid:

$$\mathcal{U} = \left\{ \hat{r} + \rho \hat{\Sigma}^{1/2} u : \|u\|_2 \leq 1 \right\}$$

where \hat{r} and $\hat{\Sigma}$ are the empirical mean and covariance.
The robust portfolio problem is

$$\max_x \min_{r \in \mathcal{U}} r^T x, \text{ s.t. } x \geq 0, \mathbf{1}^T x = 1$$

- This is equivalent as an SOCP:

$$\max_x \hat{r}^T x - \rho \|\hat{\Sigma}^{1/2} x\|_2, \text{ s.t. } x \geq 0, \mathbf{1}^T x = 1$$

References

Generalized
Inequality

Conic Programs

General Conic
Program

LP

SOCP

SDP

Examples of Conic
Programs

Norm Minimization

Sparse Group Lasso

**Robust Linear
Program**

- Ben-Tal & Nemirovski (2013), Chapter 1.4