


BMJ Open Effects of aspirin and omega-3 fatty acids on age-related macular degeneration in ASCEND-Eye: a randomised placebo-controlled trial in a population with diabetes

Emily Sammons , Louise Bowman, William Stevens, Georgina Buck, Imen Hammami, Sarah Parish, Jane Armitage, on behalf of the ASCEND-Eye Collaborative Group

To cite: Sammons E, Bowman L, Stevens W, *et al.* Effects of aspirin and omega-3 fatty acids on age-related macular degeneration in ASCEND-Eye: a randomised placebo-controlled trial in a population with diabetes. *BMJ Open* 2025;**15**:e090605. doi:10.1136/bmjopen-2024-090605

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-090605>).

Received 28 June 2024
Accepted 10 January 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY. Published by BMJ Group.

Nuffield Department of Population Health, University of Oxford, Oxford, UK

Correspondence to

Dr Emily Sammons;
emily.sammons@ndph.ox.ac.uk

ABSTRACT

Purpose Aspirin and omega-3 fatty acids (FAs) are potential disease modifiers of age-related macular degeneration (AMD), but previous studies have produced inconsistent findings. Randomised evidence for the efficacy and safety of aspirin and omega-3 FAs on AMD is presented in this study.

Design ASCEND-Eye is a substudy of eye effects in the 2×2 factorial design ASCEND (A Study of Cardiovascular Events in Diabetes) double-blind, randomised, placebo-controlled trial for the primary prevention of cardiovascular events. Reports of AMD diagnoses were sourced from 6 monthly ASCEND follow-up questionnaires and a Visual Function Questionnaire.

Participants 15 480 UK adults at least 40 years of age with diabetes but no evident cardiovascular disease.

Interventions 100 mg aspirin daily versus placebo and, separately, 1 g omega-3 FAs daily versus placebo.

Main outcome measure The first post-randomisation reports of AMD.

Results During 7.4 years of follow-up, 122 (1.6%) participants randomised to aspirin were reported as having AMD, compared with 138 (1.8%) randomised to placebo (rate ratio 0.88; 95% CI 0.69 to 1.12; *p*=0.31). AMD occurred in 130 (1.7%) participants randomised to omega-3 FAs, compared with 130 (1.7%) randomised to placebo (rate ratio 0.99; 95% CI 0.78 to 1.27; *p*=0.99).

Conclusion No clinically-meaningful effects of aspirin or omega-3 FAs on AMD were found. Although the study had very limited statistical power to detect clinically relevant effects, these data overcome some methodological limitations of previous observational studies, providing randomised evidence of both treatments on AMD, which could contribute to future meta-analyses.

Trial registration number ISRCTN60635500 and NCT00135226.

INTRODUCTION

Age-related macular degeneration (AMD) is a leading cause of central visual loss in older people worldwide.¹ Aspirin and omega-3

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The strengths of our study include its randomised design, well-defined exposure, a large number of participants and long duration of near-complete follow-up.
- ⇒ ASCEND (A Study of Cardiovascular Events in Diabetes)-Eye was underpowered to detect modest but plausible effects of each treatment. Therefore, it was not possible to compare the effects of aspirin or omega-3 fatty acids on prognostically important risk factors for age-related macular degeneration, including age and smoking.

fatty acids (FAs) are potential modifiers of the disease process, but previous studies of their relationships with AMD have produced inconsistent findings. The long-term use of aspirin for more than 15 years has been associated with harmful effects on the wet form of AMD in some large observational studies^{2–3}; however, two randomised trials reported non-significant trends towards a beneficial effect on their combined late AMD endpoint.^{4–5} Separately, observational studies have associated a higher consumption of oily fish or omega-3 FAs with a lower risk of AMD,^{6–10} but three previous randomised trials specifically designed to explore this relationship have reported null findings.^{11–13} A limitation of observational studies is their inability to account for residual confounding from unmeasured or unknown variables, even after applying advanced statistical methods. This problem becomes more apparent in multifactorial diseases such as AMD, where the level of risk conferred by different genetic, inflammatory, vascular or environmental factors is difficult to quantify.¹⁴ Random assignment

will distribute these characteristics evenly between each treatment group, and thereby minimise the risk of bias.

The ASCEND (A Study of Cardiovascular Events in Diabetes) trial in a diabetic population¹⁵ provided the opportunity to undertake randomised comparisons of aspirin and, separately, omega-3 FAs on participant-reported AMD events. Although they share some of the same risk factors, such as age and smoking, diabetes and AMD are not considered to be pathologically related to each other.^{16–18} The analyses we describe were part of the ASCEND-Eye substudy.¹⁹

METHODS

Trial oversight

Investigators from the Clinical Trial Service Unit at the University of Oxford designed and coordinated ASCEND-Eye. Multi-centre Research Ethics Committee (MREC) approval was granted for the substudy by the North West MREC in October 2016. Members of the writing committee vouch for the faithfulness of the substudy to the data analysis plan.

Participants and procedures

The design and main results of ASCEND have been described in detail elsewhere.^{15 20 21} Briefly, the trial used mail-based methods to follow-up 15 480 participants recruited in the UK between 2005 and 2011. They comprised men and women with any type of diabetes (excluding gestational diabetes), who were at least 40 years of age, with no previous history of cardiovascular disease, no contraindication to either study treatment and no pre-existing life-limiting medical condition. Using minimised randomisation, eligible participants were assigned in a 2×2 factorial masked design, between 100 mg aspirin daily or a matching placebo and, separately, between 1 g omega-3 fatty acid capsules (containing 460 mg eicosapentaenoic acid (EPA) and 380 mg docosahexaenoic acid (DHA)) daily or a matching placebo. Although not specifically requested on the forms, AMD events could be reported on ASCEND follow-up questionnaires sent to participants every 6 months (see online supplemental material) in response to a general question about ‘any other serious illness or admission to hospital’. Participants could also report AMD events by telephoning the central coordinating office, where a clinician clarified the diagnosis during the call.

While ASCEND was ongoing, ASCEND-Eye sought evidence from participants’ primary care providers to support the adjudication of AMD events by trial clinicians. Events were confirmed against prespecified criteria while masked to the study treatment allocations.¹⁹ This included documented evidence that an ophthalmologist had confirmed the diagnosis on a date closely matching that given by the participant. Typically, this included clinic letters describing the result of retinal imaging investigations and the subsequent treatment plan. Data on which eye was affected and whether it was the wet or dry form was

not routinely collected. Visual Function Questionnaires (VFQ) were sent to all participants who were alive and on web-based or mail-based follow-up when ASCEND ended on 31 July 2017 (see online supplemental material).¹⁹ The VFQ consisted of two parts: a bespoke first page of questions that explicitly sought new diagnoses of serious eye conditions, including AMD, followed by the standard National Eye Institute Visual Function Questionnaire-25 (NEI-VFQ-25).²² The date of the first AMD diagnosis was collected, but information was not requested regarding the phenotypic form or which eye was affected. In total, 8846 out of 11 301 (78.3%) participants who were eligible to receive the VFQ responded.¹⁹ Events originating from the VFQ were not adjudicated due to the delay between event reporting and the last communication with the participant’s primary care providers. Every randomised participant from ASCEND (n=15 480) was included in the analyses, all of whom gave their written informed consent.

Outcomes

The effects of allocation to aspirin compared with placebo, and, separately, omega-3 FAs compared with placebo, on time to first confirmed or unrefuted post-randomisation diagnosis of AMD in either eye were a prespecified secondary analysis of the ASCEND-Eye substudy, which has been described in detail previously.¹⁹ The results of our safety analyses, which included confirmed incidences of sight-threatening eye bleeds in the aspirin randomisation, have also been published elsewhere.²³

Statistical analysis

The data analysis plan was publicly available on the trial website (<https://ascend.medsci.ox.ac.uk>) before unmasking the ASCEND-Eye AMD results.²⁰

Age-related macular degeneration is a condition that becomes more common with advancing age. In a meta-analysis of studies that used comparable, standardised protocols and comprised data on nearly 25 000 individuals from high-income countries with predominantly white ethnicity, the estimated age-specific prevalence of AMD was less than 0.5% in those aged between 50 and 60, but this increased to 12% and 16% in men and women over 80 years of age, respectively.²⁴ The ASCEND population was relatively young (mean age=63.3; SD 9.2)¹⁵; therefore, ASCEND-Eye had fewer incident AMD events and thus a lower power to detect a clinically-meaningful treatment effect than would be expected in an older population. The sample size could not be increased because ASCEND-Eye was a substudy of a larger trial. Power calculations^{25 26} based on masked data indicated that there was only 24–41% power to detect 15–20% proportional reductions in AMD events at a two-sided p value of less than 0.05.

The logrank^{27 28} test was used to conduct intention-to-treat analyses of time from randomisation to AMD diagnosis between those randomised to aspirin compared with placebo and, separately, omega-3 FAs versus placebo. Average event rate ratios and their 95% CIs and two-sided

p values were calculated using the one-step method from the 'observed minus expected' numbers of events (O-E) and their variances (V) outputted from the SAS LIFETEST procedure (event rate ratio= $\exp(O-E/V)$).²⁸ The results are represented graphically in the form of Kaplan-Meier plots. It was anticipated that the factorial design of ASCEND would have little or no effect on the statistical sensitivity with which the effects of each treatment arm could be assessed.^{27 29} Moreover, no clinically significant interactions between the study treatments were anticipated.³⁰ The main comparisons of the effect of aspirin were therefore made without stratification by omega-3 FAs allocation (and vice versa for the effect of omega-3 FAs analyses), but post-hoc stratified analyses were carried out to check the validity of this assumption. As AMD was more likely to be recorded on a VFQ form than on an ASCEND follow-up form, analyses were stratified by the availability of a VFQ. No formal adjustments for multiplicity were made for these secondary analyses of ASCEND-Eye.¹⁹ Therefore, two-tailed p values of less than 0.05 and the 95% CI should be interpreted with caution.

To interpret the impact of adjudication on the AMD outcome, the percentage agreement and Cohen's Kappa (κ) test of agreement were performed between the pre-adjudicated and post-adjudicated categorisation of AMD events. These prespecified analyses were restricted to the first reports of AMD, where the information source was the participant's 6month follow-up questionnaire for ASCEND. Phoned-in events were excluded from the agreement analyses because they may have been more accurately recorded than events reported on in-trial follow-up questionnaires or the VFQ without physician assistance. A post-hoc exploratory analysis assessed the proportion of AMD events derived from the VFQ only, in-trial follow-up questionnaires only, or both.

The Clinical Trial Service Unit at the University of Oxford holds the trial database. All analyses were performed using SAS V. 9.4 and R V. 4.1.2 (R Core Team 2021-11-01) statistical software.

Patient and public involvement

It was not appropriate or possible to involve patients or the public in the design, or conduct, or reporting, or dissemination plans of our research.

RESULTS

The baseline characteristics of the ASCEND population and those who returned a VFQ have been described elsewhere.¹⁹ Key prognostic variables, such as age, smoking status and ethnicity were well-balanced between the randomised groups (table 1 and online supplemental table S1). For the full randomised population of ASCEND, the mean duration of follow-up was 7.4 years (online supplemental table S2), with approximately 114 000 person-years of follow-up split evenly between the treatment arms (online supplemental table S3). Complete morbidity follow-up was available for 15 341 (99.1%)

of the randomised population (online supplemental table S4). The study's average adherence to aspirin and omega-3 FAs was 68.0% and 75.6%, respectively, with similar proportions in the corresponding placebo arms (online supplemental table S5). Adherence declined in the later years of follow-up (online supplemental table S5), with increasing use of non-study aspirin, alternative antiplatelet agents or anticoagulants in the aspirin randomisation (non-study antiplatelet therapy use was 7.9% in the placebo arm and 6.9% in the aspirin arm; 9.6% and 8.4% respectively for non-study antiplatelet or anticoagulant therapy, online supplemental table S6). Study average adherence rates in prespecified baseline characteristic groups were similar in each treatment arm (online supplemental table S7). In both randomisations, 'Participant wishes' represented the main reason for discontinuing treatment (online supplemental table S8). There were no significant differences in the number of participants who stopped taking active treatment as compared with the placebo, in either randomisation, overall, or for specific reasons, except that stopping due to minor bleeding or bruising was more prevalent among those who were randomised to aspirin (3.0%) compared with placebo (1.8%).

The baseline characteristics of VFQ respondents were representative of the full randomised ASCEND population; however, there were statistically significant differences between those included and excluded from the exercise for some characteristics (online supplemental table S9). For example, those excluded were slightly older, and more frequently they were current or former smokers. VFQ forms were completed with a mean of 8.6 years from randomisation (online supplemental table S10). Those who returned a VFQ had better study average adherence, but similar follow-up and patterns of adherence to the overall ASCEND population (online supplemental tables S11–S14).

Effects of the study treatments on AMD events

AMD was recorded on the VFQ form by 213 out of 8846 (2.4%) participants, but fewer cases were captured by the ASCEND follow-up forms (80/15 480, 0.5%), with 33 AMD events reported by both methods (online supplemental table S15). The proportion of participants who reported an AMD diagnosis on an ASCEND follow-up questionnaire was the same for the VFQ respondents (46/8846, 0.5%) and non-recipient or non-respondents (34/6634, 0.5%, online supplemental table S13). Out of a total of 260 AMD events, 73 were confirmed by adjudication, seven were unrefuted because supporting evidence was not provided by the participants' General Practitioner (GP) to adjudicate the event, and 180 were unadjudicated, having derived from the VFQ alone.

There was no significant effect of randomisation to aspirin compared with placebo on AMD diagnoses, which were reported by 122 (1.6%) participants assigned aspirin, compared with 138 (1.8%) assigned placebo (rate ratio 0.88; 95% CI 0.69 to 1.12; $p=0.31$; figure 1). The survival

Table 1 Baseline characteristics					
Baseline characteristic	Aspirin randomisation		Omega-3 FAs randomisation		
	Active (n=7740)	Placebo (n=7740)	Active (n=7740)	Placebo (n=7740)	Overall (n=15480)
Age at randomisation (years)					
Mean (SD)	63.2±9.2	63.3±9.2	63.3±9.2	63.3±9.2	63.3±9.2
Sex					
Male	4843 (62.6)	4841 (62.5)	4842 (62.6)	4842 (62.6)	9684 (62.6)
Female	2897 (37.4)	2899 (37.5)	2898 (37.4)	2898 (37.4)	5796 (37.4)
Type of diabetes*					
Type 1	458 (5.9)	453 (5.9)	460 (5.9)	451 (5.8)	911 (5.9)
Type 2	7282 (94.1)	7287 (94.1)	7280 (94.1)	7289 (94.2)	14569 (94.1)
Duration of diabetes (years)					
Median (IQR)	7 (3–13)	7 (3–13)	7 (3–12)	7 (3–13)	7 (3–13)
Systolic blood pressure (mm Hg)†					
Mean (SD)	136.1±15.2	136.2±15.3	136.2±15.4	136.2±15.1	136.2±15.3
Diastolic blood pressure (mm Hg)†					
Mean (SD)	77.0±9.4	77.2±9.5	77.1±9.5	77.1±9.5	77.1±9.5
Body mass index (kg/m²)‡					
Mean (SD)	30.8±6.3	30.6±6.3	30.7±6.3	30.8±6.2	30.7±6.3
Cigarette smoking					
Current	639 (8.3)	640 (8.3)	639 (8.3)	640 (8.3)	1279 (8.3)
Former	3526 (45.6)	3525 (45.5)	3527 (45.6)	3524 (45.5)	7051 (45.5)
Never	3489 (45.1)	3488 (45.1)	3489 (45.1)	3488 (45.1)	6977 (45.1)
Unknown	86 (1.1)	87 (1.1)	85 (1.1)	88 (1.1)	173 (1.1)
Non-study medication					
ACE-inhibitor or ARB	4520 (58.4)	4535 (58.6)	4569 (59.0)	4486 (58.0)	9055 (58.5)
Aspirin use before screening	2740 (35.4)	2768 (35.8)	2744 (35.5)	2764 (35.7)	5508 (35.6)
Thiazide or related diuretic	1480 (19.1)	1477 (19.1)	1448 (18.7)	1509 (19.5)	2957 (19.1)
Calcium channel blocker	1926 (24.9)	1847 (23.9)	1912 (24.7)	1861 (24.0)	3773 (24.4)
Statin	5854 (75.6)	5799 (74.9)	5791 (74.8)	5862 (75.7)	11653 (75.3)
Total cholesterol (mmol/L)					
Mean (SD)	4.2±0.9	4.2±0.9	4.2±0.9	4.2±0.9	4.2±0.9
HDL cholesterol (mmol/L)					
Mean (SD)	1.3±0.4	1.3±0.4	1.3±0.4	1.3±0.4	1.3±0.4

Continued

Table 1 Continued

Baseline characteristic	Aspirin randomisation		Omega-3 FAs randomisation	
	Active (n=7740)	Placebo (n=7740)	Active (n=7740)	Placebo (n=7740)
Non-HDL cholesterol (mmol/L)				
Mean (SD)	2.9±0.9	2.9±0.8	2.9±0.9	2.9±0.8
Glycosylated haemoglobin				
IFCC (mmol/mol) mean (SD)	54.7±12.9	54.9±12.9	54.9±13.0	54.8±12.9
CKD-EPI estimated GFR (mL/min/1.73 m ²)§				
Mean (SD)	85.2±21.1	85.2±21.1	85.2±21.3	85.2±21.1
Urinary albumin:creatinine ratio (mg/mmol)¶				
Median (IQR)	0.56 (0.00–1.33)	0.55 (0.18–1.34)	0.56 (0.18–1.36)	0.55 (0.14–1.32)
Ethnic origin				
White	7467	7468	7467	7468
	(96.5%)	(96.5%)	(96.5%)	(96.5%)
	14 935	14 935	14 935	14 935
	(96.5%)	(96.5%)	(96.5%)	(96.5%)

Figures presented are counts with percentages unless otherwise stated. Percentages may not total 100 because of rounding.

*The presence of type 2 diabetes was based on a broad clinical definition involving the participant's age at the diagnosis of diabetes, the use of insulin within 1 year after diagnosis and the body mass index.

†From blood and urine consent forms, generally before randomisation.

‡The body mass index (the weight in kilograms divided by the square of the height in metres) was based on values for height and weight the participants reported on their randomisation questionnaires.

§Calculated from blood cystatin c concentration using the CKD-EPI formula (Inker LA, Schmid CH, Tighiouart H, *et al.* Estimating glomerular filtration rate from serum creatinine and cystatin C. *New England Journal of Medicine* 2012; 367(1): 20–9).

¶There was an analysis rule in ASCEND which stated that those with a below-detectable threshold albumin component of their urinary albumin creatinine ratio would be recorded as zero. This applied to just over 25% of participants in the active arm of each treatment randomisation, and to just under 25% of participants in the placebo arm of each treatment randomisation. Hence the IQR included zero.

ACE, angiotensin-converting enzyme; ARB, angiotensin receptor blockers; ASCEND, A Study of Cardiovascular Events in Diabetes; CKD-EPI, Chronic Kidney Disease Epidemiology Collaboration; FA, fatty acids; GFR, glomerular filtration rate; HDL, high-density lipoprotein; IFCC, International Federation of Clinical Chemistry.




Figure 1 Kaplan-Meier plot of time to confirmed or unrefuted diagnoses of age-related macular degeneration by aspirin allocation. The number of participants at risk at the start of each year of follow-up is shown. The rate ratio is for confirmed or unrefuted AMD diagnoses among participants in the aspirin arm, as compared with those in the placebo arm, stratified by the availability of a Visual Functioning Questionnaire. The corresponding unstratified rate ratio was 0.88 (95% CI 0.69 to 1.13; p=0.31). AMD, age-related macular degeneration.

curves appear to separate after approximately 6 years of follow-up, but the difference did not become statistically significant. Similarly, there was no significant effect of randomisation to omega-3 FAs compared with placebo.

AMD occurred in 130 (1.7%) participants allocated to omega-3 FAs, compared with 130 (1.7%) allocated to placebo (rate ratio 0.99; 95% CI 0.78 to 1.27; p=0.97; [figure 2](#)). In exploratory analyses, the proportional effects




Figure 2 Kaplan-Meier plot of time to confirmed or unrefuted diagnoses of age-related macular degeneration by omega-3 fatty acid allocation. The number of participants at risk at the start of each year of follow-up is shown. The rate ratio is for confirmed or unrefuted AMD diagnoses among participants in the active omega-3 FAs arm, compared with those in the placebo arm, stratified by availability of a Visual Functioning Questionnaire. The corresponding unstratified rate ratio was 1.00 (95% CI 0.78 to 1.27; p=0.99). AMD, age-related macular degeneration; FAs, fatty acids.

Table 2 Effect of adjudication on first participant-reported age-related macular degeneration events

	Adjudicator		Total
	AMD	No AMD	
Participant			
AMD	50	12	62
No AMD	23	595	618
Total	73	607	680

% of agreement=94.9%; Cohen's κ : 0.71.
 Events were restricted to the first reports of AMD recorded on the participant's 6 monthly follow-up questionnaires for ASCEND and not the VFQ. Phoned-in events and those where supporting evidence was not provided by the participants' General Practitioner to adjudicate the event were excluded.
 AMD, age-related macular degeneration; ASCEND, A Study of Cardiovascular Events in Diabetes; VFQ, Visual Function Questionnaire.

of aspirin and, separately, omega-3 FAs on AMD events did not vary by other treatment assignment ($p=0.74$ for interaction in the aspirin model and $p=0.76$ in the omega-3 FA model; online supplemental table S16).

Effect of adjudication on AMD event categorisation

GPs provided evidence to support the adjudication of 680 out of 725 eye-related adverse events recorded by participants on their ASCEND 6 monthly follow-up forms. The clinical adjudicator agreed with the participant that an AMD event had or had not occurred in 94.9% of cases, with a κ score of 0.71, indicating substantial inter-rater reliability (table 2).³¹

DISCUSSION

In the ASCEND-Eye substudy of a large randomised placebo-controlled trial, there were no clinically-meaningful effects of allocation to aspirin 100mg daily or 1g omega-3 FAs daily for 7.4 years on confirmed or unrefuted AMD diagnoses.

Aspirin has been considered of potential benefit in AMD owing to its complementary, non-selective and irreversible inhibitive effects on two isoforms of the COX enzyme: COX-1 and COX-2. The long-term suppression of platelet aggregation via acetylation of COX-1 slows atherosclerotic cardiovascular disease progression, which might also protect against retinal arteriolar narrowing and the deposition of lipids in Bruch's membrane.³² Meanwhile, COX-2 inhibition may reduce the platelet-mediated release of vascular endothelial growth factors^{33–35} and the expression of pro-inflammatory prostaglandins involved in the biogenesis of drusen.^{36 37} Collectively, these actions might protect against choroidal neovascularisation and disruption of the retinal pigment epithelium. Conversely, the vasoconstrictive effect of COX-2-suppressed prostacyclin synthesis may cause hypoxia in older people with narrowed choroidal blood vessels, becoming the stimulus

for neovascularisation and the development of the wet form of AMD.³

Our study complements the findings of three other large, double-masked, randomised placebo-controlled trials of aspirin for the primary prevention of cardiovascular events^{4 5} or death, dementia and persistent physical disability.^{38–40} The Physician's Health Study tested 325 mg of aspirin on alternate days in US Physicians aged 40–84 years.⁴ The trial, which stopped early because there were clear reductions in myocardial infarction events, included 22 071 men who did not report having AMD at baseline.⁴ After 5 years of treatment, there were fewer AMD diagnoses in those assigned aspirin compared with those in the placebo group (51 (0.5%) vs 66 (0.6%); rate ratio 0.77; 95% CI 0.54 to 1.11).⁴ Similar results were reported in the Women's Health Study, which tested 100 mg aspirin on alternate days in female healthcare professionals, aged 45 years or older.⁵ After 10 years of follow-up, among 39 421 women without AMD at baseline, there was a non-significant trend towards a beneficial effect with the active treatment (111 (0.5%) vs 134 (0.7%); HR 0.82; 95% CI 0.64 to 1.06).⁵ Like ASCEND-Eye, in which there is a suggestion that the survival curves may separate in favour of aspirin after 6 years of follow-up, there was a separation of the curves after 3 years of follow-up in the Women's Health Study, but overall, neither trial reached statistical significance. Finally, a substudy of the ASPirin in Reducing Events in the Elderly (ASPREE) randomised trial tested 100 mg of aspirin daily in 3171 people aged 70 years and older without dementia, independence-limiting physical disability or cardiovascular disease.⁴¹ AMD status was ascertained by experienced graders using retinal photography and the Beckman classification⁴² of AMD severity. After 3 years of treatment, there were no significant effects of aspirin on cumulative AMD incidence among 1983 participants without AMD at baseline (195 of 1004 (19.4%) vs 187 of 979 (19.1%); relative risk 1.02; 95% CI 0.85 to 1.22).⁴¹ However, all four trials were underpowered, and the duration of follow-up may have needed to be longer to corroborate observational evidence of harm with more prolonged aspirin exposure (>15 years) and wet AMD.^{2 3} Therefore, obtaining further data regarding the effects of aspirin on AMD over a more extended follow-up period may be important. It may be possible to obtain long-term follow-up data of our participants by linking to their electronic UK National Ophthalmology Database AMD Audit records if an application to secure Section 251 exemption to access patient identifiers is approved.⁴³ If it was confirmed that harmful effects take longer to emerge, it might be appropriate to offer older people prescribed aspirin more frequent eye examinations, given the potential to prevent rapid visual loss from wet AMD with anti-Vascular Endothelial Growth Factor (VEGF) therapies.

Experimental studies have shown that omega-3 FAs, in particular DHA, could have a protective role against the development or progression of AMD via complex antiangiogenic and antioxidant pathways, localised to

the retina.⁴⁴ In support of this, most, but not all, observational studies have found that affected individuals tend to recall a lower consumption of fish or omega-3 FAs.^{7 9 45 46} However, our results are consistent with three previous randomised trials of omega-3 FAs, which also failed to demonstrate any effects.^{11–13} The Age-Related Eye Disease Study-2 (AREDS2) trial randomised 4203 men and women aged 50–85 with bilateral early disease or late AMD in only one eye, in a 2×2 factorial design between 10mg lutein plus 2mg zeaxanthin daily or a matching placebo and, separately, between omega-3 FA capsules (containing 650mg EPA and 350mg DHA) daily or a matching placebo.¹¹ The primary outcome was incident wet or dry AMD identified using annual mydriatic fundus photographs.¹¹ During 4.9 years of follow-up, allocation to active omega-3 FAs did not prevent AMD progression: 979 out of 3491 eyes (28.0%) in the active treatment group developed late disease, compared with 961 out of 3400 eyes (28.3%) in the placebo group (HR 0.98; 95% CI 0.89 to 1.08; *p*=0.74).¹¹ Similar results were reported in the Vitamin D and omega-3 triAL (VITAL-AMD) study, which randomised 25 871 men over 50 and women over 55 years of age, in a 2×2 factorial masked design, to 2000 units of cholecalciferol daily or a matching placebo and, separately, to an omega-3 FA supplement (containing 460mg EPA and 380mg DHA) daily or a matching placebo.¹³ The primary outcome of total AMD was a composite of incident AMD or AMD progression based on participant-reported events confirmed by medical record review.¹³ Over 5.3 years of follow-up, allocation to active omega-3 FAs did not reduce the total number of participants who developed AMD compared with the placebo: 157 (1.2%) versus 167 (1.3%), respectively; HR 0.93; 95% CI 0.73 to 1.17.¹³ Finally, the Nutritional AMD Treatment-2 (NAT-2) study compared the efficacy of DHA-enriched supplementation for the prevention of wet AMD.¹² In total, 263 men and women aged 55–85 years with bilateral drusen or wet AMD in just one eye were randomised to a daily omega-3 FA supplement (containing 270mg EPA and 840mg DHA) or matching placebo.¹² Participants underwent baseline and annual fluorescein angiography examinations over a 3-year follow-up period.¹² There were no beneficial effects of omega-3 FAs on the primary efficacy outcome of time to the first occurrence of wet AMD: 19.5±10.9 months in the active group versus 18.7±10.6 months in the placebo group (HR 0.89; 95% CI 0.55 to 1.42).¹² However, only a per-protocol analysis was performed instead of intention-to-treat, which invalidates the underlying randomisation and may be prone to bias.

ASCEND-Eye has several strengths that facilitated a robust assessment of the effects of the study treatments on AMD, including its randomised design, well-defined exposure, a large number of participants and long duration of near-complete follow-up. Our study also has some limitations. Although the study average adherence was relatively good, poor adherence in the later years of follow-up, along with drop-ins to treatment in the placebo arm of the aspirin randomisation by those prescribed non-study

antiplatelet therapy, may have shifted risk estimates towards the null. Like the earlier studies, ASCEND-Eye was underpowered to detect modest but plausible effects of each treatment. The lack of power and lack of overall effect meant it was not appropriate to consider the effects of aspirin or omega-3 FAs in subgroups of important prognostic risk factors, such as age, gender or smoking.

AMD diagnoses that were eligible to be adjudicated were confirmed by medical record review, but confirmation by retinal photography remains the gold standard for detailed phenotyping of early and late disease, and wet and dry AMD subtypes. Since the AMD status of participants at baseline was unknown, we could not ascertain their disease progression or assess any possible treatment effects on disease progression. However, it seems unlikely that a treatment may have a large effect on progression and no effect on incidence.

The adjudication process had little effect on the categorisation of AMD events, which supported the decision not to adjudicate events derived from the VFQ. However, there was a difference in the psychometric performance of the in-trial and VFQ questionnaires. While the proportion of AMD events recorded on in-trial questionnaires was the same among those who did and did not receive the VFQ, the VFQ captured substantially more events than the in-trial questionnaires. This probably reflects the lack of any explicit questions about AMD on the ASCEND follow-up questionnaire. Therefore, the possibility of some under-reporting of events by participants who were ineligible to receive or did not respond to the VFQ cannot be excluded. These differences would further reduce the study's power, but they would not be expected to bias the results in favour of a particular treatment.

Next, although our findings are internally valid, their generalisability in a real-world setting may be limited by a lack of ethnic diversity in the ASCEND study population. Finally, although diabetes and AMD are not considered to be pathologically related to each other,¹⁷ the possibility cannot be excluded that the pharmacodynamic effects of each study treatment differ in those with diabetes compared with those without it.

In conclusion, we found no clinically-meaningful effects of taking aspirin or omega-3 FAs for the prevention of AMD. Our study overcomes some of the methodological limitations of previous observational studies, providing randomised evidence that could contribute to future meta-analyses.

Acknowledgements The authors acknowledge the contributions of the ASCEND-Eye Collaborative Group (see supplementary materials) and the generosity of trial participants.

Collaborators ASCEND-Eye Collaborative Group: Rory Collins, Richard Peto, Jill Barton, David Simpson, Amanda Adler, Theingi Aung, Colin Baigent, Jonathan Bodansky, Andrew Farmer, Richard Haynes, Roger McPherson, Marion Mafham, Nilesh Samani, Peter Sleight, Peter Sandercock, Richard Gray, Lucy Fletcher, Murphy Kevin, Susan Hurley, Ryonfa Lee, Sandra Pickworth, Monica Willett, Elizabeth Wincott, Michael Lay, Ola Murawska, Allen Young, Karen Melham, Gavin Brown, Jonathan Latham-Mollart, Anna Brewer, Peter Scanlon, Praveen Patel, Mike Olson, Jude Kay, Sanjiv Banerjee, Laura Evans, Alisha Griffiths, Mark Davies, Helen Clayton, Philip Kirby, Mike Pennington, Dominic Clarke, Jenny Anslow, Alex

Hallam, Jenny Witts, Sarah Egan, Alex Wharton, Arun Sachdev, Amy Derbyshire, Elizabeth Williamson, Katherine Hepplestone, Sujit Mithra, Su Oliver, Pauline Wiatrak-Olszewska, Tasso Gazis, Kelly Alvey, Eric Wu, Helen Cook, Nick Gregory, Priyai Parkinson, John Anderson, Louis Bolter, Prema Maharajan, Rachel McFee, Lynne Allsop, Donna Sowter, John Featonby, Richard Furnival, Helen Lipinski, Hazel Benjamin, Tanisha McAfee, Emma Payne, Liz Still.

Contributors JA and LB were co-principal investigators of the ASCEND trial, and together with ES, have made substantial contributions to the design of the ASCEND-Eye study. ES obtained regulatory and information governance approvals and wrote this manuscript. WS, GB, SP and IH gave statistical advice and made significant contributions to the data analysis plan and statistical analyses for ASCEND-Eye. All authors read the final manuscript. ES is responsible for the overall content as guarantor.

Funding The ASCEND study was funded by grants to the University of Oxford from the British Heart Foundation (SP/03/002, SP/08/010/25939, SP/14/3/31114, PG/05/013/18296). The supply of aspirin and matching placebo was provided by Bayer Healthcare, and the supply of omega-3 fatty acids and matching placebo capsules by Abbott Products Operations (formerly Solvay Pharmaceuticals), with some funding from each company to cover drug packaging. Some funding for the ASCEND-Eye study was obtained from the Macular Society. The Clinical Trial Research Unit at the University of Oxford receives support from the UK Medical Research Council (which funds the MRC Population Health Research Unit in a strategic partnership with the University of Oxford, MC_UU_00017/3, MC_UU_00017/5), the British Heart Foundation and Cancer Research UK. The Clinical Trial Service Unit & Epidemiological Studies Unit (CTSU) of the Nuffield Department of Population Health at the University of Oxford has a staff policy of not taking any personal payments directly or indirectly from industry (with reimbursement sought only for the costs of travel and accommodation to attend scientific meetings) <http://www.ctsu.ox.ac.uk/about-ctsu/documents/independent-research>. Financial sources had no input into the design of the study, nor the collection, analysis or interpretation of data, nor on the decision to submit manuscripts for publication.

Competing interests None declared.

Patient and public involvement statement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants and was approved by North West—Haydock Multicentre Research Ethics Committee, Reference: 03/8/087. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. All requests for data sharing should be addressed to JA and LB, co-principal investigators of ASCEND. These will be handled in line with the data access and sharing policy of the Nuffield Department of Population Health, University of Oxford (www.ndph.ox.ac.uk/about/data-access-policy).

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID ID

Emily Sammons <http://orcid.org/0000-0002-8647-0366>

REFERENCES

- Steinmetz JD, Bourne RRA, Briant PS, *et al.* Causes of blindness and vision impairment in 2020 and trends over 30 years, and prevalence of avoidable blindness in relation to VISION 2020: the Right to Sight: an analysis for the Global Burden of Disease Study. *Lancet Glob Health* 2021;9:e144–60.
- Liew G, Mitchell P, Wong TY, *et al.* The association of aspirin use with age-related macular degeneration. *JAMA Intern Med* 2013;173:258–64.
- Klein BEK, Howard KP, Gangnon RE, *et al.* Long-term Use of Aspirin and Age-Related Macular Degeneration. *JAMA* 2012;308:2469.
- Christen WG, Glynn RJ, Ajani UA, *et al.* Age-related maculopathy in a randomized trial of low-dose aspirin among US physicians. *Arch Ophthalmol* 2001;119:1143–9.
- Christen WG, Glynn RJ, Chew EY, *et al.* Low-dose aspirin and medical record-confirmed age-related macular degeneration in a randomized trial of women. *Ophthalmology* 2009;116:2386–92.
- Tan JSL, Wang JJ, Flood V, *et al.* Dietary fatty acids and the 10-year incidence of age-related macular degeneration: the Blue Mountains Eye Study. *Arch Ophthalmol* 2009;127:656–65.
- Karger AB, Guan W, Nomura SO, *et al.* ASSOCIATION OF PLASMA ω -3 FATTY ACIDS WITH EARLY AGE-RELATED MACULAR DEGENERATION IN THE MULTI-ETHNIC STUDY OF ATHEROSCLEROSIS. *Retina (Philadelphia, Pa)* 2022;42:1384–91.
- Agrón E, Mares J, Clemons TE, *et al.* Dietary Nutrient Intake and Progression to Late Age-Related Macular Degeneration in the Age-Related Eye Disease Studies 1 and 2. *Ophthalmology* 2021;128:425–42.
- Wu J, Cho E, Giovannucci EL, *et al.* Dietary Intakes of Eicosapentaenoic Acid and Docosahexaenoic Acid and Risk of Age-Related Macular Degeneration. *Ophthalmology* 2017;124:634–43.
- Cho E, Hung S, Willett WC, *et al.* Prospective study of dietary fat and the risk of age-related macular degeneration. *Am J Clin Nutr* 2001;73:209–18.
- Age-Related Eye Disease Research Group. Lutein + zeaxanthin and omega-3 fatty acids for age-related macular degeneration: the Age-Related Eye Disease Study 2 (AREDS2) randomized clinical trial. *J Am Med Assoc* 2013;309:2005–15.
- Souied EH, Delcourt C, Querques G, *et al.* Oral docosahexaenoic acid in the prevention of exudative age-related macular degeneration: the Nutritional AMD Treatment 2 study. *Ophthalmology* 2013;120:1619–31.
- Christen WG, Cook NR, Manson JE, *et al.* Effect of Vitamin D and ω -3 Fatty Acid Supplementation on Risk of Age-Related Macular Degeneration: An Ancillary Study of the VITAL Randomized Clinical Trial. *JAMA Ophthalmol* 2020;138:1280–9.
- Klein R, Peto T, Bird A, *et al.* The epidemiology of age-related macular degeneration. *Am J Ophthalmol* 2004;137:486–95.
- Bowman L, Mafham M, Stevens W, *et al.* ASCEND: A Study of Cardiovascular Events in Diabetes: Characteristics of a randomized trial of aspirin and of omega-3 fatty acid supplementation in 15,480 people with diabetes. *Am Heart J* 2018;198:135–44.
- Coleman HR, Chan C-C, Ferris FL III, *et al.* Age-related macular degeneration. *The Lancet* 2008;372:1835–45.
- Klein R, Klein BEK, Marino EK, *et al.* Early age-related maculopathy in the cardiovascular health study. *Ophthalmology* 2003;110:25–33.
- The Atherosclerosis Risk in Communities Study Investigators. Prevalence of Age-related Maculopathy in the Atherosclerosis Risk in Communities Study. *Arch Ophthalmol* 1999;117:1203.
- Sammons E, Bowman L, Stevens W, *et al.* ASCEND-Eye: Rationale, design and baseline characteristics for a sub-study of the ASCEND randomised trial, exploring the effects of aspirin and omega-3 fatty acids on diabetic retinopathy and age-related macular degeneration. *Contemp Clin Trials Commun* 2023;35:101184.
- The ASCEND Study Collaborative Group. Effects of Aspirin for Primary Prevention in Persons with Diabetes Mellitus. *N Engl J Med* 2018;379:1529–39.
- Group AC. Effects of n–3 Fatty Acid Supplements in Diabetes Mellitus. *N Engl J Med* 2018;379:1540–50.
- Mangione CM, Lee PP, Gutierrez PR, *et al.* Development of the 25-item National Eye Institute Visual Function Questionnaire. *Arch Ophthalmol* 2001;119:1050–8.
- Sammons EL, Buck G, Bowman LJ, *et al.* ASCEND-Eye: Effects of Aspirin on Diabetic Retinopathy. *Ophthalmology* 2024;131:771–9.
- Friedman DS, O'Colmain BJ, Muñoz B, *et al.* Prevalence of age-related macular degeneration in the United States. *Arch Ophthalmol* 2004;122:564–72.
- Fleiss JL, Tytun A, Ury HK. A simple approximation for calculating sample sizes for comparing independent proportions. *Biometrics* 1980;36:343–6.

- 26 Armitage P, Berry G, Matthews JNS. *Statistical Methods in Medical Research*. 4th edn. Blackwell Publishing, 2002.
- 27 Peto R, Pike MC, Armitage P, *et al*. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. II. analysis and examples. *Br J Cancer* 1977;35:1–39.
- 28 Yusuf S, Peto R, Lewis J, *et al*. Beta blockade during and after myocardial infarction: an overview of the randomized trials. *Prog Cardiovasc Dis* 1985;27:335–71.
- 29 Peto R, Pike MC, Armitage P, *et al*. Design and analysis of randomized clinical trials requiring prolonged observation of each patient. I. Introduction and design. *Br J Cancer* 1976;34:585–612.
- 30 Sapieha P, Stahl A, Chen J, *et al*. 5-Lipoxygenase metabolite 4-HDHA is a mediator of the antiangiogenic effect of ω -3 polyunsaturated fatty acids. *Sci Transl Med* 2011;3:69ra12.
- 31 Cohen J. A Coefficient of Agreement for Nominal Scales. *Educ Psychol Meas* 1960;20:37–46.
- 32 Snow KK, Seddon JM. Do age-related macular degeneration and cardiovascular disease share common antecedents? *Ophthalmic Epidemiol* 1999;6:125–43.
- 33 Christen WG, Chew EY. Does long-term aspirin use increase the risk of neovascular age-related macular degeneration? *Expert Opin Drug Saf* 2014;13:421–9.
- 34 Nowak JZ. Aspirin and age-related macular degeneration: positives versus negatives. *Expert Opin Drug Saf* 2014;13:687–90.
- 35 Chong EW, Guymer RH, Robman LD. Does aspirin increase the risk of age-related macular degeneration? *Expert Opin Drug Saf* 2014;13:691–3.
- 36 Seddon JM, George S, Rosner B, *et al*. Progression of age-related macular degeneration: prospective assessment of C-reactive protein, interleukin 6, and other cardiovascular biomarkers. *Arch Ophthalmol* 2005;123:774–82.
- 37 Anderson DH, Mullins RF, Hageman GS, *et al*. A role for local inflammation in the formation of drusen in the aging eye. *Am J Ophthalmol* 2002;134:411–31.
- 38 McNeil JJ, Nelson MR, Woods RL, *et al*. Effect of Aspirin on All-Cause Mortality in the Healthy Elderly. *N Engl J Med* 2018;379:1519–28.
- 39 McNeil JJ, Woods RL, Nelson MR, *et al*. Effect of Aspirin on Disability-free Survival in the Healthy Elderly. *N Engl J Med* 2018;379:1499–508.
- 40 McNeil JJ, Wolfe R, Woods RL, *et al*. Effect of Aspirin on Cardiovascular Events and Bleeding in the Healthy Elderly. *N Engl J Med* 2018;379:1509–18.
- 41 Robman LD, Wolfe R, Woods RL, *et al*. Effect of Low-Dose Aspirin on the Course of Age-Related Macular Degeneration: A Secondary Analysis of the ASPREE Randomized Clinical Trial. *JAMA Ophthalmol* 2024;142:627–35.
- 42 Ferris FL III, Wilkinson CP, Bird A, *et al*. Clinical Classification of Age-related Macular Degeneration. *Ophthalmology* 2013;120:844–51.
- 43 The Royal College of Ophthalmologists. National Ophthalmology Database Audit, Available: <https://www.nodaudit.org.uk> [Accessed 13 Mar 2023].
- 44 SanGiovanni JP, Chew EY. The role of omega-3 long-chain polyunsaturated fatty acids in health and disease of the retina. *Prog Retin Eye Res* 2005;24:87–138.
- 45 Joachim N, Mitchell P, Burlutsky G, *et al*. The Incidence and Progression of Age-Related Macular Degeneration over 15 Years: The Blue Mountains Eye Study. *Ophthalmology* 2015;122:2482–9.
- 46 Augood C, Chakravarthy U, Young I, *et al*. Oily fish consumption, dietary docosahexaenoic acid and eicosapentaenoic acid intakes, and associations with neovascular age-related macular degeneration. *Am J Clin Nutr* 2008;88:398–406.