

**Dig-In:**

## The Squeeze Theorem

*The Squeeze theorem allows us to exchange difficult functions for easy functions.*

In mathematics, sometimes we can study complex functions by exchanging them for simpler functions. The *Squeeze Theorem* tells us one situation where this is possible.

**Theorem 1** (Squeeze Theorem). *Suppose that*

$$g(x) \leq f(x) \leq h(x)$$

*for all  $x$  close to  $a$  but not necessarily equal to  $a$ . If*

$$\lim_{x \rightarrow a} g(x) = L = \lim_{x \rightarrow a} h(x),$$

*then  $\lim_{x \rightarrow a} f(x) = L$ .*

**Question 1** *I'm thinking of a function  $f$ . I know that for all  $x$*

$$0 \leq f(x) \leq x^2.$$

*What is  $\lim_{x \rightarrow 0} f(x)$ ?*

**Multiple Choice:**


- (a)  $f(x)$
- (b)  $f(0)$
- (c)  $0$  ✓
- (d) *impossible to say*

**Example 1.** *Consider the function*

$$f(x) = \begin{cases} \sqrt[5]{x} \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0, \\ 0 & \text{if } x = 0, \end{cases}$$

Learning outcomes: Understand the Squeeze Theorem and how it can be used to find limit values. Calculate limits using the Squeeze Theorem.

## The Squeeze Theorem



Is this function continuous at  $x = 0$ ?

**Explanation.** We must show that  $\lim_{x \rightarrow 0} f(x) = \boxed{0}$ . Note

$$-\sqrt[5]{x} \leq f(x) \leq \sqrt[5]{x}.$$

Since

$$\lim_{x \rightarrow 0} -\sqrt[5]{x} = \boxed{0} = \lim_{x \rightarrow 0} \sqrt[5]{x},$$


we see by the Squeeze Theorem, Theorem, that  $\lim_{x \rightarrow 0} f(x) = \boxed{0}$ . Hence  $f(x)$  is continuous.

Here we see how the informal definition of continuity being that you can “draw it” without “lifting your pencil” differs from the formal definition.

**Example 2.** Compute:

$$\lim_{\theta \rightarrow 0} \frac{\sin(\theta)}{\theta}$$

**Explanation.** To compute this limit, use the Squeeze Theorem. First note that we only need to examine  $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$  and for the present time, we’ll assume that  $\theta$  is positive. Consider the diagrams below:



From our diagrams above we see that

$$\text{Area of Triangle A} \leq \text{Area of Sector} \leq \text{Area of Triangle B}$$

and computing these areas we find

$$\frac{\cos(\theta) \sin(\theta)}{2} \leq \frac{\theta}{2} \leq \frac{\tan(\theta)}{2}.$$

## The Squeeze Theorem

Multiplying through by 2, and recalling that  $\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$  we obtain

$$\cos(\theta) \sin(\theta) \leq \theta \leq \frac{\sin(\theta)}{\cos(\theta)}.$$

Dividing through by  $\sin(\theta)$  and taking the reciprocals (reversing the inequalities), we find

$$\cos(\theta) \leq \frac{\sin(\theta)}{\theta} \leq \frac{1}{\cos(\theta)}.$$

Note,  $\cos(-\theta) = \cos(\theta)$  and  $\frac{\sin(-\theta)}{-\theta} = \frac{\sin(\theta)}{\theta}$ , so these inequalities hold for all  $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ . Additionally, we know

$$\lim_{\theta \rightarrow 0} \cos(\theta) = \boxed{1}_{\text{given}} = \lim_{\theta \rightarrow 0} \frac{1}{\cos(\theta)},$$


and so we conclude by the Squeeze Theorem,  $\lim_{\theta \rightarrow 0} \frac{\sin(\theta)}{\theta} = \boxed{1}_{\text{given}}.$

When solving a problem with the Squeeze Theorem, one must write a sort of mathematical poem. You have to tell your friendly reader exactly which functions you are using to “squeeze-out” your limit.

**Example 3.** Compute:

$$\lim_{x \rightarrow 0} \left( \sin(x) e^{\cos(\frac{1}{x^3})} \right)$$

**Explanation.** Let's graph this function to see what's going on:



The function  $\sin(x) e^{\cos(\frac{1}{x^3})}$  has two factors:

goes to zero as  $x \rightarrow 0$

$$\overbrace{\sin(x)} \cdot \underbrace{e^{\cos(\frac{1}{x^3})}}$$

bounded between  $e^{-1}$  and  $e$

## The Squeeze Theorem

Hence we have that when  $x > 0$

$$\sin(x) \boxed[e^{-1}]{\text{given}} \leq \sin(x) e^{\cos(\frac{1}{x^3})} \leq \sin(x) \boxed[e]{\text{given}}$$

and we see

$$\lim_{x \rightarrow 0^+} \sin(x) \boxed[e^{-1}]{\text{given}} = \boxed[0]{\text{given}} = \lim_{x \rightarrow 0^+} \sin(x) \boxed[e]{\text{given}}$$

and so by the Squeeze theorem,

$$\lim_{x \rightarrow 0^+} \left( \sin(x) e^{\cos(\frac{1}{x^3})} \right) = \boxed[0]{\text{given}}.$$

In a similar fashion, when  $x < 0$ ,

$$\sin(x) \boxed[e]{\text{given}} \leq \sin(x) e^{\cos(\frac{1}{x^3})} \leq \sin(x) \boxed[e^{-1}]{\text{given}}$$

and so

$$\lim_{x \rightarrow 0^-} \sin(x) \boxed[e]{\text{given}} = \boxed[0]{\text{given}} = \lim_{x \rightarrow 0^-} \sin(x) \boxed[e^{-1}]{\text{given}},$$

and again by the Squeeze Theorem  $\lim_{x \rightarrow 0^-} \left( \sin(x) e^{\cos(\frac{1}{x^3})} \right) = 0$ . Hence we see that

$$\lim_{x \rightarrow 0} \left( \sin(x) e^{\cos(\frac{1}{x^3})} \right) = \boxed[0]{\text{given}}.$$