


Intact Brain Network Function in a COVID-19 Patient with Persistent Unresponsiveness


David Fischer, M.D., Zachary D. Threlkeld, M.D., Yelena G. Bodien, Ph.D.,
John Kirsch, Ph.D., Susie Y. Huang, M.D., Ph.D., Pamela W. Schaefer, M.D., Otto Rapalino, M.D.,
Leigh R. Hochberg, M.D., Ph.D., Bruce R. Rosen, M.D., Ph.D., and Brian L. Edlow, M.D.

Resting-state fMRI Sequence Parameters



Resting-state fMRI Sequence Parameters

Parameter	Value
sequence type	FID-EPI
receiver coil	32 channel head coil
repetition time, TR	1250 ms
echo time, TE	30 ms
flip angle, FA	65 degs
# slices	72
interslice gap	0% (contiguous)
Fat suppression	yes
sampling bandwidth	1684 Hz/pixel
slice thickness, TH	2 mm
number of acquisition, NEX	1
slice orientation	Transverse, axial
phase encoding direction	A-P
Field-of-view readout, FOV(r)	212 mm
Field-of-view phase, FOV(p)	212 mm
matrix readout, MA(r)	106
matrix phase, MA(p)	106
partial fourier (phase)	87.5% (7/8)
readout acceleration	GRAPPA, 2x
slice acceleration	4x
# measurements (time points)	482





https://github.com/ComaRecoveryLab/COVID-19_rsfMRI

www.ComaRecoveryLab.org

 @ComaRecoveryLab

