
Oxygenation Severity in Those with both PF and SF
SpO2/FiO2 Defined
PaO2/FiO2 Defined
161 (3.5%)
489 (10.6%)
1221 (26.5%)
2744 (59.5%)
633 (13.7%)
934 (20.2%)
1811 (39.2%)
1237 (26.8%)
Not Hypoxemic
Mild Hypoxemia
Moderate Hypoxemia
Severe Hypoxemia
Prevalence
Higher
Lower


Death or Hospice by Day 28, (n=7943)
SpO2/FiO2 Defined
PaO2/FiO2 Defined
12.8%
15.7%
17.0%
26.1%
29.1%
33.0%
32.8%
40.8%
Not Hypoxemic
Mild Hypoxemia
Moderate Hypoxemia
Severe Hypoxemia
Higher
Lower


Death or Hospice by Day 28, (n=7943)
SpO2/FiO2 Defined
PaO2/FiO2 Defined
13.1%
14.3%
18.1%
19.7%
25.4%
29.1%
33.6%
32.1%
35.0%
43.3%
0
1
2
3
4
SOFA-2 Respiratory Score
Higher
Lower


Death or Hospice by Day 28, (n=4615),
Both PF and SF Measured
SpO2/FiO2 Defined
PaO2/FiO2 Defined
19.3%
27.6%
30.5%
38.4%
30.5%
33.3%
32.3%
40.7%
Not Hypoxemic
Mild Hypoxemia
Moderate Hypoxemia
Severe Hypoxemia
Higher
Lower


Death or Hospice by Day 28, (n=4615),
Both PF and SF Measured
SpO2/FiO2 Defined
PaO2/FiO2 Defined
22.7%
28.6%
28.8%
29.8%
39.5%
30.5%
34.5%
31.5%
34.7%
43.5%
0
1
2
3
4
SOFA-2 Respiratory Score
Higher
Lower


Death or Hospice by Day 28, (n=4615),
PF and SF Measured Contemparaneously
SpO2/FiO2 Defined
PaO2/FiO2 Defined
20.2%
27.3%
30.5%
38.4%
30.5%
33.3%
32.3%
40.7%
Not Hypoxemic
Mild Hypoxemia
Moderate Hypoxemia
Severe Hypoxemia
Higher
Lower


Death or Hospice by Day 28, (n=4615),
PF and SF Measured Contemparaneously
SpO2/FiO2 Defined
PaO2/FiO2 Defined
23.4%
28.6%
28.6%
29.7%
39.5%
30.5%
34.5%
31.5%
34.7%
43.5%
0
1
2
3
4
SOFA-2 Respiratory Score
Higher
Lower