

Randomized Algorithms for Max-SAT

Given a graph $G(V, E)$, find a cut A, B with maximum weight
 $w(A, B) = |(A \times B) \cap E|$.

Given a graph $G(V, E)$, find a cut A, B with maximum weight $w(A, B) = |(A \times B) \cap E|$.

Generalizations to weighted cases straightforward and will not be considered in the following.

Given a graph $G(V, E)$, find a cut A, B with maximum weight $w(A, B) = |(A \times B) \cap E|$.

Generalizations to weighted cases straightforward and will not be considered in the following.

Min-Cut is solvable in polynomial time, Max-Cut is NP-hard.

A Simple Randomized Algorithm

Dumb Rounding

1. For every node v flip a fair coin.
2. If heads, place v in A .
3. If tails, place v in B .

A Simple Randomized Algorithm

Dumb Rounding

1. For every node v flip a fair coin.
2. If heads, place v in A .
3. If tails, place v in B .

Theorem

Dumb rounding is a $\frac{1}{2}$ -approximation.

A Simple Randomized Algorithm

Dumb Rounding

1. For every node v flip a fair coin.
2. If heads, place v in A .
3. If tails, place v in B .

Theorem

Dumb rounding is a $\frac{1}{2}$ -approximation.

Corollary

The Max-Cut always contains at least $\frac{|E|}{2}$ edges.

(We'll only prove the theorem.)

Analysis of Dumb Rounding

Theorem

Dumb rounding is a $\frac{1}{2}$ -approximation.

Consider an arbitrary edge $e = (u, v)$. We have the following four cases:

- $u \in A, v \in A \Rightarrow e \notin \text{cut}$
- $u \in A, v \in B \Rightarrow e \in \text{cut}$
- $u \in B, v \in A \Rightarrow e \in \text{cut}$
- $u \in B, v \in B \Rightarrow e \notin \text{cut}$

Analysis of Dumb Rounding

Theorem

Dumb rounding is a $\frac{1}{2}$ -approximation.

Consider an arbitrary edge $e = (u, v)$. We have the following four cases:

- $u \in A, v \in A \Rightarrow e \notin \text{cut}$
- $u \in A, v \in B \Rightarrow e \in \text{cut}$
- $u \in B, v \in A \Rightarrow e \in \text{cut}$
- $u \in B, v \in B \Rightarrow e \notin \text{cut}$

All cases occur with equal probability. Hence

$$\mathbb{E}[|(A \times B) \cap E|] = \sum_{e \in E} \mathbb{E}[e \in \text{cut}]$$

Analysis of Dumb Rounding

Theorem

Dumb rounding is a $\frac{1}{2}$ -approximation.

Consider an arbitrary edge $e = (u, v)$. We have the following four cases:

- $u \in A, v \in A \Rightarrow e \notin \text{cut}$
- $u \in A, v \in B \Rightarrow e \in \text{cut}$
- $u \in B, v \in A \Rightarrow e \in \text{cut}$
- $u \in B, v \in B \Rightarrow e \notin \text{cut}$

All cases occur with equal probability. Hence

$$\mathbb{E}[|(A \times B) \cap E|] = \sum_{e \in E} \mathbb{E}[e \in \text{cut}] = \sum_{e \in E} \mathbb{P}[e \in \text{cut}]$$

Analysis of Dumb Rounding

Theorem

Dumb rounding is a $\frac{1}{2}$ -approximation.

Consider an arbitrary edge $e = (u, v)$. We have the following four cases:

- $u \in A, v \in A \Rightarrow e \notin \text{cut}$
- $u \in A, v \in B \Rightarrow e \in \text{cut}$
- $u \in B, v \in A \Rightarrow e \in \text{cut}$
- $u \in B, v \in B \Rightarrow e \notin \text{cut}$

All cases occur with equal probability. Hence

$$\begin{aligned}\mathbb{E}[|(A \times B) \cap E|] &= \sum_{e \in E} \mathbb{E}[e \in \text{cut}] = \sum_{e \in E} \mathbb{P}[e \in \text{cut}] \\ &= \sum_{e \in E} \frac{1}{2} = \frac{|E|}{2}\end{aligned}$$

Analysis of Dumb Rounding

Theorem

Dumb rounding is a $\frac{1}{2}$ -approximation.

Consider an arbitrary edge $e = (u, v)$. We have the following four cases:

- $u \in A, v \in A \Rightarrow e \notin \text{cut}$
- $u \in A, v \in B \Rightarrow e \in \text{cut}$
- $u \in B, v \in A \Rightarrow e \in \text{cut}$
- $u \in B, v \in B \Rightarrow e \notin \text{cut}$

All cases occur with equal probability. Hence

$$\begin{aligned}\mathbb{E}[|(A \times B) \cap E|] &= \sum_{e \in E} \mathbb{E}[e \in \text{cut}] = \sum_{e \in E} \mathbb{P}[e \in \text{cut}] \\ &= \sum_{e \in E} \frac{1}{2} = \frac{|E|}{2} \geq \frac{OPT}{2}\end{aligned}$$

The output is only a 2-approximation on expectation. How can we be sure?

The output is only a 2-approximation on expectation. How can we be sure?

First Idea: Repeat the algorithm a few times and take the best one. $\log n$ repetitions yield a 2 approximation with probability $1 - 1/n$.

The output is only a 2-approximation on expectation. How can we be sure?

First Idea: Repeat the algorithm a few times and take the best one. $\log n$ repetitions yield a 2 approximation with probability $1 - 1/n$.

Remove the randomness from the algorithm.

We will use the method of conditional expectations. Let X be a function of Bernoulli random variables X_1, \dots, X_n . Then

$$\mathbb{E}[X] = \mathbb{E}[X|X_1 = 0] \cdot \mathbb{P}[X_1 = 0] + \mathbb{E}[X|X_1 = 1] \cdot \mathbb{P}[X_1 = 1].$$

We will use the method of conditional expectations. Let X be a function of Bernoulli random variables X_1, \dots, X_n . Then

$$\mathbb{E}[X] = \mathbb{E}[X|X_1 = 0] \cdot \mathbb{P}[X_1 = 0] + \mathbb{E}[X|X_1 = 1] \cdot \mathbb{P}[X_1 = 1].$$

This implies $\max(\mathbb{E}[X|X_1 = 0], \mathbb{E}[X|X_1 = 1]) > \mathbb{E}[X]$.

Derandomization

We will use the method of conditional expectations. Let X be a function of Bernoulli random variables X_1, \dots, X_n . Then

$$\mathbb{E}[X] = \mathbb{E}[X|X_1 = 0] \cdot \mathbb{P}[X_1 = 0] + \mathbb{E}[X|X_1 = 1] \cdot \mathbb{P}[X_1 = 1].$$

This implies $\max(\mathbb{E}[X|X_1 = 0], \mathbb{E}[X|X_1 = 1]) > \mathbb{E}[X]$.

Suppose without loss of generality that

$\max(\mathbb{E}[X|X_1 = 0], \mathbb{E}[X|X_1 = 1]) = \mathbb{E}[X|X_1 = 0]$. By the same argument

$$\max(\mathbb{E}[X|X_1 = 0, X_2 = 0], \mathbb{E}[X|X_1 = 0, X_2 = 1]) > \mathbb{E}[X].$$

Evaluation of the Expectation

Denote by X_i the random coin toss of node v_i . Let X_1, \dots, X_{i-1} be the fixed coin tosses (i.e. the nodes that were already placed in A or B). Let $m_1 = |(A \times \{v_i\}) \cap E|$ and $m_2 = |(B \times \{v_i\}) \cap E|$ and $k = |(A \times B) \cap E|$ and $\ell = |((A \times A) \cup (B \times B)) \cap E|$.

Evaluation of the Expectation

Denote by X_i the random coin toss of node v_i . Let X_1, \dots, X_{i-1} be the fixed coin tosses (i.e. the nodes that were already placed in A or B). Let $m_1 = |(A \times \{v_i\}) \cap E|$ and $m_2 = |(B \times \{v_i\}) \cap E|$ and $k = |(A \times B) \cap E|$ and $\ell = |((A \times A) \cup (B \times B)) \cap E|$.

Then

$$\mathbb{E}[X | X_1, \dots, X_{i-1} \text{ fixed}, X_i = 0] = k + m_1 + \frac{|E| - k - m_1 - m_2 - \ell}{2}$$

$$\mathbb{E}[X | X_1, \dots, X_{i-1} \text{ fixed}, X_i = 1] = k + m_2 + \frac{|E| - k - m_1 - m_2 - \ell}{2}$$

Evaluation of the Expectation

Denote by X_i the random coin toss of node v_i . Let X_1, \dots, X_{i-1} be the fixed coin tosses (i.e. the nodes that were already placed in A or B). Let $m_1 = |(A \times \{v_i\}) \cap E|$ and $m_2 = |(B \times \{v_i\}) \cap E|$ and $k = |(A \times B) \cap E|$ and $\ell = |((A \times A) \cup (B \times B)) \cap E|$.

Then

$$\mathbb{E}[X | X_1, \dots, X_{i-1} \text{ fixed}, X_i = 0] = k + m_1 + \frac{|E| - k - m_1 - m_2 - \ell}{2}$$

$$\mathbb{E}[X | X_1, \dots, X_{i-1} \text{ fixed}, X_i = 1] = k + m_2 + \frac{|E| - k - m_1 - m_2 - \ell}{2}$$

Greedy Max-Cut (via Derandomization)

1. $A \leftarrow \{v_1\}, B \leftarrow \emptyset$
2. For $i = 2$ to n
 place v_i into the set that maximizes the current cut size.

Maximum Satisfiability

Given a boolean formula in conjunctive normal form, satisfy as many clauses as possible.

- Formula $(x_1 \vee \overline{x_2} \vee \overline{x_3}) \wedge (x_2 \vee \overline{x_3} \vee x_4 \vee x_5) \wedge \overline{x_5} \wedge (w_3 \vee x_5) \dots$
- Clause $(x_1 \vee \overline{x_2} \vee \overline{x_3})$
- Literal $x_1, \overline{x_2}$

Maximum Satisfiability

Given a boolean formula in conjunctive normal form, satisfy as many clauses as possible.

- Formula $(x_1 \vee \overline{x_2} \vee \overline{x_3}) \wedge (x_2 \vee \overline{x_3} \vee x_4 \vee x_5) \wedge \overline{x_5} \wedge (w_3 \vee x_5) \dots$
- Clause $(x_1 \vee \overline{x_2} \vee \overline{x_3})$
- Literal $x_1, \overline{x_2}$

Max 2-SAT (every clause has exactly two literals) is a special case of the Max-Cut problem.

Integer Linear Program Formulation

- For each clause $(x_1 \vee x_2 \vee \overline{x_3})$ add a variable z
- For each literal x add a binary variable y .
- c is satisfied if and only if $y_1 + y_2 + (1 - y_3) \geq 1$.

Integer Linear Program Formulation

- For each clause $(x_1 \vee x_2 \vee \overline{x}_3)$ add a variable z
- For each literal x add a binary variable y .
- c is satisfied if and only if $y_1 + y_2 + (1 - y_3) \geq 1$.

$$\begin{aligned} & \text{maximize } \sum_{\text{clause } c} w_c \cdot z_c && \text{such that} \\ & \sum_{x_i \in c} y_i + \sum_{\overline{x}_i \in c} (1 - y_i) \geq z_c && \text{for all clauses } c \\ & y_i \in \{0, 1\} \\ & 0 \leq z_i \leq 1 \end{aligned}$$

Algorithm

$$\begin{array}{ll} \text{maximize} & \sum_{\text{clause } c} w_c \cdot z_c \\ & \text{such that} \\ & \sum_{x_i \in c} y_i + \sum_{\bar{x}_i \in c} (1 - y_i) \geq z_c \\ & \text{for all clauses } c \end{array}$$

$$y_i \in \{0, 1\}$$

$$0 \leq z_i \leq 1$$

Randomized Rounding

1. Solve LP relaxation ($0 \leq y_i \leq 1$)
2. Set each x_i to 1 with probability y_i

Algorithm

$$\text{maximize } \sum_{\text{clause } c} w_c \cdot z_c \quad \text{such that}$$

$$\sum_{x_i \in c} y_i + \sum_{\bar{x}_i \in c} (1 - y_i) \geq z_c \quad \text{for all clauses } c$$

$$\text{(simplify notation...)} \quad \sum_{x_i^* \in c} y_i^* \geq z_c \quad \text{for all clauses } c$$

$$y_i \in \{0, 1\}$$

$$0 \leq z_i \leq 1$$

$$y_i^* = \begin{cases} y_i & \text{if } x_i \in c \\ 1 - y_i & \text{if } \bar{x}_i \in c \end{cases}$$

Randomized Rounding

1. Solve LP relaxation ($0 \leq y_i \leq 1$)
2. Set each x_i to 1 with probability y_i

A Useful Lemma

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

Proof: By induction over n .

A Useful Lemma

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

Proof: By induction over n . For $n = 1$ the statement is trivial

A Useful Lemma

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

Proof: By induction over n . For $n = 1$ the statement is trivial

Suppose the statement holds for any n non-negative numbers

A Useful Lemma

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

Proof: By induction over n . For $n = 1$ the statement is trivial

Suppose the statement holds for any n non-negative numbers

If all $n + 1$ numbers are equal the two means are equal

A Useful Lemma

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$\frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

Proof: By induction over n . For $n = 1$ the statement is trivial

Suppose the statement holds for any n non-negative numbers

If all $n + 1$ numbers are equal the two means are equal

If not all numbers are equal, at least one number is greater than the arithmetic mean and one number is smaller than the arithmetic mean.

Without loss of generality assume that $x_{n+1} < a := \frac{x_1 + x_2 + \dots + x_{n+1}}{n+1} < x_1$

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$a = \frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$a = \frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

$$\Rightarrow 0 < (a - x_{n+1})(x_1 - a)$$

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$a = \frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

$$\Rightarrow 0 < (a - x_{n+1})(x_1 - a)$$

$$(n+1) \cdot a = x_1 + x_2 + \dots + x_{n+1} \Leftrightarrow n \cdot a = x_2 + \dots + x_1 + x_{n+1} - a$$

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$a = \frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

$$\Rightarrow 0 < (a - x_{n+1})(x_1 - a)$$

$(n+1) \cdot a = x_1 + x_2 + \dots + x_{n+1} \Leftrightarrow n \cdot a = x_2 + \dots + x_1 + x_{n+1} - a$, i.e. a is also the arithmetic mean of the numbers $\{x_2, \dots, x_n, x_1 + x_{n+1} - a\}$

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$a = \frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

$$\Rightarrow 0 < (a - x_{n+1})(x_1 - a)$$

$(n+1) \cdot a = x_1 + x_2 + \dots + x_{n+1} \Leftrightarrow n \cdot a = x_2 + \dots + x_1 + x_{n+1} - a$, i.e. a is also the arithmetic mean of the numbers $\{x_2, \dots, x_n, x_1 + x_{n+1} - a\}$

$$(x_1 + x_{n+1} - a)a - x_1 x_{n+1} = (x_1 - a)(a - x_{n+1}) > 0$$

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$a = \frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

$$\Rightarrow 0 < (a - x_{n+1})(x_1 - a)$$

$(n+1) \cdot a = x_1 + x_2 + \dots + x_{n+1} \Leftrightarrow n \cdot a = x_2 + \dots + x_1 + x_{n+1} - a$, i.e. a is also the arithmetic mean of the numbers $\{x_2, \dots, x_n, x_1 + x_{n+1} - a\}$

$$(x_1 + x_{n+1} - a)a - x_1 x_{n+1} = (x_1 - a)(a - x_{n+1}) > 0 \text{ which implies} \\ (x_1 + x_{n+1} - a)a > x_1 x_{n+1}$$

We apply the inductive hypothesis on $\{x_2, \dots, x_n, x_1 + x_{n+1} - a\}$
 $a^{n+1} \geq x_2 \cdot x_3 \cdot \dots \cdot x_n \cdot (x_1 + x_{n+1} - a)a$

AM-GM inequality

Let y_1, \dots, y_n be a list of n nonnegative numbers. Then

$$a = \frac{x_1 + x_2 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdot x_2 \cdot \dots \cdot x_n}$$

$$\Rightarrow 0 < (a - x_{n+1})(x_1 - a)$$

$(n+1) \cdot a = x_1 + x_2 + \dots + x_{n+1} \Leftrightarrow n \cdot a = x_2 + \dots + x_1 + x_{n+1} - a$, i.e. a is also the arithmetic mean of the numbers $\{x_2, \dots, x_n, x_1 + x_{n+1} - a\}$

$$(x_1 + x_{n+1} - a)a - x_1 x_{n+1} = (x_1 - a)(a - x_{n+1}) > 0 \text{ which implies} \\ (x_1 + x_{n+1} - a)a > x_1 x_{n+1}$$

We apply the inductive hypothesis on $\{x_2, \dots, x_n, x_1 + x_{n+1} - a\}$


$$a^{n+1} \geq x_2 \cdot x_3 \cdot \dots \cdot x_n \cdot (x_1 + x_{n+1} - a)a \geq x_2 \cdot x_3 \cdot \dots \cdot x_1 x_{n+1}$$

Concave Functions

Definition

A function f is concave on $[l, r]$ if for any $0 \leq \alpha \leq 1$

$$f((1 - \alpha) \cdot l + \alpha \cdot r) \geq (1 - \alpha) \cdot f(l) + \alpha \cdot f(r)$$



We will use the special case $l = 0$, $r = 1$ and $f(0) = 0$:

$$f(\alpha \cdot r) \geq \alpha \cdot f(1)$$

Analysis of Randomized Rounding

Consider any clause c with k literals. The associated constraint is $\sum_{x_i \in c} y_i + \sum_{\bar{x}_i \in c} (1 - y_i) = \sum_{x_i^* \in c} y_i^* \geq z_c$.

$$\mathbb{P}[c \text{ is satisfied}] = 1 - \prod_{i=1}^k (1 - y_i^*)$$

Analysis of Randomized Rounding

Consider any clause c with k literals. The associated constraint is $\sum_{x_i \in c} y_i + \sum_{\bar{x}_i \in c} (1 - y_i) = \sum_{x_i^* \in c} y_i^* \geq z_c$.

$$\mathbb{P}[c \text{ is satisfied}] = 1 - \prod_{i=1}^k (1 - y_i^*)$$

$$\text{AM-GM-inequality} \geq 1 - \left(\frac{k - \sum_{i=1}^k y_i^*}{k} \right)^k$$

Analysis of Randomized Rounding

Consider any clause c with k literals. The associated constraint is $\sum_{x_i \in c} y_i + \sum_{\bar{x}_i \in c} (1 - y_i) = \sum_{x_i^* \in c} y_i^* \geq z_c$.

$$\mathbb{P}[c \text{ is satisfied}] = 1 - \prod_{i=1}^k (1 - y_i^*)$$

$$\text{AM-GM-inequality} \geq 1 - \left(\frac{k - \sum_{i=1}^k y_i^*}{k} \right)^k$$

$$\text{LP-constraint} \geq 1 - \left(1 - \frac{z_c}{k} \right)^k$$

Analysis of Randomized Rounding

Consider any clause c with k literals. The associated constraint is $\sum_{x_i \in c} y_i + \sum_{\bar{x}_i \in c} (1 - y_i) = \sum_{x_i^* \in c} y_i^* \geq z_c$.

$$\mathbb{P}[c \text{ is satisfied}] = 1 - \prod_{i=1}^k (1 - y_i^*)$$

$$\text{AM-GM-inequality} \geq 1 - \left(\frac{k - \sum_{i=1}^k y_i^*}{k} \right)^k$$

$$\text{LP-constraint} \geq 1 - \left(1 - \frac{z_c}{k} \right)^k$$

$$\text{concave function } (\alpha \equiv z_c) \geq \left(1 - \left(1 - \frac{1}{k} \right)^k \right) z_c$$

For any clause, we have $\mathbb{P}[c \text{ is satisfied}] \geq \left(1 - \left(1 - \frac{1}{k}\right)^k\right) z_c$.

$$\mathbb{E}\left[\sum_{\text{clause } c} w_c \cdot z_c\right] = \sum_{\text{clause } c} w_c \cdot \mathbb{P}[c \text{ is satisfied}]$$

For any clause, we have $\mathbb{P}[c \text{ is satisfied}] \geq \left(1 - \left(1 - \frac{1}{k}\right)^k\right) z_c$.

$$\begin{aligned}\mathbb{E}\left[\sum_{\text{clause } c} w_c \cdot z_c\right] &= \sum_{\text{clause } c} w_c \cdot \mathbb{P}[c \text{ is satisfied}] \\ &\geq \min_k \left(1 - \left(1 - \frac{1}{k}\right)^k\right) \sum_{\text{clause } c} w_c z_c\end{aligned}$$

For any clause, we have $\mathbb{P}[c \text{ is satisfied}] \geq \left(1 - \left(1 - \frac{1}{k}\right)^k\right) z_c$.

$$\begin{aligned}\mathbb{E}\left[\sum_{\text{clause } c} w_c \cdot z_c\right] &= \sum_{\text{clause } c} w_c \cdot \mathbb{P}[c \text{ is satisfied}] \\ &\geq \min_k \left(1 - \left(1 - \frac{1}{k}\right)^k\right) \sum_{\text{clause } c} w_c z_c \\ &\geq \min_k \left(1 - \left(1 - \frac{1}{k}\right)^k\right) OPT\end{aligned}$$

For any clause, we have $\mathbb{P}[c \text{ is satisfied}] \geq \left(1 - \left(1 - \frac{1}{k}\right)^k\right) z_c$.

$$\begin{aligned}\mathbb{E}\left[\sum_{\text{clause } c} w_c \cdot z_c\right] &= \sum_{\text{clause } c} w_c \cdot \mathbb{P}[c \text{ is satisfied}] \\ &\geq \min_k \left(1 - \left(1 - \frac{1}{k}\right)^k\right) \sum_{\text{clause } c} w_c z_c \\ &\geq \min_k \left(1 - \left(1 - \frac{1}{k}\right)^k\right) OPT \\ &\geq \left(1 - \frac{1}{e}\right) OPT\end{aligned}$$

Dumb Rounding

1. Set literals to **true** with probability $1/2$

For a clause with k literals, the probability of being satisfied is

$$\mathbb{P}[c \text{ is satisfied}] \geq \left(1 - \left(\frac{1}{2}\right)^k\right).$$

Further Improvements

Dumb Rounding

1. Set literals to **true** with probability $1/2$

For a clause with k literals, the probability of being satisfied is $\mathbb{P}[c \text{ is satisfied}] \geq \left(1 - \left(\frac{1}{2}\right)^k\right)$.

Combined Rounding

1. Run Dumb Rounding $\rightarrow W_1$
2. Run Randomized Rounding $\rightarrow W_2$
3. Output the better of the two

$$\mathbb{E}[\max(W_1, W_2)] \geq \mathbb{E}\left[\frac{1}{2}W_1 + \frac{1}{2}W_2\right]$$

$$\begin{aligned}\mathbb{E}[\max(W_1, W_2)] &\geq \mathbb{E}\left[\frac{1}{2}W_1 + \frac{1}{2}W_2\right] \\ \text{Linearity} &\geq \sum_{\text{clause } c} w_c \cdot \left(\frac{1}{2}\mathbb{P}[c \text{ is satisfied by Dumb}] \right. \\ &\quad \left. + \frac{1}{2}\mathbb{P}[c \text{ is satisfied by Random}]\right)\end{aligned}$$

$$\begin{aligned}\mathbb{E}[\max(W_1, W_2)] &\geq \mathbb{E}\left[\frac{1}{2}W_1 + \frac{1}{2}W_2\right] \\ \text{Linearity} &\geq \sum_{\text{clause } c} w_c \cdot \left(\frac{1}{2}\mathbb{P}[c \text{ is satisfied by Dumb}] \right. \\ &\quad \left. + \frac{1}{2}\mathbb{P}[c \text{ is satisfied by Random}]\right) \\ &\geq \sum_{\text{clause } c} w_c \cdot \frac{1}{2} \left(1 - \left(\frac{1}{2}\right)^k\right) + \frac{1}{2} \left(1 - \left(1 - \frac{1}{k}\right)^k\right) z_c\end{aligned}$$

$$\mathbb{E}[\max(W_1, W_2)] \geq \mathbb{E}\left[\frac{1}{2}W_1 + \frac{1}{2}W_2\right]$$

$$\begin{aligned} \text{Linearity} &\geq \sum_{\text{clause } c} w_c \cdot \left(\frac{1}{2} \mathbb{P}[c \text{ is satisfied by Dumb}] \right. \\ &\quad \left. + \frac{1}{2} \mathbb{P}[c \text{ is satisfied by Random}] \right) \end{aligned}$$

$$\geq \sum_{\text{clause } c} w_c \cdot \frac{1}{2} \left(1 - \left(\frac{1}{2} \right)^k \right) + \frac{1}{2} \left(1 - \left(1 - \frac{1}{k} \right)^k \right) z_c$$

$$\begin{aligned} \text{boring calculations} &\geq \sum_{\text{clause } c} w_c \cdot \frac{3}{4} z_j \end{aligned}$$

$$\mathbb{E}[\max(W_1, W_2)] \geq \mathbb{E}\left[\frac{1}{2}W_1 + \frac{1}{2}W_2\right]$$

$$\begin{aligned} \text{Linearity} &\geq \sum_{\text{clause } c} w_c \cdot \left(\frac{1}{2} \mathbb{P}[c \text{ is satisfied by Dumb}] \right. \\ &\quad \left. + \frac{1}{2} \mathbb{P}[c \text{ is satisfied by Random}] \right) \end{aligned}$$

$$\geq \sum_{\text{clause } c} w_c \cdot \frac{1}{2} \left(1 - \left(\frac{1}{2} \right)^k \right) + \frac{1}{2} \left(1 - \left(1 - \frac{1}{k} \right)^k \right) z_c$$

$$\text{boring calculations} \geq \sum_{\text{clause } c} w_c \cdot \frac{3}{4} z_j$$

$$\geq \frac{3}{4} \sum_{\text{clause } c} w_c z_j$$

$$\geq \frac{3}{4} OPT$$

Boring Calculations

Concavity of $1 - \left(1 - \frac{x}{k}\right)^k$ in $[0, 1]$:

First Derivative: $\left(1 - \frac{x}{k}\right)^{k-1}$

Second Derivative: $-\frac{k-1}{k} \left(1 - \frac{x}{k}\right)^{k-2} \leq 0$ (for $k \geq 1$)

For $k = 1$: $\frac{1}{2}\left(1 - \frac{1}{2}\right) + \frac{1}{2}z_c = \frac{1}{4} + \frac{1}{2}z_c \geq \frac{3}{4}z_c$

For $k = 2$: $\frac{1}{2}\left(1 - \frac{1}{4}\right) + \frac{1}{2}\left(1 - \left(1 - \frac{1}{2}\right)^2\right)z_c \geq \frac{3}{8} + \frac{3}{8}z_c \geq \frac{3}{4}z_c$

For $k \geq 3$: $\frac{1}{2}\left(1 - \frac{1}{8}\right) + \frac{1}{2}\left(1 - \frac{1}{e}\right)z_c \geq \frac{7}{16} + \frac{1}{2}(1 - 1/e)z_c \geq 0.753z_c$