

and the leakage current, V_i are the equilibrium potentials for each of the currents, and α_j and β_j are nonlinear functions of V . We use the notation n , m , and h for the gate functions because the notation is universally used in the literature. These gate functions are empirical attempts to describe how the membrane controls the flow of ions into and out of the nerve cell. Hodgkin and Huxley found the following empirical forms for α_j and β_j :

$$\alpha_n = 0.01(V + 10)/[e^{(1+V/10)} - 1] \quad (4.31a)$$

$$\beta_n = 0.125 e^{V/80} \quad (4.31b)$$

$$\alpha_m = 0.01(V + 25)/[e^{(2.5+V/10)} - 1] \quad (4.31c)$$

$$\beta_m = 4 e^{V/18} \quad (4.31d)$$

$$\alpha_h = 0.07 e^{V/20} \quad (4.31e)$$

$$\beta_h = 1/[e^{(3+V/10)} + 1]. \quad (4.31f)$$

The parameter values are $C = 1.0 \mu\text{F}/\text{cm}^2$, $g_K = 36 \text{ mmho}/\text{cm}^2$, $g_{Na} = 120 \text{ mmho}/\text{cm}^2$, $g_L = 0.3 \text{ mmho}/\text{cm}^2$, $V_K = 12 \text{ mV}$, $V_{Na} = -115 \text{ mV}$, and $V_L = 10.6 \text{ mV}$. The unit mho represents ohm^{-1} , and the unit of time is milliseconds (ms). These parameters assume that the resting potential of the nerve cell is zero; however, we now know that the resting potential is about -70 mV .

We can use the ODE solver to solve (4.30) with the state vector $\{V, n, m, h, t\}$; the rates are given by the right-hand side of (4.30). The following questions ask you to explore the properties of the model.

- Write a program to plot n , m , and h as a function of V in the steady state (for which $\dot{n} = \dot{m} = \dot{h} = 0$). Describe how these gates are operating.
- Write a program to simulate the nerve cell membrane potential and plot $V(t)$. You can use a simple Euler algorithm with a time step of 0.01 ms . Describe the behavior of the potential when the external current is 0 .
- Consider a current that is zero except for a one millisecond interval. Try a current spike amplitude of $7 \mu\text{A}$ (that is, the external current equals 7 in our units). Describe the resulting nerve impulse $V(t)$. Is there a threshold value for the current below which there is no large spike but only a broad peak?
- A constant current should produce a train of spikes. Try different amplitudes for the current and determine if there is a threshold current and how the spacing between spikes depends on the amplitude of the external current.
- Consider a situation where there is a steady external current I_1 for 20 ms and then the current increases to $I_2 = I_1 + \Delta I$. There are three types of behavior depending on I_2 and ΔI . Describe the behavior for the following four situations: (1) $I_1 = 2.0 \mu\text{A}$, $\Delta I = 1.5 \mu\text{A}$; (2) $I_1 = 2.0 \mu\text{A}$, $\Delta I = 5.0 \mu\text{A}$; (3) $I_1 = 7.0 \mu\text{A}$, $\Delta I = 1.0 \mu\text{A}$; and (4) $I_1 = 7.0 \mu\text{A}$, $\Delta I = 4.0 \mu\text{A}$. Try other values of I_1 and ΔI as well. In which cases do you obtain a steady spike train? Which cases produce a single spike? What other behavior do you find?
- Once a spike is triggered, it is frequently difficult to trigger another spike. Consider a current pulse at $t = 20 \text{ ms}$ of $7 \mu\text{A}$ that lasts for one millisecond. Then give a second

current pulse of the same amplitude and duration at $t = 25 \text{ ms}$. What happens? What happens if you add a third pulse at 30 ms ? ■

REFERENCES AND SUGGESTIONS FOR FURTHER READING

- F. S. Acton, *Numerical Methods That Work* (The Mathematical Association of America, 1999), Chapter 5.
- G. L. Baker and J. P. Gollub, *Chaotic Dynamics: An Introduction*, 2nd ed. (Cambridge University Press, 1996). A good introduction to the notion of phase space.
- Eugene I. Butikov, "Square-wave excitation of a linear oscillator," *Am. J. Phys.* **72**, 469–476 (2004).
- A. Douglas Davis, *Classical Mechanics* (Saunders College Publishing, 1986). The author gives simple numerical solutions of Newton's equations of motion. Much emphasis is given to the harmonic oscillator problem.
- S. Eubank, W. Miner, T. Tajima, and J. Wiley, "Interactive computer simulation and analysis of Newtonian dynamics," *Am. J. Phys.* **57**, 457–463 (1989).
- Richard P. Feynman, Robert B. Leighton, and Matthew Sands, *The Feynman Lectures on Physics*, Vol. 1 (Addison-Wesley, 1963). Chapters 21 and 23–25 are devoted to various aspects of harmonic motion.
- A. P. French, *Newtonian Mechanics* (W. W. Norton & Company, 1971). An introductory level text with a good discussion of oscillatory motion.
- M. Gitterman, "Classical harmonic oscillator with multiplicative noise," *Physica A* **352**, 309–334 (2005). The analysis is analytical and at the graduate level. However, it would be straightforward to reproduce most of the results after you learn about random processes in Chapter 7.
- A. L. Hodgkin and A. F. Huxley, "A quantitative description of ion currents and its applications to conduction and excitation in nerve membranes," *J. Physiol. (Lond.)* **117**, 500–544 (1952).
- Charles Kittel, Walter D. Knight, and Malvin A. Ruderman, *Mechanics*, 2nd ed., revised by A. Carl Helmholz and Burton J. Moyer (McGraw-Hill, 1973).
- R. Lefever and G. Nicolis, "Chemical instabilities and sustained oscillations," *J. Theor. Biol.* **30**, 267 (1971).
- Jerry B. Marion and Stephen T. Thornton, *Classical Dynamics*, 5th ed. (Harcourt, 2004). Excellent discussion of linear and nonlinear oscillators.
- M. F. McInerney, "Computer-aided experiments with the damped harmonic oscillator," *Am. J. Phys.* **53**, 991–996 (1985).
- William H. Press, Saul A. Teukolsky, William T. Vetterling, and Brian P. Flannery, *Numerical Recipes*, 2nd ed. (Cambridge University Press, 1992). Chapter 16 discusses the integration of ordinary differential equations.
- Scott Hamilton, *An Analog Electronics Companion* (Cambridge University Press, 2003). A good discussion of the physics and mathematics of basic circuit design including an extensive introduction to circuit simulation using the PSpice simulation program.
- S. C. Zilio, "Measurement and analysis of large-angle pendulum motion," *Am. J. Phys.* **50**, 450–452 (1982).