

Numerical projects in Quantum mechanics

Morten Hjorth-Jensen^{1,2}

¹Department of Physics, University of Oslo

²Department of Physics and Astronomy and National Superconducting Cyclotron Laboratory, Michigan State University

2017

Quantum dots in three dimensions, one ad two electrons

We are first interested in the solution of the radial part of Schroedinger's equation for one electron. This equation reads

$$-\frac{\hbar^2}{2m} \left(\frac{1}{r^2} \frac{d}{dr} r^2 \frac{d}{dr} - \frac{l(l+1)}{r^2} \right) R(r) + V(r)R(r) = ER(r).$$

In our case $V(r)$ is the harmonic oscillator potential $(1/2)kr^2$ with $k = m\omega^2$ and E is the energy of the harmonic oscillator in three dimensions. The oscillator frequency is ω and the energies are

$$E_{nl} = \hbar\omega \left(2n + l + \frac{3}{2} \right),$$

with $n = 0, 1, 2, \dots$ and $l = 0, 1, 2, \dots$.

Spherical coordinates

Since we have made a transformation to spherical coordinates it means that $r \in [0, \infty)$. The quantum number l is the orbital momentum of the electron. Then we substitute $R(r) = (1/r)u(r)$ and obtain

$$-\frac{\hbar^2}{2m} \frac{d^2}{dr^2} u(r) + \left(V(r) + \frac{l(l+1)}{r^2} \frac{\hbar^2}{2m} \right) u(r) = Eu(r).$$

The boundary conditions are $u(0) = 0$ and $u(\infty) = 0$.

Making the equations dimensionless

We introduce a dimensionless variable $\rho = (1/\alpha)r$ where α is a constant with dimension length and get

$$-\frac{\hbar^2}{2m\alpha^2} \frac{d^2}{d\rho^2} u(\rho) + \left(V(\rho) + \frac{l(l+1)}{\rho^2} \frac{\hbar^2}{2m\alpha^2} \right) u(\rho) = Eu(\rho).$$

In project 2 we choose $l = 0$. Inserting $V(\rho) = (1/2)k\alpha^2\rho^2$ we end up with

$$-\frac{\hbar^2}{2m\alpha^2} \frac{d^2}{d\rho^2} u(\rho) + \frac{k}{2} \alpha^2 \rho^2 u(\rho) = Eu(\rho).$$

We multiply thereafter with $2m\alpha^2/\hbar^2$ on both sides and obtain

$$-\frac{d^2}{d\rho^2} u(\rho) + \frac{mk}{\hbar^2} \alpha^4 \rho^2 u(\rho) = \frac{2m\alpha^2}{\hbar^2} Eu(\rho).$$

And introducing a natural length scale

We have thus

$$-\frac{d^2}{d\rho^2} u(\rho) + \frac{mk}{\hbar^2} \alpha^4 \rho^2 u(\rho) = \frac{2m\alpha^2}{\hbar^2} Eu(\rho).$$

The constant α can now be fixed so that

$$\frac{mk}{\hbar^2} \alpha^4 = 1,$$

or

$$\alpha = \left(\frac{\hbar^2}{mk} \right)^{1/4}.$$

Defining

$$\lambda = \frac{2m\alpha^2}{\hbar^2} E,$$

we can rewrite Schroedinger's equation as

$$-\frac{d^2}{d\rho^2} u(\rho) + \rho^2 u(\rho) = \lambda u(\rho).$$

This is the first equation to solve numerically. In three dimensions the eigenvalues for $l = 0$ are $\lambda_0 = 3, \lambda_1 = 7, \lambda_2 = 11, \dots$

Discretizing I

We use the by now standard expression for the second derivative of a function u

$$u'' = \frac{u(\rho + h) - 2u(\rho) + u(\rho - h)}{h^2} + O(h^2), \quad (1)$$

where h is our step. Next we define minimum and maximum values for the variable ρ , $\rho_{\min} = 0$ and ρ_{\max} , respectively. You need to check your results for the energies against different values ρ_{\max} , since we cannot set $\rho_{\max} = \infty$.

Discretizing II

With a given number of steps, n_{step} , we then define the step h as

$$h = \frac{\rho_{\text{max}} - \rho_{\text{min}}}{n_{\text{step}}}.$$

Define an arbitrary value of ρ as

$$\rho_i = \rho_{\text{min}} + ih \quad i = 0, 1, 2, \dots, n_{\text{step}}$$

we can rewrite the Schrödinger equation for ρ_i as

$$-\frac{u(\rho_i + h) - 2u(\rho_i) + u(\rho_i - h)}{h^2} + \rho_i^2 u(\rho_i) = \lambda u(\rho_i),$$

or in a more compact way

$$-\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} + \rho_i^2 u_i = -\frac{u_{i+1} - 2u_i + u_{i-1}}{h^2} + V_i u_i = \lambda u_i,$$

where $V_i = \rho_i^2$ is the harmonic oscillator potential.

Discretizing III

Define first the diagonal matrix element

$$d_i = \frac{2}{h^2} + V_i,$$

and the non-diagonal matrix element

$$e_i = -\frac{1}{h^2}.$$

In this case the non-diagonal matrix elements are given by a mere constant. *All non-diagonal matrix elements are equal.*

With these definitions the Schroedinger equation takes the following form

$$d_i u_i + e_{i-1} u_{i-1} + e_{i+1} u_{i+1} = \lambda u_i,$$

where u_i is unknown. We can write the latter equation as a matrix eigenvalue problem

$$\begin{pmatrix} d_1 & e_1 & 0 & 0 & \dots & 0 & 0 \\ e_1 & d_2 & e_2 & 0 & \dots & 0 & 0 \\ 0 & e_2 & d_3 & e_3 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & \dots & d_{n_{\text{step}}-2} & e_{n_{\text{step}}-1} \\ 0 & \dots & \dots & \dots & \dots & e_{n_{\text{step}}-1} & d_{n_{\text{step}}-1} \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ \dots \\ \dots \\ u_{n_{\text{step}}-1} \end{pmatrix} = \lambda \begin{pmatrix} u_1 \\ u_2 \\ \dots \\ \dots \\ \dots \\ u_{n_{\text{step}}-1} \end{pmatrix} \quad (2)$$

or if we wish to be more detailed, we can write the tridiagonal matrix as

$$\begin{pmatrix} \frac{2}{h^2} + V_1 & -\frac{1}{h^2} & 0 & 0 & \dots & 0 & 0 \\ -\frac{1}{h^2} & \frac{2}{h^2} + V_2 & -\frac{1}{h^2} & 0 & \dots & 0 & 0 \\ 0 & -\frac{1}{h^2} & \frac{2}{h^2} + V_3 & -\frac{1}{h^2} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & \dots & \frac{2}{h^2} + V_{n_{\text{step}}-2} & -\frac{1}{h^2} \\ 0 & \dots & \dots & \dots & \dots & -\frac{1}{h^2} & \frac{2}{h^2} + V_{n_{\text{step}}-1} \end{pmatrix} \quad (3)$$

Recall that the solutions are known via the boundary conditions at $i = n_{\text{step}}$ and at the other end point, that is for ρ_0 . The solution is zero in both cases.

From one to two electrons, studying the role of the repulsive Coulomb force

We are going to study two electrons in a harmonic oscillator well which also interact via a repulsive Coulomb interaction. Let us start with the single-electron equation written as

$$-\frac{\hbar^2}{2m} \frac{d^2}{dr^2} u(r) + \frac{1}{2} kr^2 u(r) = E^{(1)} u(r),$$

where $E^{(1)}$ stands for the energy with one electron only. For two electrons with no repulsive Coulomb interaction, we have the following Schroedinger equation

$$\left(-\frac{\hbar^2}{2m} \frac{d^2}{dr_1^2} - \frac{\hbar^2}{2m} \frac{d^2}{dr_2^2} + \frac{1}{2} kr_1^2 + \frac{1}{2} kr_2^2 \right) u(r_1, r_2) = E^{(2)} u(r_1, r_2).$$

Rewriting the equations

Note that we deal with a two-electron wave function $u(r_1, r_2)$ and two-electron energy $E^{(2)}$.

With no interaction this can be written out as the product of two single-electron wave functions, that is we have a solution on closed form.

We introduce the relative coordinate $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$ and the center-of-mass coordinate $\mathbf{R} = 1/2(\mathbf{r}_1 + \mathbf{r}_2)$. With these new coordinates, the radial Schroedinger equation reads

$$\left(-\frac{\hbar^2}{m} \frac{d^2}{dr^2} - \frac{\hbar^2}{4m} \frac{d^2}{dR^2} + \frac{1}{4} kr^2 + kR^2 \right) u(r, R) = E^{(2)} u(r, R).$$

Center of mass and relative coordinates

The equations for r and R can be separated via the ansatz for the wave function $u(r, R) = \psi(r)\phi(R)$ and the energy is given by the sum of the relative energy E_r and the center-of-mass energy E_R , that is

$$E^{(2)} = E_r + E_R.$$

We add then the repulsive Coulomb interaction between two electrons, namely a term

$$V(r_1, r_2) = \frac{\beta e^2}{|\mathbf{r}_1 - \mathbf{r}_2|} = \frac{\beta e^2}{r},$$

with $\beta e^2 = 1.44 \text{ eVnm}$.

The equation for the relative motion

Adding this term, the r -dependent Schroedinger equation becomes

$$\left(-\frac{\hbar^2}{m} \frac{d^2}{dr^2} + \frac{1}{4} k r^2 + \frac{\beta e^2}{r} \right) \psi(r) = E_r \psi(r).$$

This equation is similar to the one we had previously in parts (a) and (b) and we introduce again a dimensionless variable $\rho = r/\alpha$. Repeating the same steps, we arrive at

$$-\frac{d^2}{d\rho^2} \psi(\rho) + \frac{mk}{4\hbar^2} \alpha^4 \rho^2 \psi(\rho) + \frac{m\alpha\beta e^2}{\rho\hbar^2} \psi(\rho) = \frac{m\alpha^2}{\hbar^2} E_r \psi(\rho).$$

Further manipulations

We want to manipulate this equation further to make it as similar to the one-electron case as possible. We define a 'frequency'

$$\omega_r^2 = \frac{1}{4} \frac{mk}{\hbar^2} \alpha^4,$$

and fix the constant α by requiring

$$\frac{m\alpha\beta e^2}{\hbar^2} = 1$$

or

$$\alpha = \frac{\hbar^2}{m\beta e^2}.$$

The new equation to solve

Defining

$$\lambda = \frac{m\alpha^2}{\hbar^2} E,$$

we can rewrite Schroedinger's equation as

$$-\frac{d^2}{d\rho^2} \psi(\rho) + \omega_r^2 \rho^2 \psi(\rho) + \frac{1}{\rho} \psi(\rho) = \lambda \psi(\rho).$$

We treat ω_r as a parameter which reflects the strength of the oscillator potential.

With no repulsive Coulomb interaction you should get a result which corresponds to the relative energy of a non-interacting system. Make sure your results are stable as functions of ρ_{\max} and the number of steps.

We are only interested in the ground state with $l = 0$. We omit the center-of-mass energy.

For specific oscillator frequencies, the above equation has analytic answers, see the article by M. Taut, Phys. Rev. A 48, 3561 - 3566 (1993). The article can be retrieved from the following web address http://prola.aps.org/abstract/PRA/v48/i5/p3561_1.

The Python code

The code sets up the Hamiltonian matrix by defining the minimum and maximum values of r with a maximum value of integration points. These are set in the initialization function. It plots the eigenfunctions of the three lowest eigenstates.

```
#Program which solves the one-particle Schrodinger equation
#for a potential specified in function
#potential(). This example is for the harmonic oscillator in 3d

from matplotlib import pyplot as plt
import numpy as np
#Function for initialization of parameters
def initialize():
    RMin = 0.0
    RMax = 10.0
    lOrbital = 0
    Dim = 400
    return RMin, RMax, lOrbital, Dim
# Here we set up the harmonic oscillator potential
def potential(r):
    return r*r

#Get the boundary, orbital momentum and number of integration points
RMin, RMax, lOrbital, Dim = initialize()

#Initialize constants
Step = RMax/(Dim+1)
DiagConst = 2.0 / (Step*Step)
NondiagConst = -1.0 / (Step*Step)
OrbitalFactor = lOrbital * (lOrbital + 1.0)

#Calculate array of potential values
v = np.zeros(Dim)
r = np.linspace(RMin,RMax,Dim)
for i in xrange(Dim):
    r[i] = RMin + (i+1) * Step;
    v[i] = potential(r[i]) + OrbitalFactor/(r[i]*r[i]);

#Setting up a tridiagonal matrix and finding eigenvectors and eigenvalues
Hamiltonian = np.zeros((Dim,Dim))
Hamiltonian[0,0] = DiagConst + v[0];
```

```

Hamiltonian[0,1] = NondiagConst;
for i in xrange(1,Dim-1):
    Hamiltonian[i,i-1] = NondiagConst;
    Hamiltonian[i,i] = DiagConst + v[i];
    Hamiltonian[i,i+1] = NondiagConst;
Hamiltonian[Dim-1,Dim-2] = NondiagConst;
Hamiltonian[Dim-1,Dim-1] = DiagConst + v[Dim-1];
# diagonalize and obtain eigenvalues, not necessarily sorted
EigValues, EigVectors = np.linalg.eig(Hamiltonian)
# sort eigenvectors and eigenvalues
permute = EigValues.argsort()
EigValues = EigValues[permute]
EigVectors = EigVectors[:,permute]
# now plot the results for the three lowest lying eigenstates
for i in xrange(3):
    print EigValues[i]
FirstEigvector = EigVectors[:,0]
SecondEigvector = EigVectors[:,1]
ThirdEigvector = EigVectors[:,2]
plt.plot(r, FirstEigvector**2, 'b-', r, SecondEigvector**2, 'g-', r, ThirdEigvector**2, 'r-')
plt.axis([0,4.6,0.0, 0.025])
plt.xlabel(r'$r$')
plt.ylabel(r'Radial probability $r^2|R(r)|^2$')
plt.title(r'Radial probability distributions for three lowest-lying states')
plt.savefig('eigenvector.pdf')
plt.show()

```