

Quantum Computing and Quantum Machine Learning

Morten Hjorth-Jensen¹

Department of Physics, University of Oslo¹

March 4, 2026

© 1999-2026, Morten Hjorth-Jensen. Released under CC Attribution-NonCommercial 4.0 license

Plans for the week of March 2-6

1. Reminder on basics of the VQE method and how to perform measurements for the simpler one- and two-qubit Hamiltonians
2. Simulating efficiently Hamiltonians on quantum computers with the VQE method and gradient descent to optimize the state function ansatz
3. Introducing the Lipkin model
4. Work on project 1

Readings and notebooks

- ▶ For the discussion of one-qubit, two-qubit and other gates, sections 2.6-2.11 and 3.1-3.4 of Hundt's book **Quantum Computing for Programmers**, contain most of the relevant information.
- ▶ The VQE algorithm is discussed in Hundt's section 6.11, note that the solution of the two-qubit system is outdated, measuring on one qubit only is not the present standard
- ▶ See also the VQE review article by Tilly et al.
- ▶ Jupyter-notebook on VQE and single-qubit problem at https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week6/ipynb/single_qubit_vqe_modified.ipynb
- ▶ Jupyter-notebook on VQE and two-qubit problem at https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week6/ipynb/two_qubit_vqe_modified.ipynb

More material

- ▶ Notes on why we measure on all qubits and not on one as in Hundt's chapter 6.11, see additional pdf file at <https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week7/pdf/measurenotes.pdf>
- ▶ Jupyter-notebook on VQE for the Lipkin model at <https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week7/ipynb/lipkinvqe.ipynb>

More background material

For the Lipkin model, we recommend strongly the work of LaRose and collaborators, see <https://journals.aps.org/prc/abstract/10.1103/PhysRevC.106.024319>, the article is also available at <https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/articles/PhysRevC.106.024319.pdf>. See in particular section 3.

Two-qubit Hamiltonian

Here we discuss how to rewrite the two-qubit Hamiltonian defined by the following Hamiltonian matrix (project 1)

$$\mathcal{H} = \begin{bmatrix} \epsilon_1 + V_z & 0 & 0 & V_x \\ 0 & \epsilon_2 - V_z & V_x & 0 \\ 0 & H_x & \epsilon_3 - V_z & 0 \\ H_x & 0 & 0 & \epsilon_4 + V_z \end{bmatrix}.$$

This Hamiltonian can be rewritten in terms of various one-qubit matrices.

Definitions

We define

$$\epsilon_{II} = \frac{\epsilon_1 + \epsilon_2 + \epsilon_3 + \epsilon_4}{4},$$

$$\epsilon_{ZI} = \frac{\epsilon_1 + \epsilon_2 - \epsilon_3 - \epsilon_4}{4},$$

$$\epsilon_{IZ} = \frac{\epsilon_1 - \epsilon_2 + \epsilon_3 - \epsilon_4}{4},$$

$$\epsilon_{ZZ} = \frac{\epsilon_1 - \epsilon_2 - \epsilon_3 + \epsilon_4}{4}.$$

The Hamiltonian in terms of Pauli-**X** and Pauli-**Z** matrices

With these definitions we can rewrite our two-qubit Hamiltonian as

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_I$$

with

$$\mathcal{H}_0 = \epsilon_{II} I \otimes I + \epsilon_{ZI} Z \otimes I + \epsilon_{IZ} I \otimes Z + \epsilon_{ZZ} Z \otimes Z,$$

and

$$\mathcal{H}_I = V_z Z \otimes Z + V_x X \otimes X.$$

Why don't we measure on one qubit?

- ▶ Notes on why we measure on all qubits and not on one as in Hundt's chapter 6.11, see additional pdf file at <https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week7/pdf/measurenotes.pdf>
- ▶ * Jupyter-notebook on VQE and two-qubit problem at https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week6/ipynb/two_qubit_vqe_modified.ipynb
- ▶ See also additional notes from last week at <https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week6/pdf/additional2week6.pdf>

Lipkin model

We will study a schematic model (the Lipkin model, see Nuclear Physics **62** (1965) 188), for the interaction among 2 and more fermions that can occupy two different energy levels.

In project 1 we consider a two-fermion case and a four-fermion case. For four fermions, the case we consider in the examples here, each level has degeneration $d = 4$, leading to different total spin values. The two levels have quantum numbers $\sigma = \pm 1$, with the upper level having $2\sigma = +1$ and energy $\varepsilon_1 = \varepsilon/2$. The lower level has $2\sigma = -1$ and energy $\varepsilon_2 = -\varepsilon/2$. That is, the lowest single-particle level has negative spin projection (or spin down), while the upper level has spin up. In addition, the substates of each level are characterized by the quantum numbers $p = 1, 2, 3, 4$.

Four fermion case

We define the single-particle states (for the four fermion case which we will work on here)

$$|u_{\sigma=-1,p}\rangle = a_{-p}^{\dagger}|0\rangle \quad |u_{\sigma=1,p}\rangle = a_{+p}^{\dagger}|0\rangle.$$

The single-particle states span an orthonormal basis.

Hamiltonian

The Hamiltonian of the system is given by

$$\hat{H} = \hat{H}_0 + \hat{H}_1 + \hat{H}_2$$

$$\hat{H}_0 = \frac{1}{2}\varepsilon \sum_{\sigma,p} \sigma a_{\sigma,p}^\dagger a_{\sigma,p}$$

$$\hat{H}_1 = \frac{1}{2}V \sum_{\sigma,p,p'} a_{\sigma,p}^\dagger a_{\sigma,p'}^\dagger a_{-\sigma,p'} a_{-\sigma,p}$$

$$\hat{H}_2 = \frac{1}{2}W \sum_{\sigma,p,p'} a_{\sigma,p}^\dagger a_{-\sigma,p'}^\dagger a_{\sigma,p'} a_{-\sigma,p}$$

where V and W are constants. The operator H_1 can move pairs of fermions while H_2 is a spin-exchange term. The latter moves a pair of fermions from a state $(p\sigma, p' - \sigma)$ to a state $(p - \sigma, p'\sigma)$.

Quasispin operators

We are going to rewrite the above Hamiltonian in terms of so-called quasispin operators

$$\hat{J}_+ = \sum_p a_{p+}^\dagger a_{p-}$$

$$\hat{J}_- = \sum_p a_{p-}^\dagger a_{p+}$$

$$\hat{J}_z = \frac{1}{2} \sum_{p\sigma} \sigma a_{p\sigma}^\dagger a_{p\sigma}$$

$$\hat{J}^2 = J_+ J_- + J_z^2 - J_z$$

These operators obey the commutation relations for angular momentum.

Including the number operator

We can in turn express $\hat{H}$ in terms of the above quasispin operators and the number operator

$$\hat{N} = \sum_{p\sigma} a_{p\sigma}^\dagger a_{p\sigma}.$$

Final Hamiltonian

Using the quasispin operators we can rewrite the Hamiltonian as

$$H = \varepsilon J_z + \frac{1}{2} V (J_+^2 + J_-^2) + \frac{1}{2} W (-N + J_+ J_- + J_- J_+). \quad (1)$$

Original Lipkin Hamiltonian

The Lipkin model Hamiltonian in terms of quasispin operators reads

$$H = \varepsilon J_z + \frac{1}{2} V (J_+^2 + J_-^2) + \frac{1}{2} W (-N + J_+ J_- + J_- J_+).$$

We use the standard commutation relations

$$[J_+, J_-] = 2J_z,$$

and the Casimir operator

$$J^2 = J_z^2 + \frac{1}{2}(J_+ J_- + J_- J_+).$$

More compact expression

From this we obtain

$$J_+ J_- = J^2 - J_z(J_z - 1), \quad J_- J_+ = J^2 - J_z(J_z + 1).$$

Adding the two expressions:

$$J_+ J_- + J_- J_+ = 2(J^2 - J_z^2).$$

Simplified Hamiltonian

Using

$$J_+ J_- + J_- J_+ = 2(J^2 - J_z^2)$$

and insert into the Hamiltonian:

$$H = \varepsilon J_z + \frac{1}{2} V (J_+^2 + J_-^2) + \frac{1}{2} W (-N + 2(J^2 - J_z^2)).$$

This gives

$$H = \varepsilon J_z + \frac{V}{2} (J_+^2 + J_-^2) + W (J^2 - J_z^2) - \frac{WN}{2}.$$

Physical interpretation

- ▶ The Hamiltonian is now expressed in terms of:
 - ▶ Linear term: εJ_z
 - ▶ Pairing term: $J_+^2 + J_-^2$
 - ▶ Quadratic term: J_z^2
 - *Casimir term: J^2
- ▶ The constant $-\frac{WN}{2}$ only shifts the spectrum.

Fixed quasispin sector ($j = N/2$)

In the Lipkin model we typically work in a fixed irreducible representation:

$$J^2 = j(j+1), \quad j = \frac{N}{2}.$$

Thus

$$H = \varepsilon J_z + \frac{V}{2}(J_+^2 + J_-^2) - WJ_z^2 + Wj(j+1) - \frac{WN}{2}.$$

Final simplified form

Using $N = 2j$, the constant simplifies:

$$Wj(j+1) - \frac{WN}{2} = Wj^2.$$

Final result

$$H = \varepsilon J_z + \frac{V}{2}(J_+^2 + J_-^2) - WJ_z^2 + Wj^2.$$

The term Wj^2 is a constant shift and can often be omitted.

Solving on a quantum computer

To solve the Lipkin model with a quantum computer, the first step is to map the system to a set of qubits. We will restrict ourselves here to the half-filled case where the number of particles N equals the degeneracy of the states Ω . One could assign each possible state (n, σ) a qubit such that the qubit being in the state $|1\rangle$ or $|0\rangle$ would imply that the state (n, σ) is occupied or unoccupied, respectively. This mapping scheme (which we will call occupation mapping) requires 2Ω qubits.

Number of qubits

However, because there are only two energy levels in the Lipkin model, any other natural mapping is possible. In this mapping scheme (which we will call level mapping) each doublet $((n, +1), (n, -1))$ would be assigned a qubit such that the qubit being in the state $|0\rangle$ or $|1\rangle$ would imply that the particle is in the $(n, +1)$ or $(n, -1)$ state, respectively. Note that these are the only two possible configurations of the doublet as we are restricting ourselves to the half-filled case and the Lipkin Hamiltonian only moves particles between energy levels, not degenerate states. Thus the level mapping only requires Ω qubits which is half that of the occupation mapping.

Converting to Pauli matrices

The Hamiltonian takes the form (leaving out the W term)

$$H = \epsilon J_z + \frac{1}{2} V (J_+^2 + J_-^2). \quad (2)$$

Writing in terms of one-body operators

Plugging the mapping from the total J operators to the individual one-body j operators yields

$$H = \epsilon \sum_n j_z^{(n)} + \frac{1}{2} V \left[\left(\sum_n j_+^{(n)} \right)^2 + \left(\sum_n j_-^{(n)} \right)^2 \right] \quad (3)$$

$$= \epsilon \sum_n j_z^{(n)} + \frac{1}{2} V \sum_{n,m} \left(j_+^{(n)} j_+^{(m)} + j_-^{(n)} j_-^{(m)} \right) \quad (4)$$

$$= \epsilon \sum_n j_z^{(n)} + 2V \sum_{n < m} \left(j_x^{(n)} j_x^{(m)} - j_y^{(n)} j_y^{(m)} \right), \quad (5)$$

where we have used the definitions

$$j_{\pm}^{(n)} = j_x^{(n)} \pm i j_y^{(n)}.$$

Explicit transformations

To convert to Pauli matrices, we make the transformations

$$j_x^{(n)} \rightarrow X_n/2,$$

and

$$j_y^{(n)} \rightarrow Y_n/2,$$

and finally

$$j_z^{(n)} \rightarrow Z_n/2,$$

which preserve the above $SU(2)$ commutation relations. The factor of $1/2$ is due to the eigenvalues of the Pauli matrices being ± 1 while we are dealing with spin $1/2$ particles.

Hamiltonian in terms of Pauli matrices

This transforms our Hamiltonian into

$$H = \frac{1}{2}\epsilon \sum_{k=1}^n Z_k + \frac{1}{2}V \sum_{k \neq j=1}^N (X_k X_j - Y_k Y_j).$$

With this form, we can clearly see that the first (one-body) term in the Hamiltonian returns the energy $-\epsilon/2$ or $+\epsilon/2$ if the qubit representing the particle of a doublet is in the ground ($|1\rangle$) or excited ($|0\rangle$) state, respectively.

Action of Hamiltonian

The action of the second (two-body) term in the Hamiltonian can be determined by noting that

$$\frac{1}{2}(XX - YY)|00\rangle = |11\rangle, \quad (6)$$

$$\frac{1}{2}(XX - YY)|01\rangle = 0, \quad (7)$$

$$\frac{1}{2}(XX - YY)|10\rangle = 0, \quad (8)$$

$$\frac{1}{2}(XX - YY)|11\rangle = |00\rangle. \quad (9)$$

That is, the two-body term moves a pair of particles between the ground states $|00\rangle$ and the excited states $|11\rangle$ of their respective doublets.

System to diagonalize by traditional methods

Using spin-eigenstates as the Hamiltonian basis, we define states through the normal approach $|J, J_z\rangle$ with J and J_z as spin and spin-projections, respectively. The states $J_z = \pm J$ are the easiest to construct, corresponding to a single level being completely filled. States in between can then be found using the quasi-spin ladder operators following

$$J_{\pm}|J, J_z\rangle = \sqrt{J(J+1) - J_z(J_z \pm 1)}|J, J_z \pm 1\rangle.$$

Using this basis for the quasi-spin Hamiltonian, we can rewrite the explicit matrix $H_{J_z, J'_z} = \langle J, J_z | H | J, J'_z \rangle$ can be constructed.

The $J = 1$ case

For $N = 2$ particles, we have the triplet $J = 1$, with three possible projection $J_z = 0, \pm 1$. This gives the following matrix

$$H = \begin{bmatrix} -\epsilon & 0 & V \\ 0 & W & 0 \\ V & 0 & \epsilon \end{bmatrix}.$$

By solving the eigenvalue problem, the ground state energy can be exactly found.

The $J = 2$ case

Similarly, for the $N = 4$ particles case we have $J = 2$ with five possible projections $J_z = 0, \pm 1, \pm 2$. This gives us the Hamiltonian matrix

$$H = \begin{bmatrix} -2\epsilon & 0 & \sqrt{6}V & 0 & 0 \\ 0 & -\epsilon + 3W & 0 & 3V & 0 \\ \sqrt{6}V & 0 & 4W & 0 & \sqrt{6}V \\ 0 & 3V & 0 & \epsilon + 3W & 0 \\ 0 & 0 & \sqrt{6}V & 0 & 2\epsilon \end{bmatrix}.$$

More rewriting

Following the work of LaRose and collaborators, see <https://journals.aps.org/prc/abstract/10.1103/PhysRevC.106.024319>, we can rewrite the Lipkin Hamiltonian for efficient adaptations to quantum computing.

We have

$$J_z = \sum_i^N j_z^i \quad J_{\pm} = \sum_i^N j_{\pm}^i = \sum_i^N (j_x^i \pm ij_y^i),$$

with N being the number of particles. Additionally, since we have spin-1/2 fermions, the mapping to Pauli matrices follow

$$j_x^i = \frac{1}{2}X_i, \quad j_y^i = \frac{1}{2}Y_i, \quad j_z^i = \frac{1}{2}Z_i.$$

This means that we require N qubits to calculate properties of a N particle system, if no more symmetry reductions are considered.

Lipkin Hamiltonian for quantum computing

We rewrote our Hamiltonian as

$$H_0 = \frac{\epsilon}{2} \sum_p Z_p,$$

$$H_1 = \frac{1}{2} V \sum_{p < q} (X_p X_q - Y_p Y_q),$$

$$H_2 = \frac{1}{2} W \sum_{p < q} (X_p X_q + Y_p Y_q).$$

For two particles coupling to spin $J = 1$ we have

$$H^{N=2} = \frac{\epsilon}{2} (Z_1 + Z_2) + \frac{W + V}{2} X_1 X_2 - \frac{W - V}{2} Y_1 Y_2.$$

For four fermions

Our Hamiltonian for $N = 4$ is

$$\begin{aligned} H^{N=4} = & \frac{\epsilon}{2} (Z_1 + Z_2 + Z_3 + Z_4) + \frac{W - V}{2} \left(X_1 X_2 + \right. \\ & X_1 X_3 + X_1 X_4 + X_2 X_3 + X_3 X_4 + X_3 X_4 \Big) \\ & + \frac{W + V}{2} \left(Y_1 Y_2 + Y_1 Y_3 + Y_1 Y_4 + Y_2 Y_3 \right. \\ & \left. + Y_3 Y_4 + Y_3 Y_4 \right). \end{aligned}$$

Note that well that a term like Z_1 reads $Z_1 \otimes I_2 \otimes I_3 \otimes I_4$ where the subscript points to qubit i and all matrices are 2×2 matrices.

Leaving out the W term

If we set $W = 0$ we can simplify the Hamiltonian. For this we note that if we only include the H_0 and H_1 terms, spins differing by ± 2 are the only possible non-diagonal couplings. Since H_0 is diagonal and single particle energies are degenerate, it does not break any symmetry. Instead of the $2J + 1$ spin projections, we simply have $J + 1$ relevant states. From <https://journals.aps.org/prc/abstract/10.1103/PhysRevC.106.024319>, the two-qubit Hamiltonian for $N = 4$ can be written as

$$H_{W=0}^{N=4} = \epsilon(Z_1 + Z_2) + \frac{\sqrt{6}}{2} V(X_1 + X_2 + Z_1 X_0 - X_1 Z_0).$$

This is computationally more efficient since we only need half of the qubits.

More details

Here we give some additional details behind the rewriting of the Lipkin Hamiltonian in term of Pauli matrices and the identity matrix. For the one-body term we have

$$H_0 = \epsilon J_z = \epsilon \sum_p j_z^p = \frac{\epsilon}{2} \sum_p Z_p$$

Moving on to the H_1 term, we need to expand the square of the $J_{\pm}$ operators. Starting with J_+ we find

$$\begin{aligned} J_+^2 &= \left(\sum_p j_x^p + \imath j_y^p \right)^2 = \sum_{pq} (j_x^p + \imath j_y^p)(j_x^q + \imath j_y^q) \\ &= \sum_{pq} (j_x^p j_x^q - j_y^p j_y^q + \imath j_y^p j_x^q + \imath j_x^p j_y^q) . \end{aligned}$$

Other manipulations

Similarly, the J_-^2 term yields

$$J_-^2 = \sum_{pq} (j_x^p j_x^q - j_y^p j_y^q - j_z^p j_x^q - j_z^p j_y^q).$$

Rewriting we get

$$\begin{aligned} J_+^2 + J_-^2 &= 2 \sum_{pq} j_x^p j_x^q - j_z^p j_z^q \\ &= 2 \sum_p (j_x^p)^2 - (j_y^p)^2 + 4 \sum_{p>q} (j_x^p j_x^q - j_y^p j_y^q) \\ &= \frac{1}{2} \sum_p (X_p^2 - Y_p^2) + \sum_{p>q} (X_p X_q - Y_p Y_q) \end{aligned}$$

Final expression for H_1 and H_2

The diagonal term will cancel out, since the Pauli matrices are involutory, resulting in

$$H_1 = \frac{1}{2}V \sum_{p>q} (X_p X_q - Y_p Y_q).$$

Following the same procedure as above, we have finally

$$H_2 = \frac{1}{2}W \sum_{p<q} (X_p X_q + Y_p Y_q).$$

Jupyter-notebook for the Lipkin model

- ▶ Jupyter-notebook on VQE for the Lipkin model at <https://github.com/CompPhysics/QuantumComputingMachineLearning/blob/gh-pages/doc/pub/week7/ipynb/lipkinvqe.ipynb>

Note that this notebook does not implement the VQE with shots.

Plans for next week

1. We will focus mainly on setting up the quantum circuit for the Lipkin model with $J = 1$ and $J = 2$, rewriting the Lipkin model in terms of Pauli matrices
2. Jordan-Wigner transformation
3. Thereafter, if we get time we start with quantum Fourier transforms and discussions of Quantum Phase estimation algorithm