

Lecture 8 - Linear Mappings Part 1

COMP1046 - Maths for Computer Scientists

Dr. Ferrante Neri / Dr. Tony Bellotti



By the end of this lecture we will have learned:

- ⊙ Mappings
- ⊙ Linear Mappings
- ⊙ Linear Mappings and Vector Spaces

Based on Sections 10.1 and 10.2 of text book (Neri 2018).

Mappings



Definition

Let $(E, +, \cdot)$ and $(F, +, \cdot)$ be two vector spaces defined over the scalar field \mathbb{K} . Let $f : E \rightarrow F$ be a relation.

Let U be a set such that $U \subseteq E$.

The relation f is said *mapping* when

$$\forall \mathbf{u} \in U : \exists ! \mathbf{w} \in F \text{ such that } f(\mathbf{u}) = \mathbf{w}.$$

The set U is said *domain* and is indicated with $\text{dom}(f)$.

A vector \mathbf{w} such that

$$\mathbf{w} = f(\mathbf{u})$$

is said to be the *mapped* (or *transformed*) of \mathbf{u} through f .

Definition

Let f be a mapping $E \rightarrow F$, where E and F are sets associated with the vector spaces $(E, +, \cdot)$ and $(F, +, \cdot)$. The *image* of f , indicated with $\text{Im}(f)$, is a set defined as

$$\text{Im}(f) = \{ \mathbf{w} \in F \mid \exists \mathbf{u} \in E \text{ such that } f(\mathbf{u}) = \mathbf{w} \}.$$

Example

Let $(\mathbb{R}, +, \cdot)$ be a vector space.

An example of mapping $\mathbb{R} \rightarrow \mathbb{R}$ is $f(x) = x^2 + 2x + 2$.

- ⊙ The domain of the mapping $\text{dom}(f) = \mathbb{R}$.
- ⊙ The image $\text{Im}(f) = [1, \infty[$.

Example

Let $(\mathbb{R}^2, +, \cdot)$ and $(\mathbb{R}, +, \cdot)$ be two vector spaces.

An example of mapping $\mathbb{R}^2 \rightarrow \mathbb{R}$ is $f(x, y) = x + 2y + 2$.

- ⊙ The domain of the mapping $\text{dom}(f) = \mathbb{R}^2$.
- ⊙ The image $\text{Im}(f) = \mathbb{R}$.

Example

From the vector spaces $(\mathbb{R}^2, +, \cdot)$ and $(\mathbb{R}^3, +, \cdot)$ an example of mapping $\mathbb{R}^3 \rightarrow \mathbb{R}^2$ is

$$f(x, y, z) = (x + 2y + -z + 2, 6y - 4z + 2).$$

- ⊙ The domain of the mapping $\text{dom}(f) = \mathbb{R}^3$.
- ⊙ The image $\text{Im}(f) = \mathbb{R}^2$.

Definition

Let f be a mapping $E \rightarrow F$, where E and F are sets associated with the vector spaces $(E, +, \cdot)$ and $(F, +, \cdot)$.

- ⊙ The mapping f is said *surjective* if the image of f coincides with F : $\text{Im}(f) = F$.
- ⊙ The mapping f is said *injective* if

$$\forall \mathbf{u}, \mathbf{v} \in E \text{ with } \mathbf{u} \neq \mathbf{v} \Rightarrow f(\mathbf{u}) \neq f(\mathbf{v}).$$

- ⊙ The mapping f is said *bijective* if f is injective and surjective.

Example

Consider these mappings:

- ⊙ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$:
 - Not surjective since $\text{Im}(f) = \mathbb{R}^+$.
 - Not injective because $\exists x_1, x_2$ with $x_1 \neq x_2$ such that $x_1^2 = x_2^2$.
For example if $x_1 = 3$ and $x_2 = -3$, thus $x_1 \neq x_2$, it occurs that $x_1^2 = x_2^2 = 9$.
 - Hence not bijective.
- ⊙ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = e^x$ is injective but not surjective.
Hence, this mapping is not bijective.
- ⊙ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = 2x + 2$ is injective and surjective.
Hence, this mapping is bijective.

Linear Mappings



Definition

Let f be a mapping $E \rightarrow F$, where E and F are sets associated with the vector spaces $(E, +, \cdot)$ and $(F, +, \cdot)$.

The mapping f is said *linear mapping* if the following properties are valid:

- ⊙ additivity: $\forall \mathbf{u}, \mathbf{v} \in E : f(\mathbf{u} + \mathbf{v}) = f(\mathbf{u}) + f(\mathbf{v})$
- ⊙ homogeneity: $\forall \lambda \in \mathbb{K} \text{ and } \forall \mathbf{v} \in E : f(\lambda \mathbf{v}) = \lambda f(\mathbf{v})$

Example

Let us check the linearity of the mapping $f : \mathbb{R} \rightarrow \mathbb{R}$

$$\forall x : f(x) = e^x.$$

Consider two vectors (numbers in this case) x_1 and x_2 .

Calculate $f(x_1 + x_2) = e^{x_1 + x_2}$.

From basic calculus we know that

$$e^{x_1 + x_2} \neq e^{x_1} + e^{x_2}.$$

Therefore, additivity is not verified. Hence the mapping is not linear.

Example

Let us consider the linearity of the mapping $f : \mathbb{R} \rightarrow \mathbb{R}$

$$\forall x : f(x) = 2x.$$

Consider two vectors (numbers in this case) x_1 and x_2 . Then,

$$\begin{aligned} f(x_1 + x_2) &= 2(x_1 + x_2) \\ f(x_1) + f(x_2) &= 2x_1 + 2x_2. \end{aligned}$$

It follows that $f(x_1 + x_2) = f(x_1) + f(x_2)$.

Hence, this mapping is additive.

Continued...

Example

Let us check the homogeneity by considering a generic scalar λ . We have that

$$f(\lambda x) = 2\lambda x$$

$$\lambda f(x) = \lambda 2x.$$

It follows that $f(\lambda x) = \lambda f(x)$.

Hence, since also homogeneity is verified this mapping is linear.

Definition

Let f be a mapping $E \rightarrow F$, where E and F are sets associated with the vector spaces $(E, +, \cdot)$ and $(F, +, \cdot)$.

The mapping f is said *affine mapping* if the mapping

$$g(\mathbf{v}) = f(\mathbf{v}) - f(\mathbf{o})$$

is linear.

Affine Mapping

Example

Consider the mapping $f : \mathbb{R} \rightarrow \mathbb{R}, \forall x : f(x) = x + 2$.

Consider two vectors (numbers in this case) x_1 and x_2 . Then

$$f(x_1 + x_2) = x_1 + x_2 + 2$$

$$f(x_1) + f(x_2) = x_1 + 2 + x_2 + 2 = x_1 + x_2 + 4.$$

It follows that $f(x_1 + x_2) \neq f(x_1) + f(x_2)$. Hence, this mapping is not linear. Still, $f(0) = 2$ and

$$g(x) = f(x) - f(0) = x$$

which is a linear mapping. This means that $f(x)$ is an affine mapping.

Proposition

Let f be a linear mapping $E \rightarrow F$.

Let us indicate with \mathbf{o}_E and \mathbf{o}_F the null vectors of the vector spaces $(E, +, \cdot)$ and $(F, +, \cdot)$, respectively.

It follows that

$$f(\mathbf{o}_E) = \mathbf{o}_F.$$

Proof.

$$f(\mathbf{o}_E) = f(0\mathbf{o}_E) = 0f(\mathbf{o}_E) = \mathbf{o}_F.$$



Exercise 1: Linear and Affine Mappings

Which of these mappings are linear and which are affine (or both or neither)?:

⊙ $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = 2x^2 + 2.$

⊙ $f : \mathbb{R}^2 \rightarrow \mathbb{R}, f(x, y) = 2x - 3y.$

⊙ $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, f(x, y) = (2x - y + 1, 2y - x - 2).$

Exercise 1: *Solution*

To be completed.

Exercise 1: *Solution*

To be completed.

Exercise 1: *Solution*

To be completed.

Linear Mappings and Vector Spaces

Linear Mappings and Vector Spaces

Definition

Let f be a mapping $U \subset E \rightarrow F$.

The *image of U through f* , indicated with $f(U)$, is the set

$$f(U) = \{ \mathbf{w} \in F \mid \exists \mathbf{u} \in U \text{ such that } f(\mathbf{u}) = \mathbf{w} \}.$$

Theorem

Let $f : E \rightarrow F$ be a linear mapping and $(U, +, \cdot)$ be a vector subspace of $(E, +, \cdot)$.

It follows that the triple $(f(U), +, \cdot)$ is a vector subspace of $(F, +, \cdot)$.

Proof.

In order to prove that $(f(U), +, \cdot)$ is a vector subspace of $(F, +, \cdot)$ we have to show that the set $f(U)$ is closed with respect to the two composition laws.

By definition, the fact that a vector $\mathbf{w} \in f(U)$ means that $\exists \mathbf{v} \in U$ such that $f(\mathbf{v}) = \mathbf{w}$.

Thus, if we consider two vectors $\mathbf{w}, \mathbf{w}' \in f(U)$ then

$$\mathbf{w} + \mathbf{w}' = f(\mathbf{v}) + f(\mathbf{v}') = f(\mathbf{v} + \mathbf{v}').$$

Since for hypothesis $(U, +, \cdot)$ is a vector space, then $\mathbf{v} + \mathbf{v}' \in U$. Hence, $f(\mathbf{v} + \mathbf{v}') \in f(U)$. The set $f(U)$ is therefore closed with respect to the internal composition law. *continued...*

Proof.

Let us now consider a generic scalar $\lambda \in \mathbb{K}$ and calculate

$$\lambda \mathbf{w} = \lambda f(\mathbf{v}) = f(\lambda \mathbf{v}).$$

Since for hypothesis $(U, +, \cdot)$ is a vector space, then $\lambda \mathbf{v} \in U$.

Hence, $f(\lambda \mathbf{v}) \in f(U)$. The set $f(U)$ is closed with respect to the external composition law.

Since the set $f(U)$ is closed with respect to both the composition laws the triple $(f(U), +, \cdot)$ is a vector subspace of $(F, +, \cdot)$. □

Definition

Let f be a mapping $E \rightarrow F$ and $W \subset F$.

The *inverse image of W through f* , indicated with $f^{-1}(W)$, is a set defined as

$$f^{-1}(W) = \{\mathbf{u} \in E \mid f(\mathbf{u}) \in W\}.$$

Theorem

Let $f : E \rightarrow F$ be a linear mapping.

If $(W, +, \cdot)$ is a vector subspace of $(F, +, \cdot)$,

then $(f^{-1}(W), +, \cdot)$ is a vector subspace of $(E, +, \cdot)$.

Proof.

In order to prove that $(f^{-1}(W), +, \cdot)$ is a vector subspace of $(E, +, \cdot)$ we have to prove the closure of $f^{-1}(W)$ with respect to the two composition laws.

If a vector $\mathbf{v} \in f^{-1}(W)$ then $f(\mathbf{v}) \in W$.

We can write for the linearity of f

$$f(\mathbf{v} + \mathbf{v}') = f(\mathbf{v}) + f(\mathbf{v}').$$

Since $(W, +, \cdot)$ is a vector space, $f(\mathbf{v}) + f(\mathbf{v}') \in W$.

Hence, $f(\mathbf{v} + \mathbf{v}') \in W$ and $\mathbf{v} + \mathbf{v}' \in f^{-1}(W)$.

Thus, the set $f^{-1}(W)$ is closed with respect to the first composition law.

continued...

Proof.

Let us consider a generic scalar $\lambda \in \mathbb{K}$ and calculate

$$f(\lambda \mathbf{v}) = \lambda f(\mathbf{v}).$$

Since $(W, +, \cdot)$ is a vector space, $\lambda f(\mathbf{v}) \in W$. Since $f(\lambda \mathbf{v}) \in W$ the $\lambda \mathbf{v} \in f^{-1}(W)$. Thus, the set $f^{-1}(W)$ is closed with respect to the second composition law.

Hence, $(f^{-1}(W), +, \cdot)$ is a vector subspace of $(E, +, \cdot)$. □

Example

Let us consider the linear mapping $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$$f(x, y) = (x + y, x - y)$$

and the set $U = \{(x, y) \in \mathbb{R}^2 \mid 2x - y = 0\}$.

It can be easily checked that $(U, +, \cdot)$ is a vector space. This set can be represented by means of the vector

$$U = \alpha (1, 2)$$

with $\alpha \in \mathbb{R}$.

Continued...

Note: Here we use the shorthand $\alpha \mathbf{v}$ for $\{\alpha \mathbf{v} : \alpha \in \mathbb{R}\}$.

Example

Let us now calculate $f(U)$ by replacing (x, y) with $(\alpha, 2\alpha)$:

$$f(U) = (\alpha + 2\alpha, \alpha - 2\alpha) = (3\alpha, -\alpha) = \alpha(3, -1)$$

that is a line passing through the origin. Hence also

$$(f(U), +, \cdot)$$

is a vector space.

Summary and next lecture

Summary

- ⊙ Mappings
- ⊙ Linear Mappings
- ⊙ Linear Mappings and Vector Spaces

The next lecture

We will learn about Endomorphisms, Kernels and Rank-Nullity Theorem.