

# Metagenome Analysis of Premature Birth

Jaewoong Lee    Semin Lee

Department of Biomedical Engineering  
Ulsan National Institute of Science and Technology

*jwlee230@unist.ac.kr*

2022-01-27

# Overview

1 Introduction

2 Materials

3 Methods

4 Results

5 Discussion

6 References

# 1. Introduction

# Microbiome

- Microbiota: the microorganisms which live inside & on humans (Turnbaugh et al., 2007)
- Microbiome:  $10^{13}$  to  $10^{14}$  microorganisms whose collective genome (Gill et al., 2006)



Figure: Concept of a core human microbiome (Turnbaugh et al., 2007)

# rRNA

- Ribosomal RNA
- Well-known as a key to phylogeny (Olsen & Woese, 1993)

# Premature Birth (Preterm Birth; PTB)

PTB:

- ① PTB  $<$  37 GW (Gestational week)
- ② Normal  $\geq$  37 GW

Detailed PTB:

- ① Early PTB  $<$  34 GW
- ② 34 GW  $\leq$  Late PTB  $<$  37 GW
- ③ Normal  $\geq$  37 GW

(J. Tucker & McGuire, 2004; Voronkov, Solonovych, Liashenko, & Revenko, 2018)

## 2. Materials

# 16S rRNA Sequencing

**16S rRNA sequencing** is the *reference method* for bacterial taxonomy & identification (Mignard & Flandrois, 2006)

Three main reasons (Janda & Abbott, 2007):

- 16S rRNA exists in almost all bacteria
- Functions of the 16S rRNA has not changed over evolution.
- 16S rRNA is large enough for bioinformatics

# Data Composition


- JBNU/Helixco data
  - First data
  - Second data
  - Stool data

Table: Sample Information

Data	Participants	Samples	Remarks
First	24	107	-
Second	35	288	-
Third	10	106	-
Stool	63	126	Stool

### 3. Methods

# Qiime 2 Workflow



**Figure: QIIME 2 workflow (Bolyen et al., 2019; Mandal, Van Treuren, White, Eggesbø, et al., 2015; McDonald et al., 2012)**

## 4. Results

## 4. Results

### 4.1. Data Processing with Qiime

# Filtering with Quality Score


Drawback between:

- Longer sequence read
- Higher quality value


∴ Select the maximum length  $n$  where:

$$\begin{aligned} \forall n_i \in \{n_k | \text{MedianQualityScore} \geq 30\} \\ \exists! n \in \{n_i\} : n \geq n_i \end{aligned} \tag{1}$$

# Quality Score from JBU/Helixco Data



(a) Forward



(b) Reverse

Figure: Quality Score Plot

- $\ell_{Forward} = 300$
- $\ell_{Reverse} = 245$

# Denoising Techniques

- DADA2: Amplicon Sequence Variants (ASVs) (Callahan et al., 2016)
- Deblur: Operational Taxonomic Units (OTUs) (Amir et al., 2017)




Figure: Denoising Algorithms

# Taxonomy Classification

- Greengenes (GG) (DeSantis et al., 2006)
- SILVA (Pruesse et al., 2007; Quast et al., 2012)
- Human Oral Microbiome Database (HOMD)


“A **higher** performance at taxonomic levels above *genus level*;  
but performance appears to **drop** at *species level*” (Gihawi et al., 2019)

## 4. Results

### 4.2. Abundance Test with ANCOM

- Analysis with composition of microbiome (Mandal, Van Treuren, White, Eggesbø, et al., 2015)
- ANCOM detects significantly abundant taxa, while maintain high statistical power
- Find taxa that can divide each classes


# ANCOM with Detail PTB



(a) 1-day



(b) 3-day



(c) 5-day

Figure: ANCOM for Detail PTB from Neonatal Mouth with DADA2+HOMD

## 4. Results

### 4.3. Abundance Test with LefSe

- Linear discriminant analysis Effect Size (Segata et al., 2011)
- LefSe finds the features likely to explicate differences between groups

# LefSe with Detail PTB




Figure: LefSe for Detail PTB with Deblur + Silva

## 4. Results

### 4.4. Diversity Index

# Diversity Indices






Figure: Three dimensions of phylogenetic information (C. M. Tucker et al., 2017)

- A quantitative measure that shows richness, divergence, and regularity (C. M. Tucker et al., 2017)
- Alpha diversity indices: the richness of taxa **at a single community**
- Beta diversity indices: the taxonomic differentiation **between communities**


# Violin Plot with Alpha diversity I



(a) Mother Mouth




(b) Neonate 3-day




(c) Neonate 5-day

Figure: Detail premature & Faith's PD


# Violin Plot with Alpha diversity II



(a) Mother Mouth




(b) Neonate 3-day




(c) Neonate 5-day

Figure: Detail premature & Observed Features


# Violin Plot with Alpha diversity III



(a) Mother Mouth




(b) Neonate 3-day




(c) Neonate 5-day

Figure: Detail premature & Pielou Evenness


# Violin Plot with Alpha diversity IV



(a) Mother Mouth



(b) Neonate 3-day



(c) Neonate 5-day

Figure: Detail premature & Shannon Entropy

# Cluster map with Beta diversity I






Figure: Cluster map with Bray-Curtis distance index for DADA2


# Cluster map with Beta diversity II




(a) Mouth




(b) Cervix




(c) Vagina



(d) Neonate 1-day



(e) Neonate 3-day



(f) Neonate 5-day

Figure: Clustermap with Site separation

## 4. Results

### 4.5. Taxonomy Analyses

# Shared Taxonomy with Neonates & Mothers




Figure: Shared Taxonomy with Neonates & Mothers from DADA2 & HOMD

# Correlation between Taxonomy & Clinical data I




Figure: Pearson Correlation between taxonomy & clinical Data on DADA2 + HOMD

# Correlation between Taxonomy & Clinical data II




Figure: Person Correlation with Site separation

## 5. Discussion

## 6. References

## References I

- Amir, A., McDonald, D., Navas-Molina, J. A., Kopylova, E., Morton, J. T., Xu, Z. Z., ... others (2017). Deblur rapidly resolves single-nucleotide community sequence patterns. *MSystems*, 2(2).
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., ... Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using qiime 2. *Nature Biotechnology*, 37(8), 852-857. Retrieved from <https://doi.org/10.1038/s41587-019-0209-9> doi: 10.1038/s41587-019-0209-9
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). Dada2: high-resolution sample inference from illumina amplicon data. *Nature methods*, 13(7), 581–583.

## References II

- DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., ... Andersen, G. L. (2006). Greengenes, a chimera-checked 16s rrna gene database and workbench compatible with arb. *Applied and environmental microbiology*, 72(7), 5069–5072.
- Gihawi, A., Rallapalli, G., Hurst, R., Cooper, C. S., Leggett, R. M., & Brewer, D. S. (2019). Sepath: benchmarking the search for pathogens in human tissue whole genome sequence data leads to template pipelines. *Genome biology*, 20(1), 1–15.
- Gill, S. R., Pop, M., DeBoy, R. T., Eckburg, P. B., Turnbaugh, P. J., Samuel, B. S., ... Nelson, K. E. (2006). Metagenomic analysis of the human distal gut microbiome. *science*, 312(5778), 1355–1359.
- Janda, J. M., & Abbott, S. L. (2007). 16s rrna gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. *Journal of clinical microbiology*, 45(9), 2761–2764.

## References III

- Mandal, S., Van Treuren, W., White, R. A., Eggesbø, M., Knight, R., & Peddada, S. D. (2015). Analysis of composition of microbiomes: a novel method for studying microbial composition. *Microbial ecology in health and disease*, 26(1), 27663.
- Mandal, S., Van Treuren, W., White, R. A., Eggesbø, M., Knight, R., & Peddada, S. D. (2015). Analysis of composition of microbiomes: a novel method for studying microbial composition. *Microbial ecology in health and disease*, 26(1), 27663. doi: 10.3402/mehd.v26.27663
- McDonald, D., Clemente, J. C., Kuczynski, J., Rideout, J. R., Stombaugh, J., Wendel, D., ... Caporaso, J. G. (2012). The biological observation matrix (biom) format or: how i learned to stop worrying and love the ome-ome. *GigaScience*, 1(1), 7. doi: 10.1186/2047-217X-1-7

## References IV

- Mignard, S., & Flandrois, J.-P. (2006). 16s rrna sequencing in routine bacterial identification: a 30-month experiment. *Journal of microbiological methods*, 67(3), 574–581.
- Olsen, G. J., & Woese, C. R. (1993). Ribosomal rna: a key to phylogeny. *The FASEB journal*, 7(1), 113–123.
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). Silva: a comprehensive online resource for quality checked and aligned ribosomal rna sequence data compatible with arb. *Nucleic acids research*, 35(21), 7188–7196.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... Glöckner, F. O. (2012). The silva ribosomal rna gene database project: improved data processing and web-based tools. *Nucleic acids research*, 41(D1), D590–D596.

## References V

- Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W. S., & Huttenhower, C. (2011). Metagenomic biomarker discovery and explanation. *Genome biology*, 12(6), 1–18.
- Tucker, C. M., Cadotte, M. W., Carvalho, S. B., Davies, T. J., Ferrier, S., Fritz, S. A., ... others (2017). A guide to phylogenetic metrics for conservation, community ecology and macroecology. *Biological Reviews*, 92(2), 698–715.
- Tucker, J., & McGuire, W. (2004). Epidemiology of preterm birth. *Bmj*, 329(7467), 675–678.
- Turnbaugh, P. J., Ley, R. E., Hamady, M., Fraser-Liggett, C. M., Knight, R., & Gordon, J. I. (2007). The human microbiome project. *Nature*, 449(7164), 804–810.

## References VI

Voronkov, L., Solonovych, A., Liashenko, A., & Revenko, I. (2018). Prognostic value of cognitive tests and their combination in patients with chronic heart failure and reduced left ventricular ejection fraction. *Eureka: health sciences*(6), 36–45.