

Metagenome Analysis of Premature Birth

Jaewoong Lee Semin Lee

Department of Biomedical Engineering
Ulsan National Institute of Science and Technology

jwlee230@unist.ac.kr

2021-05-26

Overview

- 1 Introduction
- 2 Materials
- 3 Methods
- 4 Results

Introduction

Microbiome

- Microbiota: the microorganisms which live inside & on humans (Turnbaugh et al., 2007)
- Microbiome: 10^{13} to 10^{14} microorganisms whose collective genome (Gill et al., 2006)



Figure: Concept of a core human microbiome (Turnbaugh et al., 2007)

- Ribosomal RNA
- Well-known as a key to phylogeny (Olsen & Woese, 1993)

Premature Birth (Preterm Birth)



Figure: Definitions of Premature (J. Tucker & McGuire, 2004)

∴ Hence, in this study,

- Premature: < 37 weeks
- Normal: ≥ 37 weeks

Materials

16S rRNA Sequencing

16S rRNA sequencing is the *reference method* for bacterial taxonomy & identification (Mignard & Flandrois, 2006)

Three main reasons (Janda & Abbott, 2007):

- 16S rRNA exists in almost all bacteria
- Functions of the 16S rRNA has not changed over time
- 16S rRNA is large enough for bioinformatics

Train/Test Data vs. Validate Data

- JBNU/Helixco data
 - First data
 - Second data
 - Stool data

Table: Sample Information

Data	Participants	Samples	Remarks
First	24	107	-
Second	35	288	-
Stool	63	126	Stool

Methods

Methods

Qiime 2 Workflow

Qiime 2 Workflow




Figure: QIIME 2 workflow (Bolyen et al., 2019; Mandal, Van Treuren, White, Eggesbø, et al., 2015; McDonald et al., 2012)

Filtering with Quality Score

Drawback between:

- Longer sequence read
- Higher quality value

∴ Select the maximum length n where:

$$\begin{aligned} \forall n_i \in \{n_k | \text{MedianQualityScore} \geq 30\} \\ \exists ! n \in \{n_i\} : n \geq n_i \end{aligned} \tag{1}$$

Denoising Techniques

- DADA2: Amplicon Sequence Variants (ASVs) (Callahan et al., 2016)
- Deblur: Operational Taxonomic Units (OTUs) (Amir et al., 2017)




Figure: Denoising Algorithms

Taxonomy Classification

- Greengenes (GG) (DeSantis et al., 2006)
- SILVA (Pruesse et al., 2007; Quast et al., 2012)




Figure: Taxonomy Classifications

“A **higher** performance at taxonomic levels above *genus level*;
but performance appears to **drop** at *species level*” (Gihawi et al., 2019)

Merging Denoising/Taxonomy

Merging multiple IDs (ASVs or OTUs) into one, which have

- Different IDs
- Identified as same taxonomy




Figure: Example Diagram for Merging Denoising/Taxonomy

Methods

Abundance Test

- Analysis of composition of microbiome (Mandal, Van Treuren, White, Eggesbø, et al., 2015)
- ANCOM detects significantly abundant taxa, while maintain high statistical power
- Find taxa that can divide each classes

Methods

Diversity Indices

Diversity Indices




Figure: Three dimensions of phylogenetic information (C. M. Tucker et al., 2017)

- A quantitative measure that shows richness, divergence, and regularity (C. M. Tucker et al., 2017)
- Alpha diversity indices: the richness of taxa **at a single community**
- Beta diversity indices: the taxonomic differentiation **between communities**

Alpha Diversity Indices

- Evenness index
- Faith's Phylogenetic Diversity (Faith PD) index
- Oberseved Features index
- Shannon's Diversity index

Beta Diversity Indices

- Bray-Curtis distance index
- Jaccard distance index
- Unweighted UniFrac distance index
- Weighted UniFrac distance index

Methods

Miscellaneous

t-distributed Stochastic Neighbor Embedding (t-SNE)



Figure: t-SNE with handwritten data (Maaten & Hinton, 2008)


- Pandas (McKinney et al., 2011)
- Scikit-Learn (Pedregosa et al., 2011)
- SciPy (Virtanen et al., 2020)
- Matplotlib (Hunter, 2007)
- Seaborn (Waskom et al., 2020)
- Statannot

Results


Results

Filtering Results

Quality Score from JBNU/Helixco Data



(a) Forward



(b) Reverse

Figure: Quality Score Plot

- $\ell_{Forward} = 300$
- $\ell_{Reverse} = 245$

Results

t-SNE with Clinical Information

Workflow for t-SNE with Site/Premature Information



Figure: Workflow of t-SNE for Site/Premature Information


Results

Bacterial Abundance Test with ANCOM


- Analysis of composition of microbiomes
- ANCOM can be used for analyzing the composition of microbiomes in multiple populations (Mandal, Van Treuren, White, Eggesbø, et al., 2015)
- Differential abundance testing
- ① clr: Centered log(*Ratio*)
- ② W: a count of the number of sub-hypothesis which have passed for given species

- Site where get the microbiome
- Premature – Before 37 weeks and After 37 weeks
- Detailed Premature – Before 34 weeks, After 37 weeks, and the other
- C-section
- PROM – Premature rupture of membranes
- Using Steroid?
- Using anti-biotic?

ANCOM with Detailed Premature



(a) DADA2 + GG




(b) Deblur + GG


Figure: ANCOM results with Detailed Premature

- *Ureaplasma* genus
- *Aerococcus* genus


ANCOM with Site



(a) DADA2 + GG



(b) DADA2 + Silva



(c) Deblur + GG

Figure: ANCOM results with Site

ANCOM with PROM






Figure: ANCOM result with PROM (DADA2 + GG)

- *Campylobacteraceae* genus *Campylobacter* species

ANCOM with Using Steroid



(a) DADA2 + GG



(b) Deblur + GG

Figure: ANCOM results with Using Steroid


- *Ureaplasma* genus
- *Aerococcus* genus

Results


Rarefaction

Rarefaction?

Rarefaction from First Data



(a) DADA2



(b) Deblur

Figure: Rarefaction from the Data

- $\min(\ell_{DADA2})$: 1046
- $\min(\ell_{Deblur})$: 4864

Results

Alpha-Diversity

Alpha-Diversity

Results

Beta-Diversity

Beta-Diversity

Results

Classification

Workflow for Classification




Figure: Workflow with Classification

Random Forest Classifier I

References I

- Amir, A., McDonald, D., Navas-Molina, J. A., Kopylova, E., Morton, J. T., Xu, Z. Z., ... others (2017). Deblur rapidly resolves single-nucleotide community sequence patterns. *MSystems*, 2(2).
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., ... Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using qiime 2. *Nature Biotechnology*, 37(8), 852-857. Retrieved from <https://doi.org/10.1038/s41587-019-0209-9> doi: 10.1038/s41587-019-0209-9
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). Dada2: high-resolution sample inference from illumina amplicon data. *Nature methods*, 13(7), 581-583.

- DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., ... Andersen, G. L. (2006). Greengenes, a chimera-checked 16s rRNA gene database and workbench compatible with arb. *Applied and environmental microbiology*, 72(7), 5069–5072.
- Dominguez-Bello, M. G., De Jesus-Laboy, K. M., Shen, N., Cox, L. M., Amir, A., Gonzalez, A., ... others (2016). Partial restoration of the microbiota of cesarean-born infants via vaginal microbial transfer. *Nature medicine*, 22(3), 250.
- Fettweis, J. M., Serrano, M. G., Brooks, J. P., Edwards, D. J., Girerd, P. H., Parikh, H. I., ... others (2019). The vaginal microbiome and preterm birth. *Nature medicine*, 25(6), 1012–1021.

References III

- Gihawi, A., Rallapalli, G., Hurst, R., Cooper, C. S., Leggett, R. M., & Brewer, D. S. (2019). Sepath: benchmarking the search for pathogens in human tissue whole genome sequence data leads to template pipelines. *Genome biology*, 20(1), 1–15.
- Gill, S. R., Pop, M., DeBoy, R. T., Eckburg, P. B., Turnbaugh, P. J., Samuel, B. S., ... Nelson, K. E. (2006). Metagenomic analysis of the human distal gut microbiome. *science*, 312(5778), 1355–1359.
- Hunter, J. D. (2007). Matplotlib: A 2d graphics environment. *Computing in science & engineering*, 9(3), 90–95.
- Janda, J. M., & Abbott, S. L. (2007). 16s rrna gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. *Journal of clinical microbiology*, 45(9), 2761–2764.
- Maaten, L. v. d., & Hinton, G. (2008). Visualizing data using t-sne. *Journal of machine learning research*, 9(Nov), 2579–2605.

- Mandal, S., Van Treuren, W., White, R. A., Eggesbø, M., Knight, R., & Peddada, S. D. (2015). Analysis of composition of microbiomes: a novel method for studying microbial composition. *Microbial ecology in health and disease*, 26(1), 27663.
- Mandal, S., Van Treuren, W., White, R. A., Eggesbø, M., Knight, R., & Peddada, S. D. (2015). Analysis of composition of microbiomes: a novel method for studying microbial composition. *Microbial ecology in health and disease*, 26(1), 27663. doi: 10.3402/mehd.v26.27663
- MARTIUS, J., KROHN, M. A., HILLIER, S. L., STAMM, W. E., HOLMES, K. K., & ESCHENBACH, D. A. (1988). Relationships of vaginal lactobacillus species, cervical chlamydia trachomatis, and bacterial vaginosis to preterm birth. *Obstetrics & Gynecology*, 71(1), 89–95.

- McDonald, D., Clemente, J. C., Kuczynski, J., Rideout, J. R., Stombaugh, J., Wendel, D., ... Caporaso, J. G. (2012). The biological observation matrix (biom) format or: how i learned to stop worrying and love the ome-ome. *GigaScience*, 1(1), 7. doi: 10.1186/2047-217X-1-7
- McKinney, W., et al. (2011). pandas: a foundational python library for data analysis and statistics. *Python for High Performance and Scientific Computing*, 14(9).
- Mignard, S., & Flandrois, J.-P. (2006). 16s rna sequencing in routine bacterial identification: a 30-month experiment. *Journal of microbiological methods*, 67(3), 574–581.
- Olsen, G. J., & Woese, C. R. (1993). Ribosomal rna: a key to phylogeny. *The FASEB journal*, 7(1), 113–123.

References VI

- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., ... others (2011). Scikit-learn: Machine learning in python. *the Journal of machine Learning research*, 12, 2825–2830.
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). Silva: a comprehensive online resource for quality checked and aligned ribosomal rna sequence data compatible with arb. *Nucleic acids research*, 35(21), 7188–7196.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., ... Glöckner, F. O. (2012). The silva ribosomal rna gene database project: improved data processing and web-based tools. *Nucleic acids research*, 41(D1), D590–D596.
- Stafford, G. P., Parker, J. L., Amabebe, E., Kistler, J., Reynolds, S., Stern, V., ... Anumba, D. O. (2017). Spontaneous preterm birth is associated with differential expression of vaginal metabolites by lactobacilli-dominated microflora. *Frontiers in physiology*, 8, 615.

- Tabatabaei, N., Eren, A., Barreiro, L., Yotova, V., Dumaine, A., Allard, C., & Fraser, W. (2019). Vaginal microbiome in early pregnancy and subsequent risk of spontaneous preterm birth: a case–control study. *BJOG: An International Journal of Obstetrics & Gynaecology*, 126(3), 349–358.
- Tucker, C. M., Cadotte, M. W., Carvalho, S. B., Davies, T. J., Ferrier, S., Fritz, S. A., ... others (2017). A guide to phylogenetic metrics for conservation, community ecology and macroecology. *Biological Reviews*, 92(2), 698–715.
- Tucker, J., & McGuire, W. (2004). Epidemiology of preterm birth. *Bmj*, 329(7467), 675–678.
- Turnbaugh, P. J., Ley, R. E., Hamady, M., Fraser-Liggett, C. M., Knight, R., & Gordon, J. I. (2007). The human microbiome project. *Nature*, 449(7164), 804–810.

References VIII

- Usui, R., Ohkuchi, A., Matsubara, S., Izumi, A., Watanabe, T., Suzuki, M., & Minakami, H. (2002). Vaginal lactobacilli and preterm birth. *Journal of perinatal medicine*, 30(6), 458–466.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., ... others (2020). Scipy 1.0: fundamental algorithms for scientific computing in python. *Nature methods*, 17(3), 261–272.
- Waskom, M., Botvinnik, O., Ostblom, J., Gelbart, M., Lukauskas, S., Hobson, P., ... Brian (2020, April). *mwaskom/seaborn: v0.10.1 (april 2020)*. Zenodo. Retrieved from <https://doi.org/10.5281/zenodo.3767070> doi: 10.5281/zenodo.3767070