Periodontitis

Seunghoon Kim Jaewoong Lee Semin Lee

Ulsan National Institute of Science and Technology *jwlee230@unist.ac.kr*

2020-11-24

Overview

- Introduction
- 2 Materials
- Methods
- Results
- 5 Discussion References

Introduction

Microbiome

- Microbiota: the micro-organisms which live inside & on humans (Turnbaugh et al., 2007)
- Microbiome: about 10^{13} micro-organisms whose which collective genome (Gill et al., 2006)



Figure: Concept of a core human microbiome (Turnbaugh et al., 2007)

rRNA

- Ribosomal RNA
- Well-known as a key to phylogeny (Olsen & Woese, 1993)

Periodontitis (Periodontal disease)

- CAL (Clinical Attachment Loss) & BL (Bone Loss) (Flemmig, 1999)
- Risk Factors (Van Dyke & Dave, 2005)
 - Smoking
 - 2 Diabetes
 - Genetic factor
 - 4 Host response

Materials

16S rRNA Sequencing

- 100 Healthy people
- 50 Chronic periodontitis Early
- 50 Chronic periodontitis Moderate
- 50 Chronic periodontitis Severe

Methods

Qiime2 Workflow




Figure: Qiime2 Workflow (Bolyen et al., 2018, 2019)

Denoising techniques

- DADA2: Amplicon Sequence Variants (ASVs) (Callahan et al., 2016)
- Deblur: Operational Taxonomic Units (OTUs) (Amir et al., 2017)




Figure: Denoising Techniques

Taxonomy Classification

- Greengenes (GG) (DeSantis et al., 2006)
- SILVA (Pruesse et al., 2007)





Figure: Taxonomy Classification

"A **higher** performance at taxonomic levels above *genus* level; but performance appears to drop at *species* level" (Gihawi et al., 2019)

Results

Quality Filter



 \therefore Maximum Sequence Length n = 250.

Discussion

References I

- Amir, A., McDonald, D., Navas-Molina, J. A., Kopylova, E., Morton, J. T., Xu, Z. Z., . . . others (2017). Deblur rapidly resolves single-nucleotide community sequence patterns. *MSystems*, *2*(2).
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C., Al-Ghalith, G. A., . . . others (2018). *Qiime 2: Reproducible, interactive, scalable, and extensible microbiome data science* (Tech. Rep.). PeerJ Preprints.
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., . . . others (2019). Reproducible, interactive, scalable and extensible microbiome data science using qiime 2. *Nature biotechnology*, *37*(8), 852–857.
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). Dada2: high-resolution sample inference from illumina amplicon data. *Nature methods*, *13*(7), 581–583.

References II

- DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., . . . Andersen, G. L. (2006). Greengenes, a chimera-checked 16s rrna gene database and workbench compatible with arb. *Applied and environmental microbiology*, 72(7), 5069–5072.
- Flemmig, T. F. (1999). Periodontitis. *Annals of Periodontology*, 4(1), 32–37.
- Gihawi, A., Rallapalli, G., Hurst, R., Cooper, C. S., Leggett, R. M., & Brewer, D. S. (2019). Sepath: benchmarking the search for pathogens in human tissue whole genome sequence data leads to template pipelines. *Genome biology*, 20(1), 1–15.
- Gill, S. R., Pop, M., DeBoy, R. T., Eckburg, P. B., Turnbaugh, P. J., Samuel, B. S., ... Nelson, K. E. (2006). Metagenomic analysis of the human distal gut microbiome. *science*, *312*(5778), 1355–1359.

References III

- Olsen, G. J., & Woese, C. R. (1993). Ribosomal rna: a key to phylogeny. *The FASEB journal*, 7(1), 113–123.
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). Silva: a comprehensive online resource for quality checked and aligned ribosomal rna sequence data compatible with arb. *Nucleic acids research*, *35*(21), 7188–7196.
- Turnbaugh, P. J., Ley, R. E., Hamady, M., Fraser-Liggett, C. M., Knight, R., & Gordon, J. I. (2007). The human microbiome project. *Nature*, 449(7164), 804–810.
- Van Dyke, T. E., & Dave, S. (2005). Risk factors for periodontitis. Journal of the International Academy of Periodontology, 7(1), 3.