

# Periodontitis

Seunghoon Kim    Jaewoong Lee    Semin Lee

Ulsan National Institute of Science and Technology

*jwlee230@unist.ac.kr*

2020-12-10

# Overview

1 Introduction

2 Materials

3 Methods

4 Results

5 Discussion

# Introduction

# Microbiome

- Microbiota: the micro-organisms which live inside & on humans (Turnbaugh et al., 2007)
- Microbiome: about  $10^{13}$  micro-organisms whose collective genome (Gill et al., 2006)



Figure: Concept of a core human microbiome (Turnbaugh et al., 2007)

# rRNA

- Ribosomal RNA
- Well-known as a key to phylogeny (Olsen & Woese, 1993)

# Periodontitis (Periodontal disease)

- Clinical Attachment Loss & Bone Loss (Flemmig, 1999)
- Risk Factors (Van Dyke & Dave, 2005)
  - ① Smoking
  - ② Diabetes
  - ③ Genetic factor
  - ④ Host response

# Materials

# 16S rRNA Sequencing

- 100 Healthy people
- 50 Chronic periodontitis – Early
- 50 Chronic periodontitis – Moderate
- 50 Chronic periodontitis – Severe

# Methods

# QIIME2 Workflow




Figure: QIIME2 Workflow (Bolyen et al., 2019, 2018)

# Denoising techniques

- DADA2: Amplicon Sequence Variants (ASVs) (Callahan et al., 2016)
- Deblur: Operational Taxonomic Units (OTUs) (Amir et al., 2017)



Figure: Denoising Techniques

# Taxonomy Classification

- Greengenes (GG) (DeSantis et al., 2006)
- SILVA (Pruesse et al., 2007)
- Human Oral Microbiome Database (HOMD) (Chen et al., 2010)




Figure: Taxonomy Classification

"A **higher** performance at taxonomic levels above *genus* level; but performance appears to drop at *species* level" (Gihawi et al., 2019)

# Merging Denosing and Taxonomy Classification

Merging multiple IDs (ASVs and OTUs) into one, which have:

- Different IDs.
- Identified as same taxonomy.




Figure: Example Diagram for Merging Denosing and Taxonomy Classification

# Rarefaction

- a statistical method of estimating the number of species expected in **a random sample** which taken from a collection (James & Rathbun, 1981)
- allows comparisons of **the species richness** among communities
- a good choice for **normalization** (Weiss et al., 2017)

# Alpha- & Beta-diversity

- Alpha-diversity: the richness of taxa **at a single community**
- Beta-diversity: the taxonomic differentiation **between communities**




Figure: Three Dimensions of Phylogenetic Information (Tucker et al., 2017)

# Alpha-diversity

- Evenness: a measurement of diversity in different type at community (Pielou, 1966)
- Faith's Phylogenetic Diversity: a qualitative measurement of community richness which prioritizes species conservation, incorporates with taxic diversity (Faith, 1992)
- Observed Features: a number of observed taxa
- Shannon's diversity index: a significant aspect of community richness (Shannon, 1948)

# Beta-diversity

- Bray-Curtis distance: a quantitative measurement of dissimilarity among communities (Sørensen, 1948)
- Jaccard distance: a measurement of local distribution among communities (Jaccard, 1912)
- Unweighted UniFrac distance: a qualitative measurement of phylogenetic distances (McDonald et al., 2018)
- Weighted UniFrac distance: a quantitative measurement of phylogenetic distances (McDonald et al., 2018)

# ANCOM

- Analysis of composition of microbiomes
- ANCOM can be used for analyzing the composition of microbiomes in multiple populations (Mandal et al., 2015)
- Differential abundance testing



Figure: Example ANCOM Volcano Plot (Bolyen et al., 2019, 2018)

- clr: Centered log Ratio
- W: a count of the number of sub-hypothesis which have passed for given species

# Python Packages

- Pandas (McKinney et al., 2011)
- Scikit-learn (Pedregosa et al., 2011)
- Matplotlib (Hunter, 2007; Barrett, Hunter, Miller, Hsu, & Greenfield, 2005)
- Seaborn (Waskom & the seaborn development team, 2020)

# t-SNE

- t-distributed stochastic neighbor embedding
- reveals high-dimensional data a location in two-dimensional map  
(Maaten & Hinton, 2008)




Figure: Visualization by t-SNE (Maaten & Hinton, 2008)

# Classification I



Figure: Workflow of Classification

## Classification Metrics:

- Accuracy
- Balanced Accuracy
- Sensitivity
- Specificity
- Precision

# Classification II





Figure: Deciding the Best Features

# Results

# Quality Filter



(a) Forward Reads


(b) Reverse Reads

Figure: Sequence Quality Plot


∴ Maximum Sequence Length  $n_{forward} = 300$ ,  $n_{reverse} = 265$

∴ The longest length which has sequence quality  $\geq 30$  at middle.

# Rarefaction



(a) DADA2




(b) Deblur

Figure: Frequency per sample

$\therefore$  p-sampling-depth  $n_{DADA2} = 3786$  and  $n_{Deblur} = 7253$


# Alpha-diversity I




(a) Evenness ( $p < 0.01$ )



(b) Faith PD ( $p < 10^{-6}$ )




(c) Observed features ( $p < 10^{-3}$ )




(d) Shannon's diversity ( $p > 0.05$ )

**Figure:** Alpha Diversity from DADA2 with Kruskal-Wallis among All Groups


# Alpha-diversity II



(a) Evenness ( $p < 0.05$ )



(b) Faith PD ( $p < 10^{-18}$ )



(c) Observed features ( $p < 10^{-12}$ ) (d) Shannon's diversity ( $p < 10^{-4}$ )




Figure: Alpha Diversity from Deblur with Kruskal-Wallis among All Groups

# Beta-diversity I




Figure: Bray-Curtis Distance with DADA2

# Beta-diversity II




Figure: Jaccard Distance with DADA2

# Beta-diversity III






Figure: Unweighted Unifrac Distance with DADA2


# Beta-diversity IV




(a) Healthy



(b) Early



(c) Moderate



(d) Severe

Figure: Weighted Unifrac Distance with DADA2

# Beta-diversity V




Figure: Bray-Curtis Distance with Deblur

# Beta-diversity VI




Figure: Jaccard Distance with Deblur

# Beta-diversity VII




Figure: Unweighted Unifrac Distance with Deblur

# Beta-diversity VIII




Figure: Weighted Unifrac Distance with Deblur


# ANCOM I



(a) GG (15 Taxa)




(b) SILVA (23 Taxa)




(c) HOMD (20 Taxa)


Figure: ANCOM Volcano Plot with DADA2



(a) GG (27 Taxa)




(b) SILVA (20 Taxa)



(c) HOMD (28 Taxa)


Figure: ANCOM Volcano Plot with Deblur

# t-SNE with Whole Microbiome I



**Figure:** t-SNE Plot with Whole Microbiome from DADA2

# t-SNE with Whole Microbiome II




(a) GG (232 Taxa)

(b) SILVA (414 Taxa)


(c) HOMD (235 Taxa)

Figure: t-SNE Plot with Whole Microbiome from Deblur


# t-SNE with ANCOM Selected I



(a) GG (15 Taxa)



(b) SILVA (23 Taxa)



(c) HOMD (20 Taxa)

**Figure:** t-SNE Plot with ANCOM Selected from DADA2

# t-SNE with ANCOM Selected II




Figure: t-SNE Plot with ANCOM Selected from Deblur

# Random Forest Classifier I




Figure: Random Forest Classifier Workflow

**DADA2 + SILVA** gives the best result with balanced accuracy.

# Random Forest Classifier II

Table: Features Order by Importances

Order	Taxonomy (Genus [Species])
0	<i>Actinomyces</i>
1	<i>Actinomyces Schaalia-odontolytica</i>
2	<i>Prevotella Prevotella-intermedia</i>
3	<i>Filifactor Filifactor-alocis</i>
4	<i>Oribacterium</i>
5	<i>Tannerella Tannerella-forsythia</i>

# Random Forest Classifier III




Figure: Metrics by Feature Count

# Random Forest Classifier IV






Figure: Random Forest Tree Plot

# Random Forest Classifier V



(a) *Actinomyces*



(b) *Actinomyces Schaalia-odontolytica*

Figure: Violin Plot by Features

# Random Forest Classifier – Merge (Healthy+Early) I




Figure: Random Forest Classifier with Merging Workflow

**DADA2 + GG** gives the best result with balanced accuracy.

# Random Forest Classifier – Merge (Healthy+Early) II

Table: Features Order by Importances

Order	Taxonomy (Genus [Species])
0	<i>Actinomyces</i>
1	<i>Filifactor</i>
2	<i>Prevotella intermedia</i>

# Random Forest Classifier – Merge (Healthy+Early) III




Figure: Metrics by Feature Count

# Random Forest Classifier – Merge (Healthy+Early) IV






Figure: Random Forest Tree Plot

# Random Forest Classifier – Merge (Healthy+Early) V



(a) *Actinomyces*



(b) *Filifactor*

Figure: Violin Plot by Features

# Random Forest Classifier – Merge (Moderate + Severe) I




Figure: Random Forest Classifier with Merging Workflow

**DADA2 + GG** gives the best result with balanced accuracy.

# Random Forest Classifier – Merge (Moderate + Severe) II

Table: Features Order by Importances

Order	Taxonomy (Genus [Species])
0	<i>Actinomyces</i>
1	<i>Filifactor</i>
2	<i>Prevotella intermedia</i>

# Random Forest Classifier – Merge (Moderate + Severe) III




Figure: Metrics by Feature Count

# Random Forest Classifier – Merge (Moderate + Severe) IV






Figure: Random Forest Tree Plot

## Random Forest Classifier – Merge (Moderate + Severe) V



(a) *Actinomyces*



(b) *Filifactor*

## Figure: Violin Plot by Features

# Random Forest Classifier – Merge (H + E) & (M + S) I




Figure: Random Forest Classifier with Merging Workflow

**Deblur + SILVA** gives the best result with balanced accuracy.

# Random Forest Classifier – Merge (H + E) & (M + S) II

Table: Features Order by Importances

Order	Taxonomy (Genus [Species])
0	<i>Porphyromonas gingivalis</i>
1	<i>Treponema denticola</i>
2	<i>Treponema medium</i>
3	<i>Metamycoplasmfaucium</i>
4	<i>Prevotella intermedia</i>
5	<i>Streptococcus constellatus</i>
6	<i>Eubacterium nodatum group Eubacterium nodatum</i>
7	<i>Treponema uncultured bacterium</i>
...	...

# Random Forest Classifier – Merge (H + E) & (M + S) III






Figure: Metrics by Feature Count

Random Forest Classifier – Merge (H + E) & (M + S) IV




## Figure: Random Forest Tree Plot

# Random Forest Classifier – Merge (H + E) & (M + S) V



(a) *Porphyromonas gingivalis*



(b) *Treponema denticola*

Figure: Violin Plot by Features

## Discussion

# *Actinomyces* Genus

- *Actinomyces*: a deceptive infection of oral cavity (Thukral, Srivastav, Mathur, Barodiya, & Srivastav, 2017)
- Prevalence of *Actinomyces* spp. in patients with chronic periodontitis (Vielkind, Jentsch, Eschrich, Rodloff, & Stingu, 2015)

# *Filifactor* Genus

- *Filifactor alocis* – a new emerging periodontal pathogen (Aruni et al., 2015)
- *In Situ* Anabolic Activity of Periodontal Pathogens *Porphyromonas gingivalis* and *Filifactor alocis* in Chronic Periodontitis (Spooner et al., 2016)

# *Porphyromonas* Genus

- Antibody responses to *Porphyromonas gingivalis* (*P. gingivalis*) in subjects with rheumatoid arthritis and periodontitis (Mikuls et al., 2009)
- Contribution of *Porphyromonas gingivalis* lipopolysaccharide to periodontitis (Jain & Darveau, 2010)

# References I

- Amir, A., McDonald, D., Navas-Molina, J. A., Kopylova, E., Morton, J. T., Xu, Z. Z., . . . others (2017). Deblur rapidly resolves single-nucleotide community sequence patterns. *MSystems*, 2(2).
- Aruni, A. W., Mishra, A., Dou, Y., Chioma, O., Hamilton, B. N., & Fletcher, H. M. (2015). Filifactor alocis—a new emerging periodontal pathogen. *Microbes and Infection*, 17(7), 517–530.
- Barrett, P., Hunter, J., Miller, J. T., Hsu, J.-C., & Greenfield, P. (2005). matplotlib—a portable python plotting package. In *Astronomical data analysis software and systems xiv* (Vol. 347, p. 91).
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C., Al-Ghalith, G. A., . . . others (2018). *Qiime 2: Reproducible, interactive, scalable, and extensible microbiome data science* (Tech. Rep.). PeerJ Preprints.

## References II

- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A., . . . others (2019). Reproducible, interactive, scalable and extensible microbiome data science using qiime 2. *Nature biotechnology*, 37(8), 852–857.
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). Dada2: high-resolution sample inference from illumina amplicon data. *Nature methods*, 13(7), 581–583.
- Chen, T., Yu, W.-H., Izard, J., Baranova, O. V., Lakshmanan, A., & Dewhirst, F. E. (2010). The human oral microbiome database: a web accessible resource for investigating oral microbe taxonomic and genomic information. *Database*, 2010.

## References III

- DeSantis, T. Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E. L., Keller, K., ... Andersen, G. L. (2006). Greengenes, a chimera-checked 16s rrna gene database and workbench compatible with arb. *Applied and environmental microbiology*, 72(7), 5069–5072.
- Faith, D. P. (1992). Conservation evaluation and phylogenetic diversity. *Biological conservation*, 61(1), 1–10.
- Flemmig, T. F. (1999). Periodontitis. *Annals of Periodontology*, 4(1), 32–37.
- Gihawi, A., Rallapalli, G., Hurst, R., Cooper, C. S., Leggett, R. M., & Brewer, D. S. (2019). Sepath: benchmarking the search for pathogens in human tissue whole genome sequence data leads to template pipelines. *Genome biology*, 20(1), 1–15.

## References IV

- Gill, S. R., Pop, M., DeBoy, R. T., Eckburg, P. B., Turnbaugh, P. J., Samuel, B. S., ... Nelson, K. E. (2006). Metagenomic analysis of the human distal gut microbiome. *science*, 312(5778), 1355–1359.
- Hunter, J. D. (2007). Matplotlib: A 2d graphics environment. *Computing in science & engineering*, 9(3), 90–95.
- Jaccard, P. (1912). The distribution of the flora in the alpine zone. 1. *New phytologist*, 11(2), 37–50.
- Jain, S., & Darveau, R. P. (2010). Contribution of porphyromonas gingivalis lipopolysaccharide to periodontitis. *Periodontology 2000*, 54(1), 53.
- James, F. C., & Rathbun, S. (1981). Rarefaction, relative abundance, and diversity of avian communities. *The Auk*, 98(4), 785–800.
- Maaten, L. v. d., & Hinton, G. (2008). Visualizing data using t-sne. *Journal of machine learning research*, 9(Nov), 2579–2605.

## References V

- Mandal, S., Van Treuren, W., White, R. A., Eggesbø, M., Knight, R., & Peddada, S. D. (2015). Analysis of composition of microbiomes: a novel method for studying microbial composition. *Microbial ecology in health and disease*, 26(1), 27663.
- McDonald, D., Vázquez-Baeza, Y., Koslicki, D., McClelland, J., Reeve, N., Xu, Z., ... Knight, R. (2018). Striped unifrac: enabling microbiome analysis at unprecedented scale. *Nature methods*, 15(11), 847–848.
- McKinney, W., et al. (2011). pandas: a foundational python library for data analysis and statistics. *Python for High Performance and Scientific Computing*, 14(9).
- Mikuls, T. R., Payne, J. B., Reinhardt, R. A., Thiele, G. M., Maziarz, E., Cannella, A. C., ... O'Dell, J. R. (2009). Antibody responses to *porphyromonas gingivalis* (*p. gingivalis*) in subjects with rheumatoid arthritis and periodontitis. *International immunopharmacology*, 9(1), 38–42.

## References VI

- Olsen, G. J., & Woese, C. R. (1993). Ribosomal rna: a key to phylogeny. *The FASEB journal*, 7(1), 113–123.
- Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., ... others (2011). Scikit-learn: Machine learning in python. *the Journal of machine Learning research*, 12, 2825–2830.
- Pielou, E. C. (1966). The measurement of diversity in different types of biological collections. *Journal of theoretical biology*, 13, 131–144.
- Pruesse, E., Quast, C., Knittel, K., Fuchs, B. M., Ludwig, W., Peplies, J., & Glöckner, F. O. (2007). Silva: a comprehensive online resource for quality checked and aligned ribosomal rna sequence data compatible with arb. *Nucleic acids research*, 35(21), 7188–7196.
- Shannon, C. E. (1948). A mathematical theory of communication. *The Bell system technical journal*, 27(3), 379–423.

## References VII

- Spooner, R., Weigel, K. M., Harrison, P. L., Lee, K., Cangelosi, G. A., & Yilmaz, Ö. (2016). In situ anabolic activity of periodontal pathogens *porphyromonas gingivalis* and *filifactor alocis* in chronic periodontitis. *Scientific reports*, 6(1), 1–8.
- Sørensen, T. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species and its application to analyses of the vegetation on danish commons. *Biol. Skr.*, 5, 1–34.
- Thukral, R., Shrivastav, K., Mathur, V., Barodiya, A., & Shrivastav, S. (2017). Actinomyces: a deceptive infection of oral cavity. *Journal of the Korean Association of Oral and Maxillofacial Surgeons*, 43(4), 282–285.
- Tucker, C. M., Cadotte, M. W., Carvalho, S. B., Davies, T. J., Ferrier, S., Fritz, S. A., ... others (2017). A guide to phylogenetic metrics for conservation, community ecology and macroecology. *Biological Reviews*, 92(2), 698–715.

## References VIII

- Turnbaugh, P. J., Ley, R. E., Hamady, M., Fraser-Liggett, C. M., Knight, R., & Gordon, J. I. (2007). The human microbiome project. *Nature*, 449(7164), 804–810.
- Van Dyke, T. E., & Dave, S. (2005). Risk factors for periodontitis. *Journal of the International Academy of Periodontology*, 7(1), 3.
- Vielkind, P., Jentsch, H., Eschrich, K., Rodloff, A. C., & Stingu, C.-S. (2015). Prevalence of actinomyces spp. in patients with chronic periodontitis. *International Journal of Medical Microbiology*, 305(7), 682–688.
- Waskom, M., & the seaborn development team. (2020, September). *mwaskom/seaborn*. Zenodo. Retrieved from <https://doi.org/10.5281/zenodo.592845> doi: 10.5281/zenodo.592845

## References IX

Weiss, S., Xu, Z. Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., ... others (2017). Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome*, 5(1), 27.