

Lung Precancer Study

Jaewoong Lee Y. Choi I. Yun S. Park Semin Lee

Department of Biomedical Engineering
Ulsan National Institute of Science and Technology

jwlee230@unist.ac.kr

2021-06-07

Overview

1 Introduction

2 Materials

3 Methods

4 Results

5 Discussion

Introduction

Introduction

Lung Cancer

Lung Cancer?

The most common cancer

The most common form of cancer (12.3 % of all cancers) (Minna, Roth, & Gazdar, 2002)

The most important factor

Tobacco


Type of Lung Cancer

Types of lung cancer:


- ① Adenocarcinoma (ADC) (40 %) ★
- ② Squamous cell carcinoma (SQC) (25 %) ★
- ③ Small cell carcinoma (20 %)
- ④ Large cell carcinoma (10 %)
- ⑤ Adenosquamous carcinoma (< 5 %)
- ⑥ Carcinoid (< 5 %)
- ⑦ Bronchioalveolar (Bronchial gland carcinoma)

(Vincent et al., 1977; Collins, Haines, Perkel, & Enck, 2007)

ADC vs. SQC I




(a) ADC




(b) SQC

Figure: ADC and SQC histology in Lung cancer (Travis, 2002)

ADC vs. SQC II



(a) All patients



(b) By cancer stages

Figure: Kaplan-Meier survival curves for ADC & SQC (Wang et al., 2020)

Findings

SQC is more dangerous than ADC. $\therefore p < 0.001$

Introduction

Study Objectives

Study Objectives

- Find different mutations
 - between WES
 - between WTS
 - from cancer
 - from precancer
- Pathway examine from the mutations
 - of WES
 - of RNA-seq
- Ultra-deep sequencing to find an *infinitesimal* quantity of Non-Circulating Tumor DNA
 - from blood
 - from urine
 - from bronchus
- Diagnostic performance

Materials

Lung Cancer Data

- WES (n=289) + Transcriptome (n=166)
- Normal + {Primary, CIS + AIS, AAH, Dysplasia, MIA}
 - Carcinoma in situ
 - Adenocarcinoma in situ
 - Atypical adenomatous hyperplasia
 - Dysplasia
 - Minimally invasive adenocarcinoma
- Squamous cell carcinoma (SQC) & Adenocarcinoma (ADC)
 - ① Normal → Dysplasia → CIS → SQC (n=80)
 - ② Normal → AAH → AIS → MIA → ADC (n=28)

Methods

Methods

Workflows

Data pre-processing for variant discovery




Figure: Data pre-processing for variant discovery (Van der Auwera et al., 2013; DePristo et al., 2011)

Somatic short variant discovery




Figure: Somatic short variant (SNVs + Indels) discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

Germline short variant discovery




Figure: Germline short variant (SNVs + Indels) discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

RNA-seq short variant discovery




Figure: RNA-seq short variant (SNVs + Indels) discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

Methods

Miscellaneous

Used Bioinformatics Tools

- FastQC (Andrews et al., 2012)
- BWA (H. Li & Durbin, 2009; H. Li, 2013)
- STAR (Dobin et al., 2013)
- Bowtie2 (Langmead & Salzberg, 2012)
- Samtools (H. Li et al., 2009)
- GATK (Van der Auwera et al., 2013; DePristo et al., 2011)
- Picard (*Picard toolkit*, 2019)
- VCF2MAF (Kandoth et al., 2018)
- BCFtools (Danecek et al., 2021)
- VEP (McLaren et al., 2016)
- RSEM (B. Li & Dewey, 2011)
- CIBERSORTx (Steen, Liu, Alizadeh, & Newman, 2020)

R Packages

- Sequenza (Favero et al., 2015)
- Copynumber (Nilsen, Liestol, & Lingjaerde, 2013; Nilsen et al., 2012)
- DESeq2 (Love, Huber, & Anders, 2014)

Python Packages

- Pandas (pandas development team, 2020; Wes McKinney, 2010)
- Sequenza-utils (Favero et al., 2015)
- Matplotlib (Hunter, 2007)
- Seaborn (Waskom & the seaborn development team, 2020)
- CoMut (Crowdis, He, Reardon, & Van Allen, 2020)
- PyClone (Roth et al., 2014)
- Statannot

Results

Results

Quality Checks with FastQC

FastQC?




Figure: Example of FastQC Result (Andrews et al., 2012)

- A quality check tool for sequence data
- Give an overview that which test may be problems

FastQC on WES





Figure: FastQC with WES data


Failure on 33P1 sample

33P1 is excluded at further analysis.

Failure on 33P1 I



(a) 33N



(b) 33P1

Figure: Per Base Sequence Quality Results

Failure on 33P1 II




Figure: Coverage Depth Plot

FastQC on WTS




Figure: FastQC with WTS data

All sample are good to analysis

∴ No sample has more than 5 failures.

Results

Copy Number Variations (CNVs) with Sequenza

Sequenza?






Figure: Representative Output of the Sequenza (Favero et al., 2015)

Cellularity & Ploidy on WES



(a) SQC Samples



(b) ADC Samples

Figure: Cellularity and Ploidy from Sequenza

Genome View on Patient #57




Figure: Dysplasia-CIS-Primary Tumor on Patient #57

CNVs of SQC




Figure: CNV Plot with SQC Patients

CNVs of ADC




Figure: CNV Plot with ADC Patients

Results

Find SNVs with Mutect2

Mutect2?




Figure: Somatic short variant discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

Witer?




Figure: Witer diagram for detecting cancer-drive genes (Jiang et al., 2019)

Somatic Variant with BWA in SQC




Figure: CoMut Plot with SQC Patients

Somatic Variant with BWA in ADC




Figure: CoMut Plot with ADC Patients

Results

Gene Expression Levels from RSEM

RSEM?




Figure: The RSEM workflow (B. Li & Dewey, 2011)

DEG Selection Strategy

Fold Change

$\log_2(\text{Fold Change}) > 1 \vee \log_2(\text{Fold Change}) < -1$

P-value


$P\text{-value} < 0.05$

Adjusted P-value


$\text{Padj} < 0.05$

DEG Volcano Plots with Bowtie2 in SQC


Normal → Dysplasia → CIS → Primary (SQC)




(a) Normal-Dysplasia




(b) Normal-CIS




(c) Normal-Primary



(d) Dysplasia-CIS



(e) Dysplasia-Primary




(f) CIS-Primary

Figure: DEG Volcano Plots with Bowtie2 in SQC

DEG Pseudo-Venn Diagram with Bowtie2 in SQC

Normal → Dysplasia → CIS → Primary (SQC)




(a) Up-regulated

(b) Down-regulated

Figure: DEG Pseudo-Venn Diagram with Bowtie2 in SQC

DEG Venn Diagram with Bowtie2 in SQC

Normal → Dysplasia → CIS → Primary (SQC)




(a) Up-regulated

(b) Down-regulated


Figure: DEG Venn Diagram with Bowtie2 in SQC

DEG Volcano Plots with Bowtie2 in ADC


Normal → AAH → AIS → MIA → Primary (ADC)




(a) Normal-AAH




(b) Normal-AIS




(c) Normal-Primary



(d) AAH-AIS



(e) AAH-Primary




(f) AIS-Primary


Figure: DEG Volcano Plots with Bowtie2 in ADC

DEG Pseudo-Venn Diagram with Bowtie2 in ADC

Normal → AAH → AIS → MIA → Primary (ADC)



(a) Up-regulated




(b) Down-regulated

Figure: DEG Pseudo-Venn Diagram with Bowtie2 in ADC

DEG Venn Diagram with Bowtie2 in ADC

Normal → AAH → AIS → MIA → Primary (ADC)



(a) Up-regulated

(b) Down-regulated

Figure: DEG Venn Diagram with Bowtie2 in ADC

Results

Tumor Evolution with MesKit

MesKit?

Results

Estimated Cell Types with CIBERSORTx

CIBERSORTx?




Figure: CIBERSORTx workflow (Steen et al., 2020)

CIBERSORTx with SQC




Figure: Estimated Cell Types with SQC Samples

CIBERSORTx with ADC




Figure: Estimated Cell Types with ADC Samples

Discussion

References I

- Andrews, S., Krueger, F., Segonds-Pichon, A., Biggins, L., Krueger, C., & Wingett, S. (2012, January). *FastQC*. Babraham Institute. Babraham, UK.
- Collins, L. G., Haines, C., Perkel, R., & Enck, R. E. (2007). Lung cancer: diagnosis and management. *American family physician*, 75(1), 56–63.
- Crowdis, J., He, M. X., Reardon, B., & Van Allen, E. M. (2020). Comut: visualizing integrated molecular information with comutation plots. *Bioinformatics*, 36(15), 4348–4349.
- Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., ... Li, H. (2021, 02). Twelve years of SAMtools and BCFtools. *GigaScience*, 10(2). Retrieved from <https://doi.org/10.1093/gigascience/giab008> (giab008) doi: 10.1093/gigascience/giab008

References II

- DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., ... others (2011). A framework for variation discovery and genotyping using next-generation dna sequencing data. *Nature genetics*, 43(5), 491.
- Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., ... Gingeras, T. R. (2013). Star: ultrafast universal rna-seq aligner. *Bioinformatics*, 29(1), 15–21.
- Favero, F., Joshi, T., Marquard, A. M., Birkbak, N. J., Krzystanek, M., Li, Q., ... Eklund, A. C. (2015). Sequenza: allele-specific copy number and mutation profiles from tumor sequencing data. *Annals of Oncology*, 26(1), 64–70.
- Hunter, J. D. (2007). Matplotlib: A 2d graphics environment. *IEEE Annals of the History of Computing*, 9(03), 90–95.

References III

- Jiang, L., Zheng, J., Kwan, J. S., Dai, S., Li, C., Li, M. J., ... others (2019). Witer: a powerful method for estimation of cancer-driver genes using a weighted iterative regression modelling background mutation counts. *Nucleic acids research*, 47(16), e96–e96.
- Kandoth, C., Gao, J., qwangmsk, Mattioni, M., Struck, A., Boursin, Y., ... Chavan, S. (2018, February). *mskcc/vcf2maf: vcf2maf v1.6.16*. Zenodo. Retrieved from <https://doi.org/10.5281/zenodo.1185418> doi: 10.5281/zenodo.1185418
- Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with bowtie 2. *Nature methods*, 9(4), 357.
- Li, B., & Dewey, C. N. (2011). Rsem: accurate transcript quantification from rna-seq data with or without a reference genome. *BMC bioinformatics*, 12(1), 1–16.

References IV

- Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with bwa-mem. *arXiv preprint arXiv:1303.3997*.
- Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with burrows-wheeler transform. *bioinformatics*, 25(14), 1754–1760.
- Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., ... Durbin, R. (2009). The sequence alignment/map format and samtools. *Bioinformatics*, 25(16), 2078–2079.
- Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for rna-seq data with deseq2. *Genome biology*, 15(12), 1–21.
- McLaren, W., Gil, L., Hunt, S. E., Riat, H. S., Ritchie, G. R., Thormann, A., ... Cunningham, F. (2016). The ensembl variant effect predictor. *Genome biology*, 17(1), 1–14.
- Minna, J. D., Roth, J. A., & Gazdar, A. F. (2002). Focus on lung cancer. *Cancer cell*, 1(1), 49–52.

References V

- Nilsen, G., Liestol, K., & Lingjaerde, O. (2013). *Copynumber: Segmentation of single-and multi-track copy number data by penalized least squares regression. R package version, 1(0)*.
- Nilsen, G., Liestøl, K., Van Loo, P., Vollan, H. K. M., Eide, M. B., Rueda, O. M., ... others (2012). Copynumber: efficient algorithms for single-and multi-track copy number segmentation. *BMC genomics*, 13(1), 1–16.
- pandas development team, T. (2020, February). *pandas-dev/pandas: Pandas*. Zenodo. Retrieved from <https://doi.org/10.5281/zenodo.3509134> doi: 10.5281/zenodo.3509134
- Picard toolkit. (2019). <http://broadinstitute.github.io/picard/>. Broad Institute.

References VI

- Roth, A., Khattra, J., Yap, D., Wan, A., Laks, E., Biele, J., ... Shah, S. P. (2014). Pyclone: statistical inference of clonal population structure in cancer. *Nature methods*, 11(4), 396–398.
- Steen, C. B., Liu, C. L., Alizadeh, A. A., & Newman, A. M. (2020). Profiling cell type abundance and expression in bulk tissues with cibersortx. In *Stem cell transcriptional networks* (pp. 135–157). Springer.
- Travis, W. D. (2002). Pathology of lung cancer. *Clinics in chest medicine*, 23(1), 65–81.
- Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine, A., ... others (2013). From fastq data to high-confidence variant calls: the genome analysis toolkit best practices pipeline. *Current protocols in bioinformatics*, 43(1), 11–10.

References VII

- Vincent, R. G., Pickren, J. W., Lane, W. W., Bross, I., Takita, H., Houten, L., ... Rzepka, T. (1977). The changing histopathology of lung cancer. a review of 1682 cases. *Cancer*, 39(4), 1647–1655.
- Wang, B.-Y., Huang, J.-Y., Chen, H.-C., Lin, C.-H., Lin, S.-H., Hung, W.-H., & Cheng, Y.-F. (2020). The comparison between adenocarcinoma and squamous cell carcinoma in lung cancer patients. *Journal of cancer research and clinical oncology*, 146(1), 43–52.
- Waskom, M., & the seaborn development team. (2020, September). *mwaskom/seaborn*. Zenodo. Retrieved from <https://doi.org/10.5281/zenodo.592845> doi: 10.5281/zenodo.592845

References VIII

Wes McKinney. (2010). Data Structures for Statistical Computing in Python. In Stéfan van der Walt & Jarrod Millman (Eds.), *Proceedings of the 9th Python in Science Conference* (p. 56 - 61). doi: 10.25080/Majora-92bf1922-00a