

Lung Precancer Study

Jaewoong Lee Y. Choi I. Yun S. Park Semin Lee

Department of Biomedical Engineering
Ulsan National Institute of Science and Technology

jwlee230@unist.ac.kr

2021-08-19

Overview

1 Introduction

2 Materials

3 Methods

4 Results

5 Discussion

6 References

Introduction

Introduction

Lung Cancer

Lung Cancer? I

The most common cancer

The most common form of cancer:

12.3 % of all cancers (Minna, Roth, & Gazdar, 2002)

The most important factor

Tobacco

Cancer Survival Rate in Korea



Figure: Common cancer survival rates (Hong et al., 2021)

Survival rate (More than 5 yr)

- Thyroid: 68.4 %
- Lung: 35.4 %

Type of Lung Cancer

Types of lung cancer:

- ① Adenocarcinoma (ADC) (40 %) ★
- ② Squamous cell carcinoma (SQC) (25 %) ★
- ③ Small cell carcinoma (20 %)
- ④ Large cell carcinoma (10 %)
- ⑤ Adenosquamous carcinoma (< 5 %)
- ⑥ Carcinoid (< 5 %)
- ⑦ Bronchioalveolar (Bronchial gland carcinoma)


(Vincent et al., 1977; Collins, Haines, Perkel, & Enck, 2007)

ADC vs. SQC I




Figure: ADC and SQC histology in Lung cancer (Travis, 2002)

ADC vs. SQC II



(a) All patients



(b) By cancer stages

Figure: Kaplan-Meier survival curves for ADC & SQC (B.-Y. Wang et al., 2020)

Findings

SQC is more dangerous than ADC. $\therefore p < 0.001$

Introduction

Study Objectives

Study Objectives

Find different mutations

- between WES vs. WTS
- from cancer vs. precancer

Pathway examine

- with the mutation of WES & RNA-seq
- with immune-depleted animal models

Ultra-deep sequencing

to find an *infinitesimal* quantity of Non-Circulating Tumor DNA

- from blood
- from urine
- from bronchus

Materials

Lung Cancer Data

- WES (n=289) + Transcriptome (n=166)
- Normal + {Primary, CIS + AIS, AAH, Dysplasia, MIA}
 - Carcinoma in situ
 - Adenocarcinoma in situ
 - Atypical adenomatous hyperplasia
 - Dysplasia
 - Minimally invasive adenocarcinoma
- Squamous cell carcinoma (SQC) & Adenocarcinoma (ADC)
 - ① Normal → Dysplasia → CIS → SQC (n=80)
 - ② Normal → AAH → AIS → MIA → ADC (n=28)

Methods

Methods

Workflows

Data pre-processing for variant discovery




Figure: Data pre-processing for variant discovery (Van der Auwera et al., 2013; DePristo et al., 2011)

Somatic short variant discovery



Figure: Somatic short variant (SNVs + Indels) discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

Germline short variant discovery



Figure: Germline short variant (SNVs + Indels) discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

RNA-seq short variant discovery



Figure: RNA-seq short variant (SNVs + Indels) discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

Results

Results

Quality Checks

FastQC?



Figure: Example of FastQC Result (Andrews et al., 2012)

- A quality check tool for sequence data
- Give an overview that which test may be problems

FastQC on WES





Figure: FastQC with WES data


Failure on 33P1 sample

33P1 is excluded at further analysis.

Failure on 33P1 I



(a) 33N



(b) 33P1

Figure: Per Base Sequence Quality Results

Failure on 33P1 II



Figure: Coverage Depth Plot

FastQC on WTS




Figure: FastQC with WTS data

All sample are good to analysis

∴ No sample has more than 5 failures.

Results

Copy Number Variations

Sequenza?






Figure: Representative Output of the Sequenza (Favero et al., 2015)

Cellularity & Ploidy on WES



(a) SQC Samples



(b) ADC Samples

Figure: Cellularity and Ploidy from Sequenza

Genome View on Patient #57




Figure: Dysplasia-CIS-Primary Tumor on Patient #57

CNVs of SQC




Figure: CNV Plot with SQC Patients

CNVs of ADC




Figure: CNV Plot with ADC Patients

SQC vs. ADC




Figure: Simple CNV Plot with SQC Patients




Figure: Simple CNV Plot with ADC Patients

Findings in Sequenza

Sequenza Findings

- SQC have more CNVs than ADC.
- SQC have aggressive CNVs on chromosome #3 and #5.
- ADC have aggressive CNVs on chromosome #5.

Results

SNVs Analysis

Mutect2?





Figure: Somatic short variant discovery workflow (Van der Auwera et al., 2013; DePristo et al., 2011)

MutEnricher?



Analysis summary:

Inputs:

- Somatic mutations
- Features of interest:
 - Coding genes
 - Non-coding regions
- Genomic covariates (optional)

Analyses:

- Background calculations:
 - global, local, or covariate clustered
- Mutation enrichments:
 - coding/non-coding modules

Outputs:

- Gene or non-coding region enrichments:
 - Overall genes/regions
 - Hotspots
 - Combined

Figure: Schematic representation of MunEnricher's analysis procedures (Soltis et al., 2020)

Somatic Variant in SQC




Figure: CoMut Plot with SQC Patients

Somatic Variant in ADC




Figure: CoMut Plot with ADC Patients

Findings in SNVs Analysis

TTN

TTN is the most mutated gene both in SQC and ADC.

Results

VAF Analysis

VAF?

- Variant allele frequency
- VAF = Alternative allele read count/Total read count
- To find tumor evolution




Figure: VAF distribution of validated mutations (Ding et al., 2012)

VAF Plots I

PyClone?




Figure: Analysis of multiple samples by PyClone (Roth et al., 2014)

PyClone Plots I

Findings in VAF Analysis

Results

Differences in Gene Expression Levels

RSEM?




Figure: RSEM workflow (Li & Dewey, 2011)

DESeq2




Figure: DESeq2 workflow (Love, Huber, & Anders, 2014)

DEG Selection Strategy

DEG: differentially expressed genes

Fold Change

$$\log_2(\text{Fold Change}) > 1 \vee \log_2(\text{Fold Change}) < -1$$

P-value


$$P\text{-value} < 0.05$$

Adjusted P-value


$$P_{adj} < 0.05$$

DEG Volcano Plots in SQC


Normal → Dysplasia → CIS → Primary (SQC)




(a) Normal-Dysplasia




(b) Normal-CIS




(c) Normal-Primary



(d) Dysplasia-CIS



(e) Dysplasia-Primary




(f) CIS-Primary

Figure: DEG Volcano Plots in SQC

DEG Venn Diagram with Bowtie2 in SQC

Normal → Dysplasia → CIS → Primary (SQC)




(a) Up-regulated

(b) Down-regulated


Figure: DEG Venn Diagram in SQC

DEG Volcano Plots with Bowtie2 in ADC


Normal → AAH → AIS → MIA → Primary (ADC)




(a) Normal-AAH




(b) Normal-AIS




(c) Normal-Primary



(d) AAH-AIS



(e) AAH-Primary



(f) AIS-Primary

Figure: DEG Volcano Plots in ADC

DEG Venn Diagram with Bowtie2 in ADC

Normal → AAH → AIS → MIA → Primary (ADC)



(a) Up-regulated

(b) Down-regulated

Figure: DEG Venn Diagram in ADC

Findings in DEG Analysis

Results

Bulk Cell Deconvolution

Single-cell data as Reference




Figure: Comprehensive dissection and clustering of 208,506 single cells from LUAD patients (Kim et al., 2020)

CIBERSORTx




Figure: Workflow for CIBERSORTx (Steen et al., 2020; Newman et al., 2019)

Cluster Plot with Bowtie2 in ADC




Figure: Cluster Plot in ADC

Cluster Plot with Bowtie2 in SQC




Figure: Cluster Plot in SQC

Benchmarking of Cell Deconvolution Tools




Figure: Memory and time requirements for the cell deconvolution methods (Cobos et al., 2020)

Top 3 Methods

- ① BisqueRNA (Jew et al., 2020)
- ② MuSiC (X. Wang, Park, Susztak, Zhang, & Li, 2019)
- ③ SCDC (Dong et al., 2021)

BisqueRNA?




Figure: Graphical overview of the Bisque decomposition methods (Jew et al., 2020)

MuSiC?





Figure: Overview of MuSiC framework (X. Wang et al., 2019)

SCDC?

Bulk RNA-seq



Bulk Sample 2




Bulk Sample 3




scRNA-seq



Single-Cell Reference 1



Single-Cell Reference 2



Deconvolution



ENSEMBLE




Figure: Overview of deconvolution by SCDC (Dong et al., 2021)

Findings in Bulk Cell Deconvolution

Results

Tumor Evolution Trajectories Analysis

Revolver?




Figure: Repeated Evolutionary Trajectories (Caravagna et al., 2018)

Findings in Tumor Evolution Trajectories Analysis

Results

Gene Fusion

title

Discussion

References

References I

- Andrews, S., Krueger, F., Segonds-Pichon, A., Biggins, L., Krueger, C., & Wingett, S. (2012, January). *FastQC*. Babraham Institute. Babraham, UK.
- Caravagna, G., Giarratano, Y., Ramazzotti, D., Tomlinson, I., Graham, T. A., Sanguinetti, G., & Sottoriva, A. (2018). Detecting repeated cancer evolution from multi-region tumor sequencing data. *Nature methods*, 15(9), 707–714.
- Cobos, F. A., Alquicira-Hernandez, J., Powell, J. E., Mestdagh, P., & De Preter, K. (2020). Benchmarking of cell type deconvolution pipelines for transcriptomics data. *Nature communications*, 11(1), 1–14.
- Collins, L. G., Haines, C., Perkel, R., & Enck, R. E. (2007). Lung cancer: diagnosis and management. *American family physician*, 75(1), 56–63.

References II

- DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., ... others (2011). A framework for variation discovery and genotyping using next-generation dna sequencing data. *Nature genetics*, 43(5), 491.
- Ding, L., Ley, T. J., Larson, D. E., Miller, C. A., Koboldt, D. C., Welch, J. S., ... others (2012). Clonal evolution in relapsed acute myeloid leukaemia revealed by whole-genome sequencing. *Nature*, 481(7382), 506–510.
- Dong, M., Thennavan, A., Urrutia, E., Li, Y., Perou, C. M., Zou, F., & Jiang, Y. (2021). Scdc: bulk gene expression deconvolution by multiple single-cell rna sequencing references. *Briefings in bioinformatics*, 22(1), 416–427.

References III

- Favero, F., Joshi, T., Marquard, A. M., Birkbak, N. J., Krzystanek, M., Li, Q., ... Eklund, A. C. (2015). Sequenza: allele-specific copy number and mutation profiles from tumor sequencing data. *Annals of Oncology*, 26(1), 64–70.
- Hong, S., Won, Y.-J., Lee, J. J., Jung, K.-W., Kong, H.-J., Im, J.-S., ... others (2021). Cancer statistics in korea: Incidence, mortality, survival, and prevalence in 2018. *Cancer Research and Treatment: Official Journal of Korean Cancer Association*, 53(2), 301.
- Jew, B., Alvarez, M., Rahmani, E., Miao, Z., Ko, A., Garske, K. M., ... Halperin, E. (2020). Accurate estimation of cell composition in bulk expression through robust integration of single-cell information. *Nature communications*, 11(1), 1–11.

References IV

- Kim, N., Kim, H. K., Lee, K., Hong, Y., Cho, J. H., Choi, J. W., ... others (2020). Single-cell rna sequencing demonstrates the molecular and cellular reprogramming of metastatic lung adenocarcinoma. *Nature communications*, 11(1), 1–15.
- Li, B., & Dewey, C. N. (2011). Rsem: accurate transcript quantification from rna-seq data with or without a reference genome. *BMC bioinformatics*, 12(1), 1–16.
- Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for rna-seq data with deseq2. *Genome biology*, 15(12), 1–21.
- Minna, J. D., Roth, J. A., & Gazdar, A. F. (2002). Focus on lung cancer. *Cancer cell*, 1(1), 49–52.

References V

- Newman, A. M., Steen, C. B., Liu, C. L., Gentles, A. J., Chaudhuri, A. A., Scherer, F., . . . others (2019). Determining cell type abundance and expression from bulk tissues with digital cytometry. *Nature biotechnology*, 37(7), 773–782.
- Roth, A., Khattra, J., Yap, D., Wan, A., Laks, E., Biele, J., . . . Shah, S. P. (2014). Pyclone: statistical inference of clonal population structure in cancer. *Nature methods*, 11(4), 396–398.
- Soltis, A. R., Dalgard, C. L., Pollard, H. B., & Wilkerson, M. D. (2020). Mutenricher: a flexible toolset for somatic mutation enrichment analysis of tumor whole genomes. *BMC bioinformatics*, 21(1), 1–8.
- Steen, C. B., Liu, C. L., Alizadeh, A. A., & Newman, A. M. (2020). Profiling cell type abundance and expression in bulk tissues with cibersortx. In *Stem cell transcriptional networks* (pp. 135–157). Springer.

References VI

- Travis, W. D. (2002). Pathology of lung cancer. *Clinics in chest medicine*, 23(1), 65–81.
- Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., Del Angel, G., Levy-Moonshine, A., ... others (2013). From fastq data to high-confidence variant calls: the genome analysis toolkit best practices pipeline. *Current protocols in bioinformatics*, 43(1), 11–10.
- Vincent, R. G., Pickren, J. W., Lane, W. W., Bross, I., Takita, H., Houten, L., ... Rzepka, T. (1977). The changing histopathology of lung cancer. a review of 1682 cases. *Cancer*, 39(4), 1647–1655.
- Wang, B.-Y., Huang, J.-Y., Chen, H.-C., Lin, C.-H., Lin, S.-H., Hung, W.-H., & Cheng, Y.-F. (2020). The comparison between adenocarcinoma and squamous cell carcinoma in lung cancer patients. *Journal of cancer research and clinical oncology*, 146(1), 43–52.

References VII

Wang, X., Park, J., Susztak, K., Zhang, N. R., & Li, M. (2019). Bulk tissue cell type deconvolution with multi-subject single-cell expression reference. *Nature communications*, 10(1), 1–9.