Bio 338 Modeling Biological Dynamics Notes

Kyle Wang

September 2026

Contents

1 Basic Modeling with ODEs **1**
1.1 Introduction . . . 1
1.2 Predation . . . 3
1.3 HIV . . . 3
1.4 Tangent Spaces . . . 5
1.5 Fixed Points and Stability . . . 7
1.6 Bifurcations . . . 8

1 Basic Modeling with ODEs

1.1 Introduction

State: numbers that describe variables (state variables) of interest
State variables: Things we care about and measure
State space: space of allowable values of a state variable
One thing to note is that units can affect the state space. For example, the state space for allowable temperatures in Celsius is $C \in [-273.15, +\infty)$ whereas the allowable temperatures in Kelvin is $K \in [0, +\infty)$.
We can express differential equations as compartment diagrams. This diagram represents linear growth and is




Figure 1: Compartment diagram of linear growth

isomorphic to the equation

$$\dot{x} = b$$

For these examples, let us use x to describe the rabbit population. In this case, the population is increasing by b rabbits per time unit (we shall say month for the example). Figure 2 is the compartment diagram for the equation

$$\dot{x} = bx$$

This effectively represents that more rabbits means more offspring which physically makes sense. We can adjust this model further by adding a death rate d. We can then draw the compartment diagram as seen in Figure 3. The isomorphic differential equation for this is

$$\dot{x} = bx - dx = (b - d)x$$




Figure 2: Compartment diagram of per capita growth

which has the solution

$$x(t) = x_0 e^{rt} \qquad r = b - d$$

If $r > 0$, then the birth rate "out competes" the death rate which leads to exponential growth. On the other hand, if $r < 0$, the death rate overrides the birth and hence the rabbit population disappears. We can make it more realistic by adding carrying capacity that only allows k rabbits in our system. This system is described by the *Euler-Lotka* Model

$$\dot{x} = r\left(1 - \frac{x}{k}\right)x$$

where r is once again the sum of the birth and death rates. The second term effectively scales the birth and death based off of how close the population is to the carrying capacity. We must have that $r > 0$ or else we see that the amount of deaths slows down as we approach the carrying capacity which is unphysical. If we think about when




Figure 3: Compartment diagram of per capita growth and death

$r < 0$, we also see that if $x > k$, we have unbounded growth since r and $\left(1 - \frac{x}{k}\right)$ is also negative leading to $\dot{x} > 0$

and hence unbounded growth. Even if it hypothetically did work, we wouldn't really care since $r < 0$ should force everything to zero and so k doesn't have any impact on our model. Checking the behavior when $r > 0$, we find that if $0 < x < k$, we should expect growth and indeed $\dot{x} > 0$. When the number of rabbits is above the carrying capacity, we should expect the population to decline and indeed that occurs as $\dot{x} < 0$.

It is also worth noting that this model assumes the number of rabbits is a continuous variable which implies fractional (or irrational) number of rabbits which is clearly not reality. We disregard this concern for large populations because the difference between 1000.5 and 1001 rabbits is trivial; however, for small populations, we would need to enforce a discrete model since the difference of one rabbit is more significant on that scale.

1.2 Predation

Suppose that foxes (y) eat rabbits (x). Also suppose that the rate of foxes is dependent on the food supply. We can then write the differential equation describing the number of foxes as

$$\dot{y} = b[\text{food}]y - Dy$$

We can define the food supply as the probability of a single predation event as p multiplied by the number of rabbits there are x. We can then multiply the food supply available by the number of wolves to feed this which hence leads to

$$\dot{y} = bpxy - Dy$$

where D is the death rate per capita for wolves. Before, we described the rabbits as just $\dot{x} = rx$. Now we have to factor in the deaths due to foxes which is once again the food supply multiplied by the number of foxes. We now have the set of differential equations

$$\dot{x} = rx - pxy$$
$$\dot{y} = bpxy - Dy$$

And if we let all the coefficients equal 1, we get the *Lotka-Volterra Model*.

$$\dot{x} = x - xy$$
$$\dot{y} = xy - y$$

1.3 HIV

What perplexed people about HIV was the timeline of the disease. The viral load would initially spike and decline within the first 8 weeks of infection. It would then stay at a low viral load for years (in what is deemed its latent phase) before slowly increasing and leading to AIDS. There were two main questions:

1. What causes the initial spike and decline?
2. What causes the sudden climb in viral load leading to AIDS?

The initial hypothesis was that there wold be rapid viral growth and then the immune system would respond (leading to increase and decline in viral load) before replicating slowly in the background. In 1994, the first antiretroviral drugs were in clinical trials. These vaccines worked by stopping viral replication rather than directly killing existing viral particles. According to the previous hypothesis, these drugs shouldn't make a significant impact since the viruses are only replicating slowly. What they noticed was an exponential decay in viral load as time progress after the treatment.

In 1996, Alan Perelson publishes *HIV-1 Dynamics in Vivo: Virion Clearance Rate, Infected Cell Life-Span, and Viral Generation Time* to explain what was observed. Let us consider a basic bathtub with a faucet and a drain. The inflow is described as Q_i and the outflow is described as Q_o and is dependent on the volume in the bathtub. We can therefore describe the amount of volume in the bathtub, V as the following differential equation

$$\dot{V} = Q_i - Q_o V$$

We especially care about the steady state condition which is when $\dot{V} = 0$. This leads to $V = \frac{Q_i}{Q_o}$ which effectively showed that the volume at steady state is dependent on the *ratio* and not the magnitudes of Q_i and Q_o. We could therefore have a faucet with high mass flow rate but also a really efficient drain or a dripping faucet and inefficient

drain. We can't tell by solely looking at the amount of water in the drain. This extremely basic model already gives the insight that the virus is replicating extremely quickly, but the immune system is also clearing them quickly too. If we solve for $V(t)$ generally, we get

$$\begin{aligned} \dot{V} &= Q_i - Q_o V \\ \int \frac{1}{Q_i - Q_o V} dV &= t + c \\ V(t) &= \frac{Q_i}{Q_o} + Ce^{-Q_o t} \end{aligned}$$

If we apply the initial condition that $V(t=0) = V_0$, then we find the constant to be $C = V_0 - \frac{Q_i}{Q_o}$. If Q_i and Q_o are small, we should expect small exponential decay and vice versa. This effectively explains the dramatic difference the antiretrovirals made. Let us now consider a more sophisticated model. One thing to note is that the line from N_v




Figure 4: Compartment diagram of a model similar to Perelson's

implies that the rate of infection is dependent on the number of virus particles. There is also a negligible consumption of N_v in this process. Shedding also depends on N_i but doesn't change N_i. We can then write these as a set of differential equations

$$\begin{aligned} \dot{N}_u &= B_u - D_u N_u - B_i N_v N_u \\ \dot{N}_i &= B_i N_v N_u - D_i N_i \\ \dot{N}_v &= B_v N_i - D_v N_v \end{aligned}$$

These equations are annoying to solve analytically but we can solve it numerically. We can then graph as seen in Figure 5. We can further adjust the model to include latent infected cells: that is, cells that are infected but do not




Figure 5: Plot of the amount of HIV in the blood over time with the parameters using McClain's model: $B_u = 0.272, B_i = 0.00027, B_v = 100, D_u = 0.00136, D_i = 0.33, D_v = 2$

shed viruses. Let us say that 10% of cells become latently infected. Let us assume the death rate of latent cells is

the same as uninfected ($D_l = D_u$). Let us also say that the latent cells become active (and begin shedding) at a rate B_a. Our set of differential equations is now

$$\begin{aligned}
\dot{N}_u &= B_u - D_u N_u - B_i N_v N_u \\
\dot{N}_i &= .9 B_i N_v N_u + B_a N_l - D_i N_i \\
\dot{N}_l &= .1 B_i N_v N_u - B_a N_l - D_u N_l \\
\dot{N}_v &= B_v N_i - N_v D_v
\end{aligned}$$




Figure 6: McClain's model including latently infected cells

1.4 Tangent Spaces

Definition 1.1 (Tangent Space)**.** *The tangent space of a* manifold *is a vector space that contains all possible directions tangent to the manifold at that point.*

Definition 1.2 (Manifold)**.** *An n-dimensional manifold is a topological space with the property that each point has a neighborhood that is homeomorphic to an open subset of n-dimensional Euclidean space.*

We can effectively think of the a state space as a manifold and each point on that manifold corresponds to a specific state in that state space. We can then think of the tangent spaces as all possible changes that can be made to a certain state in that state space. I wish I knew more about differential geometry but maybe Gábor Székelyhidi can teach me about it.

Definition 1.3 (Vector Field)**.** *A vector is a function f : points in the state space $(\mathbb{R}^n) \mapsto$ points in the tangent space $(\mathbb{R}^n)$.*

Let us look at the example $x \in [0, \infty)$ to be our state space and $\dot{x} = 2x$ as our manifold. The state space is the positive half line and so vectors in this tangent space can point in either direction. So -5 or 2 may be allowable vectors in this tangent space. However, the *vector field* is strictly enforced by $\dot{x} = 2x$ which is a *subset* of the tangent space. We can also look at the vector field for the logistic model where $x \in [0, \infty)$ and $\dot{x} = rx\left(1 - \frac{x}{k}\right),\ r > 0$. We now note for $x < k$, the vector field points in the $+\hat{x}$ whereas if $x > k$, the vector field points in the $-\hat{x}$ direction. That is because there exists a fixed point at k.

Theorem 1.4 (Banach Fixed Point Theorem)**.** *Let (X, d) be a* complete *metric space with a contraction mapping $T : X \mapsto X$. Then T admits a unique fixed point $x^* \in X$ meaning $T(x^*) = x^*$. Furthermore, X^* can be found as follows: start with an arbitrary element $x_0 \in X$ and define $x_n = T(x_{n-1})$ for $n \in \mathbb{N}$. Then,* $\lim_{x \to \infty} x_n = x^*$

Let us look at some vector fields in $\mathbb{R}^2$. Consider the Romeo and Juliet system of differential equations which models Romeo's love y and Juliet's love x as

$$\begin{aligned}
\dot{y} &= -x \\
\dot{x} &= y
\end{aligned}$$

We can also take a look at the Lotka-Volterra model which describes predator-prey dynamics and is described by the equations

$$\begin{aligned}
\dot{x} &= x - kxy \\
\dot{y} &= kxy - y
\end{aligned}$$




Figure 7: Vector field of the Romeo and Juliet set of differential equations

We can see that in Figure 8, there exists a fixed point at the origin (no predators and no prey so no dynamics) and another one somewhere in the positive plane which suggests the existence of a steady state.




Figure 8: Vector field of the Lotka-Volterra model

Definition 1.5. Trajectories *are solutions to ODEs that exist and are unique (assuming the conditions of the Picard-Lindelöf Theorem are satisfied) and are* curves *in the state space that follow the vector field.*

We can describe a vector field V to be a function $V : \mathbb{R}^n \mapsto T_pM$ where T_pM denotes the tangent space at a point p along the state space manifold M. It is worth noting that two trajectories can *never* cross since that would require the vector field to be multi valued which is not possible over $\mathbb{R}$. One way to approximate solution curves to ODEs is via Euler's method which is as follows

$$x(t + \Delta t) = x(t) + x'(t)\Delta t$$

Other numerical methods improve on Euler's method like Runge-Kutta but Euler's method is the most clear to see. Some numerical integrators can have adaptive time steps that are based on the magnitude of the local vector field. This is because small time steps offer better accuracy at the expense of compute time. Large time steps are faster to calculate but may not converge to the true solution and are prone to deviations.

1.5 Fixed Points and Stability

Definition 1.6. *A* fixed point $\mathbf{x}$ *is a point in the state space such that* $\dot{\mathbf{x}} = 0$.

The vector field at a fixed point is the zero vector. We can classify the stability of a fixed point into a few categories:

(i) A system is *stable* if it returns to its fixed point when given a slight perturbation from it

(ii) A system is *unstable* if it leaves its fixed point when given a slight perturbation

(iii) A system is *quasi-stable* or *semi-stable* if one direction is stable while the direction is unstable (saddle point). We generally consider this to be unstable since there is only an infinitesimally small line of stability and the rest of the space unstable.

It is worth noting that there are different notions of stability. We shall state two.

Definition 1.7. *Consider an autonomous nonlinear dynamical system*

$$\dot{x} = f(x(t)) \qquad x(0) = x_0$$

where $x(t) \in \mathcal{D} \subseteq \mathbb{R}^n$ *denotes the system state vector,* $\mathcal{D}$ *denotes an open set containing the origin, and* $f : \mathcal{D} \mapsto \mathbb{R}^n$ *is a continous vector field on* $\mathcal{D}$*. Suppose* f *has an equilibrium at* x_e *such that* $f(x_e) = 0$*. Then:*

(i) The fixed point is said to be Lyapunov stable *if for every* $\epsilon > 0$ $\exists \delta > 0$ *such that* $||x(0) - x_e|| < \delta$ *then for every* $t \geq 0$ *we have that* $||x(t) - x_e|| \leq \epsilon$*.*

(ii) The fixed point is said to be asymptotically stable *if it is Lyapunov stable and there exists a* $\delta > 0$ *such that if* $||x(0) - x_e|| < \delta$ *then* $\lim_{t\to\infty} ||x(t) - x_e|| = 0$

In essence, a *Lyapunov stable* system just needs to stay within an ϵ distance from the fixed point given it starts δ away. An *asymptotically stable* system implies that close enough starting conditions will lead to convergence to the fixed. This means something that just orbits the fixed point can be consider Lyapunov stable but not asymptotically stable.

Definition 1.8. *A* nullcline *is a curve in the tangent space where the vector field is 0 in a particular direction.*

In a two dimensional state space, the location where two nullclines meet is a fixed point provided the nullcines are of *different* state variables. If we look at the Lotka-Volterra models and add in competition or communicable disease, we have

$$\dot{x} = Ax - Bxy - x^2$$
$$\dot{y} = Cy - Dxy - y^2$$

We can solve for nullclines by setting $\dot{x} = \dot{y} = 0$ and solving a system of equations. If we ignore the trivial nullclines, we get that nontrivial fixed point of this model is

$$x = \frac{A - BC}{1 - BD} \qquad y = C - D\left(\frac{A - BC}{1 - BD}\right)$$

When $\frac{A}{B} > C$, this fixed point is stable but when $\frac{A}{B} < C$, this fixed point is now unstable.

Definition 1.9. *An* attactor *of a dynamical system is a set of states to which the system tends to evolve for a large number of starting states. We define "a large number" to be a set that isn't of measure* 0*.*

An attractor can be a stable fixed point or a limit cycle. It cannot be an unstable fixed point nor a saddle (which is a set of starting states that leads to a fixed point of measure 0). The *Basin of Attraction* is the set of states that will evolve to a particular attractor over time. Let us consider a logistic model with the allee effect which states that a population needs to be above some minimum value to reproduce. It is described by the following model:

$$\dot{x} = rx\left(1 - \frac{x}{k}\right)\left(\frac{x}{a} - 1\right) \qquad 0 < a < k$$

We see that there are fixed points at $x = 0, a, k$. If we take derivatives of the time deriavtives with respect to x we can classify their stabilities

$$\frac{\partial \dot{x}}{\partial x} = r\left(1 - \frac{x}{k}\right)\left(\frac{x}{a} - 1\right) - \frac{r}{k}x\left(\frac{x}{a} - 1\right) + \frac{r}{a}x\left(1 - \frac{x}{k}\right)$$

At $x = a$, we have

$$\left.\frac{\partial \dot{x}}{\partial x}\right|_{x=a} = r\left(1 - \frac{a}{k}\right) > 0$$

which therefore makes this point unstable. At $x = k$, we have

$$\left.\frac{\partial \dot{x}}{\partial x}\right|_{x=k} = r\left(\frac{k}{a} - 1\right) < 0$$

and is hence stable. At $x = 0$, the derivative evaluates to $-r < 0$ and is also stable. We therefore have two basis of attraction: one that goes towards $x = 0$ and one towards $x = k$. The set $[0, a)$ leads to the fixed point $x^* = 0$ and the set $(a, +\infty)$ leads to $x^* = k$. An attractor is an element of its basin. The boundary between basis is known as the *separatrix*. In one dimension, it is an unstable fixed point. In two dimensions, its a curve. For higher dimensions, the separatrix is always a hyperplane.

1.6 Bifurcations

Recall in the competitive Lotka-Volterra model that depending on whether $\frac{A}{B} > C$ determined the stability of the fixed point. This transition due to a change in parameters is known as a *bifurcation*. Let us look at an example in one dimension. E. coli can consume lactose but importing lactose is an energy expensive process. As such, there is a switch that causes the E. coli to import lactose or not. Let us define x to be hte intracellular lactose level. The change in it $\dot{x}$ is equal to the production minus the degradation. The production is modeled by

$$\text{production} = \frac{a + x^n}{1 + x^n}$$

and the degradation is per capita degradation and therefore modeled as rx. As such,

$$\dot{x} = \frac{a + x^n}{1 + x^n} - rx$$

We can plot the production and degradation terms separately to get Figure 9 Regions where the red line (degradation)




Figure 9: Production and degradation terms plotted separately via the over-under method

is greater than the purple sigmoid (production) is net negative change in lactose levels and vice versa.