

Math 330 Abstract Algebra Notes

Kyle Wang

September 2026

Contents

1	Group Theory	1
1.1	Modular Arithmetic	1
1.2	Introduction to Groups	2

1 Group Theory

1.1 Modular Arithmetic

Definition 1.1. Take $n > 0$ and $n \in \mathbb{Z}$. We say that $a, b \in \mathbb{Z}$ are *congruent modulo n* ($a \equiv b \pmod{n}$) if n divides $a - b$ ($n|a - b$) or if the remainder of n/a equals the remainder of n/b .

Here are some properties:

1. if $a \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$, then $a + c \equiv b + d \pmod{n}$.

Proof. $a \equiv b \pmod{n}$ implies that $n|a - b$ which implies that $\exists q \in \mathbb{Z}$ such that $a - b = nq$. The same can be said for $a \equiv b \pmod{n}$ for some $q' \in \mathbb{Z}$ such that $c - d = nq'$. Adding these equations together we have that $a - b + c - d = (a + c) - (b + d) = n(q + q')$. Since the integers are closed under addition, $q + q'$ is an integer and therefore $n|(a + c) - (b + d)$ and as a result, $a + c \equiv b + d \pmod{n}$. $\square$

2. if $a \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$, then $ac \equiv bd \pmod{n}$

Proof. $a \equiv b \pmod{n}$ implies that $n|a - b$ which implies that $\exists q \in \mathbb{Z}$ such that $a - b = nq$. The same can be said for $a \equiv b \pmod{n}$ for some $q' \in \mathbb{Z}$ such that $c - d = nq'$. We can write ac as

$$\begin{aligned} ac &= (b + nq)(d + nq') \\ &= bd + bnq' + dnq + n^2qq' \\ \Rightarrow ac - bd &= n(bq' + dq + nqq') \end{aligned}$$

and since everything in the parentheses are integers and the integers are closed under multiplication and addition, their result is an integer and therefore $n|ac - bd$ and $ac \equiv bd \pmod{n}$. $\square$

3. Congruence mod n is an equivalence relation

Proof. An equivalence relation is reflexive, symmetric, and transitive. To prove reflexivity, we show $a \equiv a \pmod{n} \rightarrow n|a - a \rightarrow n|0$ which is always true. To prove symmetry, we have that $a \equiv b \pmod{n}$ implies $(a - b) = nq$ which we can extract a negative sign such that $(b - a) = n(-q)$ and since $-q$ is also an integer, $n|b - a$ and therefore $b \equiv a \pmod{n}$. To prove transitivity, we say that $a \equiv b \pmod{n}$ implies $a - b = nq$ and $b \equiv c \pmod{n}$ implies $b - c = nr$. We now find that $a - c = b + nq + nr - b = n(q + r)$. Since the integers are closed under addition, $q + r \in \mathbb{Z}$ and therefore $n|a - c$ and $a \equiv c \pmod{n}$. $\square$

1.2 Introduction to Groups

Definition 1.2 (Group). A group is a set of elements G with a binary operation $*$ that assigns to every pair (g, h) of elements of G another element in G denoted $g * h$. Groups follow the group axioms:

1. Associativity

$$(g_1 * g_2) * g_3 = g_1 * (g_2 * g_3)$$

2. There exists an element $e \in G$ that is the identity such that

$$e * g_1 = g_1 * e = g_1$$

3. Every element has an inverse element denoted g^{-1} such that

$$g * g^{-1} = g^{-1} * g = e$$

4. The set of elements is closed under the group operation (although this is implied in the definition).

Example 1.3. Let us look at some examples

1. The set $(\mathbb{Z}, +)$. Addition is associative, the identity is 0, the inverse of each element g is $-g$.
2. The set $(\mathbb{R}_{>0}, \cdot)$ where $\cdot$ denotes regular multiplication. Regular multiplication is associative, the identity is 1, the inverse element is $\frac{1}{g}$.
3. The set $(\{-2, -1, 0, 1, 2\}, +)$. This fails due since it is not closed (take $1 + 2 = 3 \notin$ the set).
4. The set $(\{\frac{1}{2^n}\}, \cdot)$. This fails since there is no inverse.
5. Fix a positive integer n . Take the set $(\{0, 1, 2, \dots, n-1\}, + \bmod n)$. The identity is 0, addition is associative. The inverse of an element $0 \leq a \leq n-1$ is $n-a$ since $(n-a) + a \bmod n = n \bmod n = 0$. This group is denoted by Fraleigh as $\mathbb{Z}$ but more generally denoted as $\mathbb{Z}/n\mathbb{Z}$.
6. Fix an $n \in \mathbb{N}$. We denote the group $\text{GL}_n(\mathbb{R})$ as the set of invertible $n \times n$ matrices with real entries with the operation of matrix multiplication. Matrix multiplication is associative, the identity is the $n \times n$ identity matrix I_n . The inverse of every element A is the inverse matrix of A . This group is also known as the *General Linear Group*.