

Partition universality

P. Allen, D.M.C., J. Böttcher

size-Ramsey number


Ramsey number

Let's check the Ramsey number $r_2(2K_2)$. ($2K_2 = \mathfrak{I} \mathfrak{I}$)

size-Ramsey number

Ramsey number


Let's check the Ramsey number $r_2(2K_2)$. ($2K_2 = \begin{smallmatrix} \circ & \circ \\ | & | \\ \circ & \circ \end{smallmatrix}$)



size-Ramsey number

Ramsey number


Let's check the Ramsey number $r_2(2K_2)$. ($2K_2 = \begin{smallmatrix} \circ & \circ \\ \parallel & \parallel \end{smallmatrix}$)



size-Ramsey number

Ramsey number


Let's check the Ramsey number $r_2(2K_2)$. ($2K_2 = \begin{smallmatrix} \circ & \circ \\ \parallel & \parallel \end{smallmatrix}$)



size-Ramsey number

Ramsey number


Let's check the Ramsey number $r_2(2K_2)$. ($2K_2 = \begin{smallmatrix} \circ & \circ \\ \parallel & \parallel \end{smallmatrix}$)



size-Ramsey number

Ramsey number

Let's check the Ramsey number $r_2(2K_2)$. ($2K_2 = \text{two disjoint edges}$)



size-Ramsey number

Towards size-Ramsey

There are smaller $2K_2$ -Ramsey graphs.

size-Ramsey number


Towards size-Ramsey

There are **smaller** (?) $2K_2$ -Ramsey graphs.

size-Ramsey number

Towards size-Ramsey

There are smaller $2K_2$ -Ramsey graphs.



size-Ramsey number

size-Ramsey

The r -size-Ramsey number of F is the minimum $e(\Gamma)$ such that $\Gamma \rightarrow_r F$.

size-Ramsey number

size-Ramsey

The r -size-Ramsey number of F is the minimum $e(\Gamma)$ such that $\Gamma \rightarrow_r F$.


$$\hat{r}_3(2K_2)$$

size-Ramsey number

size-Ramsey

The r -size-Ramsey number of F is the minimum $e(\Gamma)$ such that $\Gamma \rightarrow_r F$.

$$\hat{r}_3(2K_2) = 4$$




size-Ramsey number

size-Ramsey

The r -size-Ramsey number of F is the minimum $e(\Gamma)$ such that $\Gamma \rightarrow_r F$.

$$\hat{r}_3(2K_2) = 4$$



Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Beck, 1983)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Friedman and Pippinger, 1987)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Haxell, Kohayakawa and Łuczak, 1995)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Clemens, Miralaei, Reding, Schacht, Taraz, 2019)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Berger, Kohayakawa, Maesaka, Martins, Mendonca, Mota, Parczyk; Kamčev, Liebenau, Wood, Yepremyan; 2021)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Rödl and Szemerédi, 2000)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.
- Graphs F with $\Delta(F) = 3$ and $\hat{r}_2(F) = \Omega(n(\log(n))^{\frac{1}{60}})$.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Tikhomirov, 2023)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.
- Graphs F with $\Delta(F) = 3$ and $\hat{r}_2(F) = \Omega(n(\log(n))^{\frac{1}{60}})$

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.
- Graphs F with $\Delta(F) = 3$ and $\hat{r}_2(F) = \Omega(n(\log(n))^{\frac{1}{60}})$
- Changes in the upper bound: $\hat{r}_r(F) = O(n^{2 - \frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}})$.

Bounds

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Nenadov, 2016)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.
- Graphs F with $\Delta(F) = 3$ and $\hat{r}_2(F) = \Omega(n(\log(n))^{\frac{1}{60}})$
- Changes in the upper bound: $\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}}(\log n)^{\frac{1}{\Delta}})$.
 - Extended for $K_{\Delta+1}$ -free F .

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Conlon, Nenadov and Trujić)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.
- Graphs F with $\Delta(F) = 3$ and $\hat{r}_2(F) = \Omega(n(\log(n))^{\frac{1}{60}})$
- Changes in the upper bound: $\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}}(\log n)^{\frac{1}{\Delta}})$.
 - Extended for $K_{\Delta+1}$ -free F .
 - Improved for $\Delta = 3$ to $O(n^{\frac{8}{5}})$.

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Draganić and Petrova, 2023)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.
- Graphs F with $\Delta(F) = 3$ and $\hat{r}_2(F) = \Omega(n(\log(n))^{\frac{1}{60}})$
- Changes in the upper bound: $\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}}(\log n)^{\frac{1}{\Delta}})$.
 - Extended for $K_{\Delta+1}$ -free F .
 - Improved for $\Delta = 3$ to $O(n^{\frac{3}{2}+o(1)})$.

Thm. (Chvátal, Rödel, Szemerédi and Trotter, 1983)

If $\Delta(F) \leq \Delta$, then:

$$e(F) \leq \hat{r}_r(F) \leq O(n^2).$$

A lot of research went to improving these bounds.

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

- We have $\hat{r}_r(F) \sim e(F)$ when F is a path, or a tree, or a cycle, or a power of a path, or has bounded treewidth.
- Graphs F with $\Delta(F) = 3$ and $\hat{r}_2(F) = \Omega(n(\log(n))^{\frac{1}{60}})$
- Changes in the upper bound: $\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}})$.
 - Extended for $K_{\Delta+1}$ -free F .
 - Improved for $\Delta = 3$ to $O(n^{\frac{3}{2}+o(1)})$.

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 1: Do it graph by graph.



Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 2 (*stronger*) : Find a graph that works for all.

Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2 - \frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 2 (*stronger*): Find a graph that works for all.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2 - \frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 2 (*stronger*): Find a graph that works for all.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2 - \frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 2 (*stronger*): Find a graph that works for all.



Partition Universality


Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2 - \frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_2(\{2K_2, K_3, C_4\})$?

Opt. 2 (*stronger*): Find a graph that works for all.



Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_r(\mathcal{G}(\Delta, n))$?

Opt. 2 (*stronger*) : Find a graph that works for all.

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

There is Γ ($e(\Gamma) = \dots$) s.t. $\Gamma \rightarrow_r F$ for each $F \in \mathcal{G}(\Delta, n)$.

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_r(\mathcal{G}(\Delta, n))$?

Opt. 3 (*stronger +*) : Find a graph s.t. for each c , one colour works for all.

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_r(\mathcal{G}(\Delta, n))$?

Opt. 3 (*stronger +*): Find a graph s.t. for each c , one colour works for all.

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

There is a graph Γ ($e(\Gamma) = \dots$) s.t. any r -colouring has a colour class χ containing $\mathcal{G}(\Delta, n)$.

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2 - \frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_r(\mathcal{G}(\Delta, n))$?

Opt. 3 (*stronger +*): Find a graph s.t. for each c , one colour works for all.

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

There is a graph Γ ($e(\Gamma) = \dots$) s.t. any r -colouring has a colour class χ containing $\mathcal{G}(\Delta, n)$.

Partition universality

We say that Γ is r -partition universal for \mathcal{G} .

Partition Universality

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

Any graph F in $\mathcal{G}(\Delta, n)$ has

$$\hat{r}_r(F) = O(n^{2-\frac{1}{\Delta}} (\log n)^{\frac{1}{\Delta}}).$$

Q: How would you upper bound $\hat{r}_r(\mathcal{G}(\Delta, n))$?

Opt. 3 (*stronger +*): Find a graph s.t. for each c , one colour works for all.

Thm. (Kohayakawa, Rödel, Schacht and Szemerédi, 2011)

For the appropriate **random graph** $G(N, p)$, a.a.s. any r -colouring has a colour class χ containing $\mathcal{G}(\Delta, n)$.

Partition universality

We say that Γ is r -partition universal for \mathcal{G} .

Summary

Summary

While studying upper bounds for $\hat{r}_r(\mathcal{G}(\Delta, n))$ one realises that the p for which we can prove $G(N, p)$ is r size-Ramsey for $\mathcal{G}(\Delta, n)$ are the same for which $G(N, p)$ is r -partition universal for $\mathcal{G}(\Delta, n)$.

Our results

GOAL: Study partition universality properties of random graphs.

Our results

GOAL: Study partition universality properties of random graphs.

Thm. (Allen, Böttcher, 2022)

A.a.s. $G(N, N^{\mu - \frac{1}{\Delta-1}})$ is r -partition universal for $\mathcal{G}(\Delta, cN)$.

Our results

GOAL: Study partition universality properties of random graphs.

Thm. (Allen, Böttcher, 2022)

A.a.s. $G(N, N^{\mu - \frac{1}{\Delta-1}})$ is r -partition universal for $\mathcal{G}(\Delta, cN)$.

Rem: Better bounds for $\Delta = 3$.

Our results

GOAL: Study partition universality properties of random graphs.

Thm. (Allen, Böttcher, 2022)

A.a.s. $G(N, N^{\mu - \frac{1}{\Delta-1}})$ is r -partition universal for $\mathcal{G}(\Delta, cN)$.

Coro. (Allen, Böttcher, 2022)

For any $F \in \mathcal{G}(\Delta, n)$,

$$\hat{r}_r(F) = O(n^{2+\mu - \frac{1}{\Delta-1}}).$$

Our results

GOAL: Study partition universality properties of random graphs.

Lemma/Thm. (Allen, Böttcher, 2022)

A.a.s. $G(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}(D, \Delta, cN)$.

Our results

GOAL: Study partition universality properties of random graphs.

Lemma/Thm. (Allen, Böttcher, 2022)

A.a.s. $G(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}(D, \Delta, cN)$.

Rem: By first moment method we cannot take $p = o(N^{-\frac{1}{D}})$.

Our results

GOAL: Study partition universality properties of random graphs.

Lemma/Thm. (Allen, Böttcher, 2022)

A.a.s. $G(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}(D, \Delta, cN)$.

Rem: By first moment method we cannot take $p = o(N^{-\frac{1}{D}})$.

Thm. (Allen, Böttcher, M.C., 2023+)

A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.

Idea of the proof


Want

A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.

Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want



A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want



A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want



A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want



A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want



A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want



A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Want


A.a.s. $G^{(k)}(N, N^{\mu - \frac{1}{D}})$ is r -partition universal for $\mathcal{G}^{(k)}(D, \Delta, cN)$.



Idea of the proof

Goal


Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .



Idea of the proof

Goal


Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .



Idea of the proof

Goal


Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .



Idea of the proof

Goal


Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .



Idea of the proof

Goal


Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .



Idea of the proof

Goal

Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .




We must be careful when embedding $N^-(y)$

Idea of the proof

Goal

Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .




When embedding x we need to consider y .

Idea of the proof

Goal

Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .



We won't embed x to *dangerous* vertices, but there are very few of them.

Idea of the proof

Goal

Find $\psi_0, \dots, \psi_{v(F)}$ growing seq. of partial homom. from F to G .

