

The Johansson-Molloy Theorem

E. Hurley, F. Pirot

arxiv.org/abs/2109.15215

A. Martinsson



arxiv.org/abs/2111.06214

UCL-LSE Reading group



Graph colouring


Let G be a graph, a **colouring** of G is a way of assigning a label (colour) to each vertex of G in such a way that the same label is never assigned to adjacent vertices.



The **chromatic number** of G (denoted with $\chi(G)$) is the number of colours required to colour G .


List colouring

Let G be a graph, a **list** for G is a way of assigning for each vertex of G a set of possible colours to use in the colouring.




List colouring

Let G be a graph, a **list** for G is a way of assigning for each vertex of G a set of possible colours to use in the colouring.




List colouring

Let G be a graph, a **list** for G is a way of assigning for each vertex of G a set of possible colours to use in the colouring.




List colouring

Let G be a graph, a **list** for G is a way of assigning for each vertex of G a set of possible colours to use in the colouring.



List colouring

Let G be a graph, a **list** for G is a way of assigning for each vertex of G a set of possible colours to use in the colouring.



The **list chromatic number** of G (denoted with $\chi_\ell(G)$) is the minimum k such that any list in which each vertex gets at least k element allows an L -proper colouring. It holds $\chi(G) \leq \chi_\ell(G) \leq \Delta(G) + 1$.

The problem at hand

Thm (Brooks, 1941)

If G is a connected graph, not an odd cycle or a clique, $\chi(G) \leq \Delta(G)$.

The problem at hand

Thm (Brooks, 1941)

If G is a connected graph, not an odd cycle or a clique, $\chi(G) \leq \Delta(G)$.

Remark

The problem of colouring is not local. There is no obvious reason for which increasing the girth should decrease the chromatic number.

The problem at hand

Thm (Brooks, 1941)

If G is a connected graph, not an odd cycle or a clique, $\chi(G) \leq \Delta(G)$.

Remark

The problem of colouring is not local. There is no obvious reason for which increasing the girth should decrease the chromatic number.

Today we examine what happens when we remove/reduce the number of triangles in the graph.

Why is this interesting?

If the pentagon conjecture is true, then for any 3-regular graph of high girth G we have $\chi(G) \leq 5$.

Original result and history

Theorem (Johansson, 1996)

$$\lim_{\Delta \rightarrow \infty} \max \{ \chi(G) : \Delta(G) = \Delta \text{ and } G \text{ is } \triangle - \text{free} \} = O\left(\frac{\Delta}{\ln(\Delta)}\right).$$

Original result and history

Theorem (Johansson, 1996)

$$\lim_{\Delta \rightarrow \infty} \max \{ \chi(G) : \Delta(G) = \Delta \text{ and } G \text{ is } \triangle - \text{free} \} = O\left(\frac{\Delta}{\ln(\Delta)}\right).$$

Theorem (Molloy, 2019)

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle - \text{free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1+\epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Original result and history

Theorem (Johansson, 1996)

$$\lim_{\Delta \rightarrow \infty} \max \{ \chi(G) : \Delta(G) = \Delta \text{ and } G \text{ is } \triangle - \text{free} \} = O\left(\frac{\Delta}{\ln(\Delta)}\right).$$

Theorem (Molloy, 2019)

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle - \text{free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1+\epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Theorem (Bernshteyn, 2019)

Same result, but using Lovász Local Lemma instead of entropy compression.

Results of the papers

Theorem (Hurley & Pirot, 2021+; Martinsson, 2021+)

Same result, but using expectation instead of Lovász Local Lemma.

Results of the papers

Theorem (Hurley & Pirot, 2021+; Martinsson, 2021+)

Same result, but using expectation instead of Lovász Local Lemma.

Theorem (Hurley & Pirot, 2021+) ★

Assume $\forall v \in V(G)$, $\bar{d}(G[N(v)]) = d \leq \frac{\Delta}{6} - 1$ and L a list such that

$$\forall v \in V(G), \quad |L(v)| \geq \left(1 + \frac{2}{\ln\left(\frac{\Delta(G)}{d+1}\right)}\right) \frac{\deg(v)}{W\left(\frac{\deg(v)}{(d+1)\ln^3\left(\frac{\Delta(G)}{d+1}\right)}\right)} ★$$

Then there are at least $\left((d+1)\ln^3\left(\frac{\Delta(G)}{d+1}\right)\right)^{|G|}$ proper L -colourings of G .

Results of the papers

Theorem (Hurley & Pirot, 2021+)

Assume $\forall v \in V(G)$, $\bar{d}(G[N(v)]) = d \leq \frac{\Delta}{6} - 1$ and L a list such that

$$\forall v \in V(G), |L(v)| \geq \left(1 + \frac{2}{\ln\left(\frac{\Delta(G)}{d+1}\right)}\right) \frac{\deg(v)}{W\left(\frac{\deg(v)}{(d+1)\ln^3\left(\frac{\Delta(G)}{d+1}\right)}\right)}.$$

Then there are at least $\left((d+1)\ln^3\left(\frac{\Delta(G)}{d+1}\right)\right)^{|G|}$ proper L -colourings of G .

Coro (Hurley & Pirot, 2021+)

Let $\Delta(G) \leq \Delta$ and $\forall v \in V$, $e(G[N(v)]) \leq \frac{\Delta^2}{f}$. Then

$$\chi_\ell(G) \leq (1 + o(1)) \frac{\Delta}{\ln(\min\{\Delta, f\})}.$$

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangleleft\text{-free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle\text{-free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Let G be a \triangle -free graph with $\Delta := \Delta(G) \geq \Delta_\epsilon$ TBA. Let $k := \left\lceil \frac{(1+\epsilon)\Delta}{\ln \Delta} \right\rceil$ and for any H , let $\mathcal{C}(H)$ be the set of proper k -colourings of H . Let $\ell := \ln^2(\Delta)$.

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle - \text{free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Let G be a \triangle -free graph with $\Delta := \Delta(G) \geq \Delta_\epsilon$ TBA. Let $k := \left\lceil \frac{(1+\epsilon)\Delta}{\ln \Delta} \right\rceil$ and for any H , let $\mathcal{C}(H)$ be the set of proper k -colourings of H . Let $\ell := \ln^2(\Delta)$.

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle\text{-free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Let G be a \triangle -free graph with $\Delta := \Delta(G) \geq \Delta_\epsilon$ TBA. Let $k := \left\lceil \frac{(1+\epsilon)\Delta}{\ln \Delta} \right\rceil$ and for any H , let $\mathcal{C}(H)$ be the set of proper k -colourings of H . Let $\ell := \ln^2(\Delta)$.

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. The case $H = \{v\}$ being trivial.

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle\text{-free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Let G be a \triangle -free graph with $\Delta := \Delta(G) \geq \Delta_\epsilon$ TBA. Let $k := \left\lceil \frac{(1+\epsilon)\Delta}{\ln \Delta} \right\rceil$ and for any H , let $\mathcal{C}(H)$ be the set of proper k -colourings of H . Let $\ell := \ln^2(\Delta)$.

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle - \text{free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Let G be a \triangle -free graph with $\Delta := \Delta(G) \geq \Delta_\epsilon$ TBA. Let $k := \left\lceil \frac{(1+\epsilon)\Delta}{\ln \Delta} \right\rceil$ and for any H , let $\mathcal{C}(H)$ be the set of proper k -colourings of H . Let $\ell := \ln^2(\Delta)$.

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

$$\textcircled{1} \quad \mathbb{E}_{\mathcal{C}(H \setminus v)} [|L_c(v)|] = \frac{\sum_{c \in \mathcal{C}(H \setminus v)} |L_c(v)|}{|\mathcal{C}(H \setminus v)|} = \frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|}.$$

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle - \text{free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Let G be a \triangle -free graph with $\Delta := \Delta(G) \geq \Delta_\epsilon$ TBA. Let $k := \left\lceil \frac{(1+\epsilon)\Delta}{\ln \Delta} \right\rceil$ and for any H , let $\mathcal{C}(H)$ be the set of proper k -colourings of H . Let $\ell := \ln^2(\Delta)$.

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

$$\textcircled{1} \mathbb{E}_{\mathcal{C}(H \setminus v)} [|L_c(v)|] = \frac{\sum_{c \in \mathcal{C}(H \setminus v)} |L_c(v)|}{|\mathcal{C}(H \setminus v)|} = \frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|}.$$

$$\textcircled{2} \text{ For } t \in \mathbb{N} \text{ TBA and } u \in N(v): \mathbb{P}_{\mathcal{C}(H \setminus v)} [|L_c(u)| \leq t] \stackrel{\star}{\leq} \frac{t \cdot |\mathcal{C}(H \setminus \{u, v\})|}{|\mathcal{C}(H \setminus \{v\})|} \leq \frac{t}{\ell}.$$

Quick proof of Johansson-Molloy Theorem

Theorem

$$\forall \epsilon > 0 \exists \Delta_\epsilon \in \mathbb{N} \forall \triangle\text{-free } G, \Delta(G) \geq \Delta_\epsilon \implies \chi(G) \leq (1 + \epsilon) \frac{\Delta(G)}{\ln(\Delta(G))}.$$

Let G be a \triangle -free graph with $\Delta := \Delta(G) \geq \Delta_\epsilon$ TBA. Let $k := \left\lceil \frac{(1+\epsilon)\Delta}{\ln \Delta} \right\rceil$ and for any H , let $\mathcal{C}(H)$ be the set of proper k -colourings of H . Let $\ell := \ln^2(\Delta)$.

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

$$\textcircled{1} \mathbb{E}_{\mathcal{C}(H \setminus v)} [|L_c(v)|] = \frac{\sum_{c \in \mathcal{C}(H \setminus v)} |L_c(v)|}{|\mathcal{C}(H \setminus v)|} = \frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|}.$$

$$\textcircled{2} \text{ For } t \in \mathbb{N} \text{ TBA and } u \in N(v): \mathbb{P}_{\mathcal{C}(H \setminus v)} [|L_c(u)| \leq t] \leq \frac{t \cdot |\mathcal{C}(H \setminus \{u, v\})|}{|\mathcal{C}(H \setminus \{v\})|} \leq \frac{t}{\ell}.$$

If $1 \ll t \ll \frac{\ell}{\ln(\Delta)}$ Markov's gives $|\{u \in N(v) : |L_c(u)| \leq t\}| \leq o(k)$ whp.

Quick proof of Johansson-Molloy Theorem

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

$$\textcircled{1} \quad \mathbb{E}_{\mathcal{C}(H \setminus v)} [|L_c(v)|] = \frac{\sum_{c \in \mathcal{C}(H \setminus v)} |L_c(v)|}{|\mathcal{C}(H \setminus v)|} = \frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|}.$$

$$\textcircled{2} \quad \text{For } t \in \mathbb{N} \text{ TBA and } u \in N(v): \mathbb{P}_{\mathcal{C}(H \setminus v)} [|L_c(u)| \leq t] \leq \frac{t \cdot |\mathcal{C}(H \setminus \{u, v\})|}{|\mathcal{C}(H \setminus \{v\})|} \leq \frac{t}{\ell}.$$

Quick proof of Johansson-Molloy Theorem

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

$$\textcircled{1} \quad \mathbb{E}_{\mathcal{C}(H \setminus v)} [|L_c(v)|] = \frac{\sum_{c \in \mathcal{C}(H \setminus v)} |L_c(v)|}{|\mathcal{C}(H \setminus v)|} = \frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|}.$$

$$\textcircled{2} \quad \text{For } t \in \mathbb{N} \text{ TBA and } u \in N(v): \mathbb{P}_{\mathcal{C}(H \setminus v)} [|L_c(u)| \leq t] \leq \frac{t \cdot |\mathcal{C}(H \setminus \{u, v\})|}{|\mathcal{C}(H \setminus \{v\})|} \leq \frac{t}{\ell}.$$

$$\star \textcircled{3} \quad |\{u \in N(v) : |L_c(u)| \leq t\}| \leq o(k) \text{ whp.}$$

Quick proof of Johansson-Molloy Theorem

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

- ① $\mathbb{E}_{\mathcal{C}(H \setminus v)} [|L_c(v)|] = \frac{\sum_{c \in \mathcal{C}(H \setminus v)} |L_c(v)|}{|\mathcal{C}(H \setminus v)|} = \frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|}$.
- ② For $t \in \mathbb{N}$ TBA and $u \in N(v)$: $\mathbb{P}_{\mathcal{C}(H \setminus v)} [|L_c(u)| \leq t] \leq \frac{t \cdot |\mathcal{C}(H \setminus \{u, v\})|}{|\mathcal{C}(H \setminus \{v\})|} \leq \frac{t}{\ell}$.
- ③ $|\{u \in N(v) : |L_c(u)| \leq t\}| \leq o(k)$ whp.
- ④ $L_c(u)$ is determined by $c|_{H \setminus \{v, N(v)\}}$ since H is \triangle -free.

Quick proof of Johansson-Molloy Theorem

Claim

ISTS that for any $H \subseteq G$ induced, $\forall v \in V(H)$ we have $\frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|} \geq \ell$.

Proof: By induction. For $c : V(H) \rightarrow [k] \cup \{\text{NaN}\}$ let $L_c(u) := [k] \setminus c(N(u))$.

- ① $\mathbb{E}_{\mathcal{C}(H \setminus v)} [|L_c(v)|] = \frac{\sum_{c \in \mathcal{C}(H \setminus v)} |L_c(v)|}{|\mathcal{C}(H \setminus v)|} = \frac{|\mathcal{C}(H)|}{|\mathcal{C}(H \setminus v)|}$.
- ② For $t \in \mathbb{N}$ TBA and $u \in N(v)$: $\mathbb{P}_{\mathcal{C}(H \setminus v)} [|L_c(u)| \leq t] \leq \frac{t \cdot |\mathcal{C}(H \setminus \{u, v\})|}{|\mathcal{C}(H \setminus \{v\})|} \leq \frac{t}{\ell}$.
- ③ $|\{u \in N(v) : |L_c(u)| \leq t\}| \leq o(k)$ whp.
- ④ $L_c(u)$ is determined by $c|_{H \setminus \{v, N(v)\}}$ since H is \triangle -free.
- ⑤ Conditioning of $c|_{H \setminus \{v, N(v)\}} = c_0$ each $u \in N(v)$ is coloured indep.

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof:  Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] \stackrel{\star}{=} \sum_{j \in [k] \setminus B} \prod_{\substack{L_i \ni j \\ |L_i| > t}} \left(1 - \frac{1}{|L_i|}\right)$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\begin{aligned} \mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] &= \sum_{j \in [k] \setminus B} \prod_{\substack{L_i \ni j \\ |L_i| > t}} \left(1 - \frac{1}{|L_i|}\right) \\ &\stackrel{\star}{\geq} (k - |B|) \left(\prod_{j \in [k] \setminus B} \prod_{\substack{L_i \ni j \\ |L_i| > t}} \left(1 - \frac{1}{|L_i|}\right) \right)^{\frac{1}{k - |B|}} \end{aligned}$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\begin{aligned} \mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] &\geq (k - |B|) \left(\prod_{j \in [k] \setminus B} \prod_{\substack{L_i \ni j \\ |L_i| > t}} \left(1 - \frac{1}{|L_i|} \right) \right)^{\frac{1}{k - |B|}} \\ &\geq \prod_{i: |L_i| > t} \prod_{j \in L_i \setminus B} \left(1 - \frac{1}{|L_i|} \right) \end{aligned}$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\begin{aligned} \mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] &\geq (k - |B|) \left(\prod_{j \in [k] \setminus B} \prod_{\substack{L_i \ni j \\ |L_i| > t}} \left(1 - \frac{1}{|L_i|} \right) \right)^{\frac{1}{k - |B|}} \\ &\geq \prod_{i: |L_i| > t} \prod_{j \in L_i \setminus B} \left(1 - \frac{1}{|L_i|} \right) \\ &\geq \left(\left(1 - \frac{1}{t} \right)^t \right)^\Delta \end{aligned}$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\begin{aligned} \mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] &\geq (k - |B|) \left(\prod_{j \in [k] \setminus B} \prod_{\substack{L_i \ni j \\ |L_i| > t}} \left(1 - \frac{1}{|L_i|} \right) \right)^{\frac{1}{k - |B|}} \\ &\geq \left(\left(1 - \frac{1}{t} \right)^t \right)^\Delta \\ &\geq \exp \left(- \left(1 + O \left(\frac{1}{t} \right) \right) \Delta \right) \end{aligned}$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\begin{aligned} \mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] &\geq (k - |B|) \left(\prod_{j \in [k] \setminus B} \prod_{\substack{L_i \ni j \\ |L_i| > t}} \left(1 - \frac{1}{|L_i|} \right) \right)^{\frac{1}{k - |B|}} \\ &\geq (k - |B|) \exp \left(- \frac{(1 + o(1))\Delta}{k - |B|} \right) \end{aligned}$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\begin{aligned}\mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] &\geq (k - |B|) \exp \left(-\frac{(1 + o(1))\Delta}{k - |B|} \right) \\ &\geq \Theta \left(\frac{\Delta}{\ln \Delta} \right) \exp \left(-\frac{1 + o(1)}{(1 + \epsilon)} \ln \Delta \right)\end{aligned}$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\begin{aligned} \mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] &\geq \Theta \left(\frac{\Delta}{\ln \Delta} \right) \exp \left(-\frac{1 + o(1)}{(1 + \epsilon)} \ln \Delta \right) \\ &\geq \Delta^{\frac{\epsilon}{(1 + \epsilon)} - o(1)} \end{aligned}$$

Quick proof of Johansson-Molloy Theorem

Lemma

Let $L_1, \dots, L_d \subseteq [k]$ for $d \leq \Delta$, s.t. all but $o(k)$ of the L_i have $|L_i| > t$. Let $X_i \in_u L_i$ be taken independently. Let $X := [k] \setminus \{X_1, \dots, X_d\}$, then whp $|X| \gg \ell$.

Proof: Let $B := \{i : |L_i| \leq t\}$. Fix $\{X_i\}_{i \in B}$ and condition on this event.

$$\mathbb{E}_{\mathcal{C}(H \setminus v) | \{X_i\}_{i \in B}} [|X|] \geq \Delta \frac{\epsilon}{(1+\epsilon)} - o(1)$$

Analysis of the extension

Theorem (Hurley & Pirot, 2021+)

Assume $\forall v \in V(G)$, $\bar{d}(G[N(v)]) = d \leq \frac{\Delta}{6} - 1$ and L a list such that

$$\forall v \in V(G), \quad |L(v)| \geq \left(1 + \frac{2}{\ln\left(\frac{\Delta(G)}{d+1}\right)}\right) \frac{\deg(v)}{W\left(\frac{\deg(v)}{(d+1)\ln^3\left(\frac{\Delta(G)}{d+1}\right)}\right)}.$$

Then there are at least $\left((d+1)\ln^3\left(\frac{\Delta(G)}{d+1}\right)\right)^{|G|}$ proper L -colourings of G .

