

Product Schur Triples in the Integers

L. Mattos, **D. Mergoni Cecchelli**, O. Parczyk



Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.



Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.



Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.



Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

The only known values are $S(1) = 2$, $S(2) = 5$, $S(3) = 14$, $S(4) = 45$ and $S(5) = 161$.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

The only known values are $S(1) = 2$, $S(2) = 5$, $S(3) = 14$, $S(4) = 45$ and $S(5) = 161$.

Thm. (Abbott and Moser, 1966)

For any k and l positive integers we have:

$$S(k+l) \geq 2S(k)S(l) + S(k) + S(l).$$

Paired with $S(5) = 161$ (Heule, 2018), this gives $S(k) \geq c \cdot 321^{k/5}$.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

The only known values are $S(1) = 2$, $S(2) = 5$, $S(3) = 14$, $S(4) = 45$ and $S(5) = 161$.

Corollary. (Abbott and Moser, 1966)

Finding additional values (e.g. $S(6)$) would improve the lower bound.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

The only known values are $S(1) = 2$, $S(2) = 5$, $S(3) = 14$, $S(4) = 45$ and $S(5) = 161$.

Corollary. (Abbott and Moser, 1966)

Finding additional values (e.g. $S(6)$) would improve the lower bound.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

The only known values are $S(1) = 2$, $S(2) = 5$, $S(3) = 14$, $S(4) = 45$ and $S(5) = 161$.

Corollary. (Abbott and Moser, 1966)

Finding additional values (e.g. $S(6)$) would improve the lower bound.

The upper bound is $\lfloor k!(e - \frac{1}{24}) \rfloor$ and due to Irving (1974).

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Abbott-Wang Variation (1977)

How large is the largest subset of $[n]$ that can be partitioned into k sum-free sets?

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Abbott-Wang Variation (1977)

How large is the largest subset of $[n]$ that can be partitioned into k sum-free sets?

Abbott-Wang Conjecture (1977)

The Abbott-Wang construction is optimal. I.e. $n - \lfloor \frac{n}{H(k)} \rfloor$

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Graham, Rödl and Ruciński (1996)

Any 2-colouring of $[n]$ contains $n^2/19$ monochromatic sums.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Schoen (1999), Robertson and Zeilberger (1999)

Any 2-colouring of $[n]$ contains $n^2/11$ monochromatic sums.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Schoen (1999), Robertson and Zeilberger (1999)

Any 2-colouring of $[n]$ contains $n^2/11$ monochromatic sums.

For $k \geq 3$ only $\Theta(n^2)$ is known.

Deterministic Schur Problems

Lemma (Schur, 1917)

For any positive integer k there is a (smallest) $S(k) \in \mathbb{N}$ such that any k -colouring of $[S(k)] := \{1, \dots, S(k)\}$ contains a monochromatic sum.

We cannot partition $[S(k)]$ in k -many sum-free sets.

Schoen (1999), Robertson and Zeilberger (1999)

Any 2-colouring of $[n]$ contains $n^2/11$ monochromatic sums.

For $k \geq 3$ only $\Theta(n^2)$ is known. Which colourings attain the minimum?

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum?

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum?

- There are $\sim n^2/4$ sums in $[n]$.

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum?

- There are $\sim n^2/4$ sums in $[n]$.
- Let X count the sums in $[n]_p$ and $\mu = \mathbb{E}[X]$. We have

$$\mu = \Theta(n^2 p^3).$$

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum?

- There are $\sim n^2/4$ sums in $[n]$.
- Let X count the sums in $[n]_p$ and $\mu = \mathbb{E}[X]$. We have

$$\mu = \Theta(n^2 p^3).$$

- For $p \ll n^{-2/3}$, $\mu = o(1)$ and thus $\mathbb{P}[X \geq 1] = o(1)$.

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum?

- There are $\sim n^2/4$ sums in $[n]$.
- Let X count the sums in $[n]_p$ and $\mu = \mathbb{E}[X]$. We have

$$\mu = \Theta(n^2 p^3).$$

- For $p \ll n^{-2/3}$, $\mu = o(1)$ and thus $\mathbb{P}[X \geq 1] = o(1)$.
- For $p \gg n^{-2/3}$ we have $\text{Var}(X) \ll \mu^2$ as

$$\text{Var}(X) \sim n^2 p^3 (1 - p^3) + p^5 n^3 + p^4 n^2.$$

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum? $p \gg n^{-2/3}$.

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum? $p \gg n^{-2/3}$.

Thm. (Graham, Rödl, Ruciński, 1996)

W.h.p. any 2-colouring of $[n]_p$ with $p \gg n^{-1/2}$ contains $\Theta(n^2 p^3)$ monochromatic sums.

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum? $p \gg n^{-2/3}$.

Thm. (Graham, Rödl, Ruciński, 1996)

W.h.p. any 2-colouring of $[n]_p$ with $p \gg n^{-1/2}$ contains $\Theta(n^2 p^3)$ monochromatic sums.

Question

Let $\alpha_n \in (0, 1)$, what p_n guarantees that if $|C_n| \geq (1 - \alpha_n)n$ then

$$\lim_{n \rightarrow \infty} \mathbb{P}[C_n \cup [n]_{p_n} \text{ is 2-Schur}] = 1?$$

Random Schur Problems

Question

For which values of p does $[n]_p$ contain a sum? $p \gg n^{-2/3}$.

Thm. (Graham, Rödl, Ruciński, 1996)

W.h.p. any 2-colouring of $[n]_p$ with $p \gg n^{-1/2}$ contains $\Theta(n^2 p^3)$ monochromatic sums.

Thm. (Das, Knierim, and Morris, 2024)

If C is dense and $p \gg n^{-2/3}$, then w.h.p. every 2-colouring of $C \cup [n]_p$ contains a monochromatic sum.

Summary.

Summary.

- Value of $S(k)$ difficult to bound, impossible to determine.

Summary.

- Value of $S(k)$ difficult to bound, impossible to determine.
- Largest subsets of $[n]$ that can be k -partitioned in sum-free sets.

Summary.

- Value of $S(k)$ difficult to bound, impossible to determine.
- Largest subsets of $[n]$ that can be k -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?

Summary.

- Value of $S(k)$ difficult to bound, impossible to determine.
- Largest subsets of $[n]$ that can be k -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?
- k -colourings of $[n]$ with minimal number of monochromatic sums.

Summary.

- Value of $S(k)$ difficult to bound, impossible to determine.
- Largest subsets of $[n]$ that can be k -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?
- k -colourings of $[n]$ with minimal number of monochromatic sums.
- Which $[n]_p$ cannot be partitioned into k sum-free sets?

Summary.

- Value of $S(k)$ difficult to bound, impossible to determine.
- Largest subsets of $[n]$ that can be k -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?
- k -colourings of $[n]$ with minimal number of monochromatic sums.
- Which $[n]_p$ cannot be partitioned into k sum-free sets?
- What's the interplay between deterministic and random?

What about PRODUCTS?

Possible directions.

Possible directions.

- Largest subsets of $[n]$ that can be k -partitioned in product-free sets.

Possible directions.

- Largest subsets of $[n]$ that can be k -partitioned in product-free sets.
- k -colourings of $[n]$ with few monochromatic products.

Possible directions.

- Largest subsets of $[n]$ that can be k -partitioned in product-free sets.
- k -colourings of $[n]$ with few monochromatic products.
- Which $[2, n]_p$ cannot be partitioned into k product-free sets?

Possible directions.

- Largest subsets of $[n]$ that can be k -partitioned in product-free sets.
- k -colourings of $[n]$ with few monochromatic products.
- Which $[2, n]_p$ cannot be partitioned into k product-free sets?
- What's the interplay between deterministic and random?

Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

For n large enough, any r -colouring of $[2, n]$ contains $n^{1/S(r-1)}$ monochromatic products.

Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

For n large enough, any r -colouring of $[2, n]$ contains $n^{1/S(r-1)}$ monochromatic products.

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

For n large enough, any r -colouring of $[2, n]$ contains $n^{1/S(r-1)}$ monochromatic products.

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Thm. (Mattos, MC, Parczyk, 2025)

$\hat{p}_\alpha(n) = n^{-1/2+o(1)}$ is the threshold for the α -randomly perturbed product Schur property (for α in a wide range).

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Upper bound first.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Upper bound first.

- For a , let $P(a) := \{a^i : i = 1, \dots, S(k)\}$.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Upper bound first.

- For a , let $P(a) := \{a^i : i = 1, \dots, S(k)\}$.
- $P(a)$ cannot be partitioned into k product-free sets.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Upper bound first.

- For a , let $P(a) := \{a^i : i = 1, \dots, S(k)\}$.
- $P(a)$ cannot be partitioned into k product-free sets.
- Fix $A' = [\frac{1}{2}\varepsilon n^{1/S(k)}, n^{1/S(k)}]$.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Upper bound first.

- For a , let $P(a) := \{a^i : i = 1, \dots, S(k)\}$.
- $P(a)$ cannot be partitioned into k product-free sets.
- Fix $A' = [\frac{1}{2}\varepsilon n^{1/S(k)}, n^{1/S(k)}]$.
- For $a, b \in A'$ distinct, $P(a) \cap P(b) = \emptyset$ and $P(a), P(b) \subseteq [n]$.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Upper bound first.

- For a , let $P(a) := \{a^i : i = 1, \dots, S(k)\}$.
- $P(a)$ cannot be partitioned into k product-free sets.
- Fix $A' = [\frac{1}{2}\varepsilon n^{1/S(k)}, n^{1/S(k)}]$.
- For $a, b \in A'$ distinct, $P(a) \cap P(b) = \emptyset$ and $P(a), P(b) \subseteq [n]$.
- Any subset of $[n]$ that can be partitioned into k product-free sets must avoid an element of $P(a)$ for each a in A' .

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Lower bound next.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Lower bound next.

- Let $\chi : [S'(k) - 1] \rightarrow [k]$ without monochromatic $a + b = c$ or $a + b = c - 1$.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Lower bound next.

- Let $\chi : [S'(k) - 1] \rightarrow [k]$ without monochromatic $a + b = c$ or $a + b = c - 1$.
- Colour $a \in (n^{1/S'(k)}, n]$ with colour $\chi(\lceil S'(k) \cdot \log_n(a) \rceil - 1)$.

Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$ and $k \in \mathbb{N}^+$. For n large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Proof. Lower bound next.

- Let $\chi : [S'(k) - 1] \rightarrow [k]$ without monochromatic $a + b = c$ or $a + b = c - 1$.
- Colour $a \in (n^{1/S'(k)}, n]$ with colour $\chi(\lceil S'(k) \cdot \log_n(a) \rceil - 1)$.
- If $ab = c$, then let $a' = \lceil S'(k) \cdot \log_n(a) \rceil - 1$, $b' = \lceil S'(k) \cdot \log_n(b) \rceil - 1$, and $c' = \lceil S'(k) \cdot \log_n(c) \rceil - 1$ and note that $\log_n(a) + \log_n(b) = \log_n(c)$ implies $a' + b' = c'$ or $a' + b' = c' - 1$.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.

b_2   r_2

b_1   r_1


Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.




Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.




Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.




Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.



Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.
- To each $\{r_1, r_2, b_1, b_2\}$ with $r_1, r_2 \in R$, $b_1, b_2 \in B$, $r_1 r_2$ blue and $b_1 b_2$ red we can associate a monochromatic product.

Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let $\varepsilon > 0$. For n large enough, any 2-colouring of $[2, n]$ contains $n^{1/3-\varepsilon}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.
- To each $\{r_1, r_2, b_1, b_2\}$ with $r_1, r_2 \in R, b_1, b_2 \in B, r_1 r_2$ blue and $b_1 b_2$ red we can associate a monochromatic product.
- Take care of double-counting.

Deterministic Product Schur II

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

For n large enough, any r -colouring of $[2, n]$ contains $n^{1/S(r-1)}$ monochromatic products.

Proof.

- **Lemma.** If $A \subseteq [2, n]$ has size $n - \frac{1}{2}\sqrt{n}$, it contains $n/8$ products.
- Let R be the red elements of $[n^{1/3}]$, B the blue ones. By Lemma, wlog we have $|R| \geq |B| \geq n^{1/6}/2$.
- Let $P_R := \{ab : a, b \in R\}$ and $P_B := \{ab : a, b \in B\}$.
- Note $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$. Also, P_R contains $n^{1/3-\varepsilon}$ blue elements and P_B contains $n^{1/3-\varepsilon}$ red elements.
- To each $\{r_1, r_2, b_1, b_2\}$ with $r_1, r_2 \in R$, $b_1, b_2 \in B$, $r_1 r_2$ blue and $b_1 b_2$ red we can associate a monochromatic product.
- Take care of double-counting.

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Lower bound first.

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Lower bound first.

- The number of products is

$$\sum_{a=2}^{\sqrt{n}} \frac{n}{a} + O(n) = (1 + o(1))n \log(n).$$

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Lower bound first.

- The number of products is

$$\sum_{a=2}^{\sqrt{n}} \frac{n}{a} + O(n) = (1 + o(1))n \log(n).$$

- The expected number of products in $[2, n]_p$ is:

$$p^3(1 + o(1))n \log(n) + p^2\sqrt{n}.$$

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Lower bound first.

- The number of products is

$$\sum_{a=2}^{\sqrt{n}} \frac{n}{a} + O(n) = (1 + o(1))n \log(n).$$

- The expected number of products in $[2, n]_p$ is:

$$p^3(1 + o(1))n \log(n) + p^2\sqrt{n}.$$

- For $p \ll (n \log(n))^{-1/3}$, there are no products.

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Upper bound next (very sketched).

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Upper bound next (very sketched).

- Take $p = (n \log(n))^{-1/3} \omega(n)$, and let q be such that $(1 - q)^2 = (1 - p)$.

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Upper bound next (very sketched).

- Take $p = (n \log(n))^{-1/3} \omega(n)$, and let q be such that $(1 - q)^2 = (1 - p)$.
- Let $A, B := [2, n]_q$, and $C = A \cup B$. Note $C \sim [2, n]_p$.

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Upper bound next (very sketched).

- Take $p = (n \log(n))^{-1/3} \omega(n)$, and let q be such that $(1 - q)^2 = (1 - p)$.
- Let $A, B := [2, n]_q$, and $C = A \cup B$. Note $C \sim [2, n]_p$.
- This reduces to show $|A^2 \cap [2, n]| \gg 1/q$ w.h.p.

Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for $[2, n]_p$ to contain a product is $(n \log(n))^{-1/3}$.

Proof. Upper bound next (very sketched).

- Take $p = (n \log(n))^{-1/3} \omega(n)$, and let q be such that $(1 - q)^2 = (1 - p)$.
- Let $A, B := [2, n]_q$, and $C = A \cup B$. Note $C \sim [2, n]_p$.
- This reduces to show $|A^2 \cap [2, n]| \gg 1/q$ w.h.p.
- A useful tool is that no c can be written as the product of elements of A in more than 2 ways w.h.p.

Open questions.

Open questions.

- Is there an ε such that $S'(k) < (1 - \varepsilon)S(k)$ for large k ?

Open questions.

- Is there an ε such that $S'(k) < (1 - \varepsilon)S(k)$ for large k ?
- What is the k -colouring of n with the fewest monochromatic products?

Open questions.

- Is there an ε such that $S'(k) < (1 - \varepsilon)S(k)$ for large k ?
- What is the k -colouring of n with the fewest monochromatic products?
- What is the threshold for any k -colouring of $[2, n]_p$ to contain a monochromatic product?

