

# Product Schur Triples in the Integers

L. Mattos, **D. Mergoni Cecchelli**, O. Parczyk



# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.



# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.



# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.



# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

The only known values are  $S(1) = 2$ ,  $S(2) = 5$ ,  $S(3) = 14$ ,  $S(4) = 45$  and  $S(5) = 161$ .

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

The only known values are  $S(1) = 2$ ,  $S(2) = 5$ ,  $S(3) = 14$ ,  $S(4) = 45$  and  $S(5) = 161$ .

## Thm. (Abbott and Moser, 1966)

For any  $k$  and  $l$  positive integers we have:

$$S(k+l) \geq 2S(k)S(l) + S(k) + S(l).$$

Paired with  $S(5) = 161$  (Heule, 2018), this gives  $S(k) \geq c \cdot 321^{k/5}$ .

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

The only known values are  $S(1) = 2$ ,  $S(2) = 5$ ,  $S(3) = 14$ ,  $S(4) = 45$  and  $S(5) = 161$ .

## Corollary. (Abbott and Moser, 1966)

Finding additional values (e.g.  $S(6)$ ) would improve the lower bound.



# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

The only known values are  $S(1) = 2$ ,  $S(2) = 5$ ,  $S(3) = 14$ ,  $S(4) = 45$  and  $S(5) = 161$ .

## Corollary. (Abbott and Moser, 1966)

Finding additional values (e.g.  $S(6)$ ) would improve the lower bound.

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

The only known values are  $S(1) = 2$ ,  $S(2) = 5$ ,  $S(3) = 14$ ,  $S(4) = 45$  and  $S(5) = 161$ .

## Corollary. (Abbott and Moser, 1966)

Finding additional values (e.g.  $S(6)$ ) would improve the lower bound.

The upper bound is  $\lfloor k!(e - \frac{1}{24}) \rfloor$  and due to Irving (1974).

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

## Abbott-Wang Variation

How large is the largest subset of  $[n]$  that can be partitioned into  $k$  sum-free sets?

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

## Abbott-Wang Variation

How large is the largest subset of  $[n]$  that can be partitioned into  $k$  sum-free sets?

## Abbott-Wang Conjecture

The Abbott-Wang construction is optimal. I.e.  $n - \lfloor \frac{n}{H(k)} \rfloor$

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

## Graham, Rödl and Ruciński (1996)

Any 2-colouring of  $[n]$  contains  $n^2/19$  monochromatic Schur triples.

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

## Schoen (1999), Robertson and Zeilberger (1999)

Any 2-colouring of  $[n]$  contains  $n^2/11$  monochromatic Schur triples.



# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

## Schoen (1999), Robertson and Zeilberger (1999)

Any 2-colouring of  $[n]$  contains  $n^2/11$  monochromatic Schur triples.

For  $k \geq 3$  only  $\Theta(n^2)$  is known.

# Deterministic Schur Problems

## Lemma (Schur, 1917)

For any positive integer  $k$  there is a (smallest)  $S(k) \in \mathbb{N}$  such that any  $k$ -colouring of  $[S(k)] := \{1, \dots, S(k)\}$  contains a monochromatic sum.

We cannot partition  $[S(k)]$  in  $k$ -many sum-free sets.

## Schoen (1999), Robertson and Zeilberger (1999)

Any 2-colouring of  $[n]$  contains  $n^2/11$  monochromatic Schur triples.

For  $k \geq 3$  only  $\Theta(n^2)$  is known. Which colourings attain the minimum?

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?

- There are  $\sim n^2/4$  sums in  $[n]$ .

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?

- There are  $\sim n^2/4$  sums in  $[n]$ .
- Let  $X$  count the sums in  $[n]_p$  and  $\mu = \mathbb{E}[X]$ . We have

$$\mu = \Theta(n^2 p^3).$$

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?

- There are  $\sim n^2/4$  sums in  $[n]$ .
- Let  $X$  count the sums in  $[n]_p$  and  $\mu = \mathbb{E}[X]$ . We have

$$\mu = \Theta(n^2 p^3).$$

- For  $p \ll n^{-2/3}$ ,  $\mu = o(1)$  and thus  $\mathbb{P}[X \geq 1] = o(1)$ .

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?

- There are  $\sim n^2/4$  sums in  $[n]$ .
- Let  $X$  count the sums in  $[n]_p$  and  $\mu = \mathbb{E}[X]$ . We have

$$\mu = \Theta(n^2 p^3).$$

- For  $p \ll n^{-2/3}$ ,  $\mu = o(1)$  and thus  $\mathbb{P}[X \geq 1] = o(1)$ .
- For  $p \gg n^{-2/3}$  we have  $\text{Var}(X) \ll \mu^2$  as

$$\text{Var}(X) \sim n^2 p^3 (1 - p^3) + p^5 n^3 + p^4 n^2.$$

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?  $p \gg n^{-2/3}$ .



# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?  $p \gg n^{-2/3}$ .

## Thm. (Graham, Rödl, Ruciński, 2000)

W.h.p. any 2-colouring of  $[n]_p$  with  $p \gg n^{-1/2}$  contains  $\Theta(n^2 p^3)$  monochromatic sums.

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?  $p \gg n^{-2/3}$ .

## Thm. (Graham, Rödl, Ruciński, 2000)

W.h.p. any 2-colouring of  $[n]_p$  with  $p \gg n^{-1/2}$  contains  $\Theta(n^2 p^3)$  monochromatic sums.

## Question

Let  $\alpha_n \in (0, 1)$ , what  $p_n$  guarantees that if  $|C_n| \geq (1 - \alpha_n)n$  then

$$\lim_{n \rightarrow \infty} \mathbb{P}[C_n \cup [n]_p \text{ is 2-Schur}] = 1?$$

# Random Schur Problems

## Question

For which values of  $p$  does  $[n]_p$  contain a Schur triple?  $p \gg n^{-2/3}$ .

## Thm. (Graham, Rödl, Ruciński, 2000)

W.h.p. any 2-colouring of  $[n]_p$  with  $p \gg n^{-1/2}$  contains  $\Theta(n^2 p^3)$  monochromatic sums.

## Thm. (Das, Knierim, and Morris, 2024)

If  $C$  is dense and  $p \gg n^{-2/3}$ , then w.h.p. every 2-colouring of  $C \cup [n]_p$  contains a monochromatic sum.

## Summary.

## Summary.

- Value of  $S(k)$  difficult to bound, impossible to determine.

## Summary.

- Value of  $S(k)$  difficult to bound, impossible to determine.
- Largest subsets of  $[n]$  that can be  $k$ -partitioned in sum-free sets.

## Summary.

- Value of  $S(k)$  difficult to bound, impossible to determine.
- Largest subsets of  $[n]$  that can be  $k$ -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?

## Summary.

- Value of  $S(k)$  difficult to bound, impossible to determine.
- Largest subsets of  $[n]$  that can be  $k$ -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?
- $k$ -colourings of  $[n]$  with minimal number of monochromatic sums.



## Summary.

- Value of  $S(k)$  difficult to bound, impossible to determine.
- Largest subsets of  $[n]$  that can be  $k$ -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?
- $k$ -colourings of  $[n]$  with minimal number of monochromatic sums.
- Which  $[n]_p$  cannot be partitioned into  $k$  sum-free sets?

## Summary.

- Value of  $S(k)$  difficult to bound, impossible to determine.
- Largest subsets of  $[n]$  that can be  $k$ -partitioned in sum-free sets.
- Is modular arithmetic the best we can do?
- $k$ -colourings of  $[n]$  with minimal number of monochromatic sums.
- Which  $[n]_p$  cannot be partitioned into  $k$  sum-free sets?
- What's the interplay between deterministic and random?

What about PRODUCTS?

# Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

# Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

# Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

Any 2-colouring of  $[2, n]$  contains  $(\frac{1}{2\sqrt{2}} - o(1))n^{1/2} \log(n)$  monochromatic products.

# Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

Any 2-colouring of  $[2, n]$  contains  $(\frac{1}{2\sqrt{2}} - o(1))n^{1/2} \log(n)$  monochromatic products.

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

# Product Schur Problems

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

Any 2-colouring of  $[2, n]$  contains  $(\frac{1}{2\sqrt{2}} - o(1))n^{1/2} \log(n)$  monochromatic products.

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

Thm. (Mattos, MC, Parczyk, 2025)

$\hat{p}_\alpha(n) = n^{-1/2+o(1)}$  is the threshold for the  $\alpha$ -randomly perturbed product Schur property (for  $\alpha$  in a wide range).



# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Upper bound first.

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Upper bound first.

- For  $a$ , let  $P(a) := \{a^i : i = 1, \dots, S(k)\}$ .

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Upper bound first.

- For  $a$ , let  $P(a) := \{a^i : i = 1, \dots, S(k)\}$ .
- $P(a)$  cannot be partitioned into  $k$  product-free sets.

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Upper bound first.

- For  $a$ , let  $P(a) := \{a^i : i = 1, \dots, S(k)\}$ .
- $P(a)$  cannot be partitioned into  $k$  product-free sets.
- Fix  $A' = [\frac{1}{2}\varepsilon n^{1/S(k)}, n^{1/S(k)}]$ .

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Upper bound first.

- For  $a$ , let  $P(a) := \{a^i : i = 1, \dots, S(k)\}$ .
- $P(a)$  cannot be partitioned into  $k$  product-free sets.
- Fix  $A' = [\frac{1}{2}\varepsilon n^{1/S(k)}, n^{1/S(k)}]$ .
- For  $a, b \in A'$  distinct,  $P(a) \cap P(b) = \emptyset$  and  $P(a), P(b) \subseteq [n]$ .

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Upper bound first.

- For  $a$ , let  $P(a) := \{a^i : i = 1, \dots, S(k)\}$ .
- $P(a)$  cannot be partitioned into  $k$  product-free sets.
- Fix  $A' = [\frac{1}{2}\varepsilon n^{1/S(k)}, n^{1/S(k)}]$ .
- For  $a, b \in A'$  distinct,  $P(a) \cap P(b) = \emptyset$  and  $P(a), P(b) \subseteq [n]$ .
- Any subset of  $[n]$  that can be partitioned into  $k$  product-free sets must avoid an element of  $P(a)$  for each  $a$  in  $A'$ .

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Lower bound next.



# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Lower bound next.

- Let  $\chi : [S'(k) - 1] \rightarrow [k]$  without monochromatic  $a + b = c$  or  $a + b = c - 1$ .

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Lower bound next.

- Let  $\chi : [S'(k) - 1] \rightarrow [k]$  without monochromatic  $a + b = c$  or  $a + b = c - 1$ .
- Colour  $a \in (n^{1/S'(k)}, n]$  with colour  $\chi(\lceil S'(k) \cdot \log_n(a) \rceil - 1)$ .

# Deterministic Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$  and  $k \in \mathbb{N}^+$ . For  $n$  large enough,

$$n - n^{1/S'(k)} \leq g_*(k, n) \leq n - (1 - \varepsilon)n^{1/S(k)}.$$

*Proof.* Lower bound next.

- Let  $\chi : [S'(k) - 1] \rightarrow [k]$  without monochromatic  $a + b = c$  or  $a + b = c - 1$ .
- Colour  $a \in (n^{1/S'(k)}, n]$  with colour  $\chi(\lceil S'(k) \cdot \log_n(a) \rceil - 1)$ .
- If  $ab = c$ , then let  $a' = \lceil S'(k) \cdot \log_n(a) \rceil - 1$ ,  $b' = \lceil S'(k) \cdot \log_n(b) \rceil - 1$ , and  $c' = \lceil S'(k) \cdot \log_n(c) \rceil - 1$  and note that  $\log_n(a) + \log_n(b) = \log_n(c)$  implies  $a' + b' = c'$  or  $a' + b' = c' - 1$ .

# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.

# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .

# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .



# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.

# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.

$b_2$     $r_2$

$b_1$     $r_1$


# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.




# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.




# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.




# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.



# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.
- To each  $\{r_1, r_2, b_1, b_2\}$  with  $r_1, r_2 \in R$ ,  $b_1, b_2 \in B$ ,  $r_1 r_2$  blue and  $b_1 b_2$  red we can associate a monochromatic product.

# Deterministic Product Schur II

Thm. (Mattos, MC, Parczyk, 2025)

Let  $\varepsilon > 0$ . For  $n$  large enough, any 2-colouring of  $[2, n]$  contains  $n^{1/3-\varepsilon}$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.
- To each  $\{r_1, r_2, b_1, b_2\}$  with  $r_1, r_2 \in R$ ,  $b_1, b_2 \in B$ ,  $r_1 r_2$  blue and  $b_1 b_2$  red we can associate a monochromatic product.
- Take care of double-counting.



# Deterministic Product Schur II

Thm. (Aragão, Chapman, Ortega, Souza, 2024+)

Any 2-colouring of  $[2, n]$  contains  $(\frac{1}{2\sqrt{2}} - o(1))n^{1/2} \log(n)$  monochromatic products.

*Proof.*

- **Lemma.** If  $A \subseteq [2, n]$  has size  $n - \frac{1}{2}\sqrt{n}$ , it contains  $n/8$  products.
- Let  $R$  be the red elements of  $[n^{1/3}]$ ,  $B$  the blue ones. By Lemma, wlog we have  $|R| \geq |B| \geq n^{1/6}/2$ .
- Let  $P_R := \{ab : a, b \in R\}$  and  $P_B := \{ab : a, b \in B\}$ .
- Note  $|P_R|, |P_B| \geq n^{1/3-\varepsilon}$ . Also,  $P_R$  contains  $n^{1/3-\varepsilon}$  blue elements and  $P_B$  contains  $n^{1/3-\varepsilon}$  red elements.
- To each  $\{r_1, r_2, b_1, b_2\}$  with  $r_1, r_2 \in R$ ,  $b_1, b_2 \in B$ ,  $r_1 r_2$  blue and  $b_1 b_2$  red we can associate a monochromatic product.
- Take care of double-counting.

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Lower bound first.

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Lower bound first.

- The number of products is

$$\sum_{a=2}^{\sqrt{n}} \frac{n}{a} + O(n) = (1 + o(1))n \log(n).$$

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Lower bound first.

- The number of products is

$$\sum_{a=2}^{\sqrt{n}} \frac{n}{a} + O(n) = (1 + o(1))n \log(n).$$

- The expected number of products in  $[2, n]_p$  is:

$$p^3(1 + o(1))n \log(n) + p^2\sqrt{n}.$$

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Lower bound first.

- The number of products is

$$\sum_{a=2}^{\sqrt{n}} \frac{n}{a} + O(n) = (1 + o(1))n \log(n).$$

- The expected number of products in  $[2, n]_p$  is:

$$p^3(1 + o(1))n \log(n) + p^2\sqrt{n}.$$

- For  $p \ll (n \log(n))^{-1/3}$ , there are no products.

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Upper bound next (very sketched).

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Upper bound next (very sketched).

- Take  $p = (n \log(n))^{-1/3} \omega(n)$ , and let  $q$  be such that  $(1 - q)^2 = (1 - p)$ .



# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Upper bound next (very sketched).

- Take  $p = (n \log(n))^{-1/3} \omega(n)$ , and let  $q$  be such that  $(1 - q)^2 = (1 - p)$ .
- Let  $A, B := [2, n]_q$ , and  $C = A \cup B$ . Note  $C \sim [2, n]_p$ .

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Upper bound next (very sketched).

- Take  $p = (n \log(n))^{-1/3} \omega(n)$ , and let  $q$  be such that  $(1 - q)^2 = (1 - p)$ .
- Let  $A, B := [2, n]_q$ , and  $C = A \cup B$ . Note  $C \sim [2, n]_p$ .
- This reduces to show  $|A^2 \cap [2, n]| \gg 1/q$  w.h.p.

# Random Product Schur

Thm. (Mattos, MC, Parczyk, 2025)

The threshold for  $[2, n]_p$  to contain a product Schur triple is  $(n \log(n))^{-1/3}$ .

*Proof.* Upper bound next (very sketched).

- Take  $p = (n \log(n))^{-1/3} \omega(n)$ , and let  $q$  be such that  $(1 - q)^2 = (1 - p)$ .
- Let  $A, B := [2, n]_q$ , and  $C = A \cup B$ . Note  $C \sim [2, n]_p$ .
- This reduces to show  $|A^2 \cap [2, n]| \gg 1/q$  w.h.p.
- A useful tool is that no  $c$  can be written as the product of elements of  $A$  in more than 2 ways w.h.p.

Open questions.

Open questions.

- Is there an  $\varepsilon$  such that  $S'(k) < (1 - \varepsilon)S(k)$  for large  $k$ ?

Open questions.

- Is there an  $\varepsilon$  such that  $S'(k) < (1 - \varepsilon)S(k)$  for large  $k$ ?
- What is the  $k$ -colouring of  $n$  with the fewest monochromatic products?

Open questions.

- Is there an  $\varepsilon$  such that  $S'(k) < (1 - \varepsilon)S(k)$  for large  $k$ ?
- What is the  $k$ -colouring of  $n$  with the fewest monochromatic products?
- What is the threshold for any  $k$ -colouring of  $[2, n]_p$  to contain a monochromatic product?

