



Chromatic profile of $\{C_3, \dots, C_{2k-1}\}$

J. Böttcher, N. Frankl, **D. Mergoni Cecchelli**, O. Parczyk, J. Skokan




Proper colouring and chromatic number




Introduction


Proper colouring and chromatic number



Proper colouring and chromatic number



Proper colouring and chromatic number



The **chromatic number** $\chi(G)$ is the number of colours needed to colour G .

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?


- $H = K_{1,k}$

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓ $\chi(G) \leq k$



Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?


- $H = K_{1,k}$ ✓
- $H = C^{\text{Odd}}$

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = C^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = C^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = C^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = C^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = C^{\text{Odd}}$




Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = C^{\text{Odd}}$ ✓ $\chi(G) \leq 2$



Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$ ✓
- $H = K_k$

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$ ✓
- $H = K_k$



Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$ ✓
- $H = K_3$ ✗

Thm. (Tutte, 1940's). Mycielski, Burling, ...

There are K_3 -free graphs of arbitrarily high chromatic number.

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$ ✓
- $H = K_3$ ✗
- $H = \mathcal{C}_{\leq 2k-1}^{\text{Odd}}$

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$ ✓
- $H = K_3$ ✗
- $H = \mathcal{C}_{\leq 2k-1}^{\text{Odd}}$ ✗

Thm. (Erdős, 1959). Kneser+Lovász, Alon et al., ...

There are graphs with arbitrarily high chromatic number and girth.

Bounding $\chi(G)$ in H -free graphs

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$ ✓
- $H = K_3$ ✗
- $H = \mathcal{C}_{\leq 2k-1}^{\text{Odd}}$ ✗

Thm. (Erdős, 1959). Kneser+Lovász, Alon et al., ...

There are graphs with arbitrarily high chromatic number and girth.

One of the first applications of the Probabilistic method.

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Start point

Can we bound $\chi(G)$ if G avoids H ?

- $H = K_{1,k}$ ✓
- $H = \mathcal{C}^{\text{Odd}}$ ✓
- $H = K_3$ ✗
- $H = \mathcal{C}_{\leq 2k-1}^{\text{Odd}}$ ✗

Prbl. (Erdős, Simonovits, 1973; insp. Hajnal)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973; insp. Hajnal)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Conj. (Erdős, Simonovits, 1973)

If G is K_3 -free and $\delta(G) > \frac{1}{3}|G|$, then $\chi(G) \leq 3$.


Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973; insp. Hajnal)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Conj. (Erdős, Simonovits, 1973), **Thm.** (Häggkvist, 1982)

If G is K_3 -free and $\delta(G) > \frac{1}{3}|G|$, then $\chi(G) \leq 3$.




Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973; insp. Hajnal)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Conj. (Erdős, Simonovits, 1973), Thm. (Häggkvist, 1982)

If G is K_3 -free and $\delta(G) > \frac{1}{3}|G|$, then $\chi(G) \leq 3$.




Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973; insp. Hajnal)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Conj. (Erdős, Simonovits, 1973), Thm. (Häggkvist, 1982)

If G is K_3 -free and $\delta(G) > \frac{1}{3}|G|$, then $\chi(G) \leq 3$.




Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973; insp. Hajnal)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Conj. (Erdős, Simonovits, 1973), Thm. (Häggkvist, 1982)

If G is K_3 -free and $\delta(G) > \frac{1}{3} \stackrel{?}{\frac{10}{29}} |G|$, then $\chi(G) \leq 3$.




Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973; insp. Hajnal)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Conj. (Erdős, Simonovits, 1973), Thm. (Häggkvist, 1982)

If G is K_3 -free and $\delta(G) > \frac{3}{8}|G|$, then $\chi(G) \leq 3$.



Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Thm. (Andrásfai, 1964)

- If G is K_3 -free and $\delta(G) > \frac{2}{5}|G|$, then $\chi(G) \leq 2$;

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Thm. (Häggkvist, 1982)

- If G is K_3 -free and $\delta(G) > \frac{2}{5}|G|$, then $\chi(G) \leq 2$;
- If G is K_3 -free and $\delta(G) > \frac{3}{8} \overset{?}{\frac{10}{29}} |G|$, then $\chi(G) \leq 3$;

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Thm. (Jin, 1995)

- If G is K_3 -free and $\delta(G) > \frac{2}{5}|G|$, then $\chi(G) \leq 2$;
- If G is K_3 -free and $\delta(G) > \frac{10}{29}|G|$, then $\chi(G) \leq 3$;

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Thm. (Thomassen, 2002)

- If G is K_3 -free and $\delta(G) > \frac{2}{5}|G|$, then $\chi(G) \leq 2$;
- If G is K_3 -free and $\delta(G) > \frac{10}{29}|G|$, then $\chi(G) \leq 3$;
- If G is K_3 -free and $\delta(G) > (\frac{1}{3} + \varepsilon)|G|$ then $\chi(G) \leq C_\varepsilon$;

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Thm. (Brandt, Thomassé, 2005)

- If G is K_3 -free and $\delta(G) > \frac{2}{5}|G|$, then $\chi(G) \leq 2$;
- If G is K_3 -free and $\delta(G) > \frac{10}{29}|G|$, then $\chi(G) \leq 3$;
- If G is K_3 -free and $\delta(G) > \frac{1}{3}|G|$ then $\chi(G) \leq 4$;

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Thm. (Hajnal graphs)

- If G is K_3 -free and $\delta(G) > \frac{2}{5}|G|$, then $\chi(G) \leq 2$;
- If G is K_3 -free and $\delta(G) > \frac{10}{29}|G|$, then $\chi(G) \leq 3$;
- If G is K_3 -free and $\delta(G) > \frac{1}{3}|G|$ then $\chi(G) \leq 4$;
- $\forall k, \varepsilon > 0$, \exists a K_3 -free G with: $\chi(G) \geq k$ and $\delta(G) \geq (\frac{1}{3} - \varepsilon)|G|$.

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Prbl. (Erdős, Simonovits, 1973)

What can we say if G is K_3 -free and $\delta(G) \geq \alpha|G|$?

Summary (VV.AA., '70s - 2005)


Let G be a K_3 -free graph on n vertices.

$\delta(G) >$	$\frac{2}{5}n$	$\frac{10}{29}n$	$\frac{1}{3}n$	$(\frac{1}{3} - \varepsilon)n$
$\chi(G) \leq$	2	3	4	∞

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Summary (VV.AA., '70s - 2005)


Let G be a K_3 -free graph on n vertices.

$\delta(G) >$	$\frac{1}{2}n$	$\frac{2}{5}n$	$\frac{10}{29}n$	$\frac{1}{3}n$	$(\frac{1}{3} - \varepsilon)n$
$\chi(G) \leq$		2	3	4	∞

Bounding $\chi(G)$ in H -free graphs with $\delta(G) > \alpha|G|$

Summary (VV.AA., '70s - 2005)

Let G be a K_3 -free graph on n vertices.

$\delta(G) >$	$\frac{1}{2}n$	$\frac{2}{5}n$	$\frac{10}{29}n$	$\frac{1}{3}n$	$(\frac{1}{3} - \varepsilon)n$
$\chi(G) \leq$		2	3	4	∞

Thm. (Allen, Böttcher, Griffiths, Kohayakawa, Morris, 2013)

If G is H -free and $\delta(G) > (f(H) + \varepsilon)|G|$, then $\chi(G) \leq C_{\varepsilon, H}$ (optimal)

Our result

Conj. (Letzter, Snyder, '19; Ebsen, Schacht, '20)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Our result

For k large enough (≥ 600)

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Idea of the proof

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Idea of the proof

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

Idea of the proof

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

2) This can be shown in an auxiliary graph; (Thomassen, 2007)

Idea of the proof

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

2) This can be shown in an auxiliary graph; (Thomassen, 2007)

3) Work in the auxiliary graph.

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

2) This property can be shown in an auxiliary (edge-weighted) graph;

3) Work in the auxiliary graph.




If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

2) This property can be shown in an auxiliary (edge-weighted) graph;

3) Work in the auxiliary graph.




If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

2) This property can be shown in an auxiliary (edge-weighted) graph;

3) Work in the auxiliary graph.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

2) This property can be shown in an auxiliary (edge-weighted) graph;

3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Sufficient condition for $\chi(G) \leq 3$;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.


- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.



If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v$, $G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) This property can be shown in an auxiliary (edge-weighted) graph;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Work in the auxiliary graph.




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Work in the auxiliary graph.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Work in the auxiliary graph.

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Work in the auxiliary graph.
 - a) $|H| \leq 2k - 2$;

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Work in the auxiliary graph.
 - a) $|H| \leq 2k - 2$;
 - b) $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$;

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Work in the auxiliary graph.
 - a) $|H| \leq 2k - 2$;
 - b) $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$;
 - c) Spanning 3-tree T .

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Work in the auxiliary graph.
 - a) $|H| \leq 2k - 2$;
 - b) $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$;
 - c) Spanning 3-tree T .

Want: Good partition of $V(H)$.

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $H[V \setminus A]$ weigh. bipartite.

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $H[V \setminus A]$ weigh. bipartite.

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $H[V \setminus A]$ weigh. bipartite.

○ t

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $H[V \setminus A]$ weigh. bipartite.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- **Connected**;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $H[V \setminus A]$ weigh. bipartite.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $H[V \setminus A]$ weigh. bipartite.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $H[V \setminus A]$ weigh. bipartite.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)


If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case A: Two odd cycles C and D do not intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case A: Two odd cycles C and D do not intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case B : Any two odd cycles intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case B : Any two odd cycles intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case B : Any two odd cycles intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case B : Any two odd cycles intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case B : Any two odd cycles intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case B : Any two odd cycles intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let H be:

- $\{3\}$ -weighted;
- $|H| \leq 2k - 2$;
- $\text{girth}_w^{\text{Odd}}(H) \geq 2k - 1$.

Want $A \subseteq V(H)$:

- Connected;
- $H[A \cup \{v\}]$ weigh. bipartite;
- $|A| \geq \frac{4k}{3}$.

Case B : Any two odd cycles intersect.



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.


Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.


- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;



If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.


- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;



If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;




Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;

Open problems:

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;

Open problems:

- What happens for small k ?

Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;

Open problems:

- What happens for small k ?
- Lower bounds? Explicit constructions?

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;

Open problems:

- What happens for small k ?
- Lower bounds? Explicit constructions?
- What about C_{2k-1} -free graphs?

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;

Open problems:

- What happens for small k ?
- Lower bounds? Explicit constructions?
- What about C_{2k-1} -free graphs?



Thm. (Böttcher, Frankl, M., Parczyk, Skokan, '23)

If G is $\{C_3, C_5, \dots, C_{2k-1}\}$ -free and $\delta(G) \geq \frac{1}{2k-1}|G|$, then $\chi(G) \leq 3$.

Let G be a $\{C_3, C_5, \dots, C_{2k-1}\}$ -free graph with $\delta(G) \geq \frac{1}{2k-1}|G|$.

- 1) Find a connected A s.t. $\forall v, G[A \cup \{v\}]$ and $G[V \setminus A]$ are bipartite;
- 2) Consider the weighted $H = (\{v_1, \dots, v_h\}, \{v_i v_j : d(v_i, v_j) \leq 5\})$;
- 3) Find a ball of size $\geq \frac{4}{3}k$ in H ;

Open problems:

- What happens for small k ?
- Lower bounds? Explicit constructions?
- What about C_{2k-1} -free graphs?

