

Numbers in Non-Integer Bases

Connor Baker
Co-Work with Dr. Tyler White

Northern Virginia Community College

VMATYC, Spring 2017

1 Whole Numbers

- Definitions
- Examples
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- Examples
- Properties of Non-Integer Bases
- Language Generated by Non-Integer Bases

3 Questions and Answers

Overview

1 Whole Numbers

- Definitions
- Examples
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- Examples
- Properties of Non-Integer Bases
- Language Generated by Non-Integer Bases

3 Questions and Answers

Base

A base (or radix) is a mathematical building block used to describe a number system whose digits are used to represent numbers.

Definition

Base

A base (or radix) is a mathematical building block used to describe a number system whose digits are used to represent numbers.

Example (Different Whole Number Bases)

Definition

Base

A base (or radix) is a mathematical building block used to describe a number system whose digits are used to represent numbers.

Example (Different Whole Number Bases)

125_{10}

Definition

Base

A base (or radix) is a mathematical building block used to describe a number system whose digits are used to represent numbers.

Example (Different Whole Number Bases)

125_{10}

888_9

Definition

Base

A base (or radix) is a mathematical building block used to describe a number system whose digits are used to represent numbers.

Example (Different Whole Number Bases)

125_{10}

888_9

48.5_8

Definition

Base

A base (or radix) is a mathematical building block used to describe a number system whose digits are used to represent numbers.

Example (Different Whole Number Bases)

125_{10}

888_9

$48.5_8 ?$

Maximum Allowed Numeral in Base

The radix is the number of numerals allowed in a base, and it includes zero.

Maximum Allowed Numeral in Base

The radix is the number of numerals allowed in a base, and it includes zero. The maximum numeral in a base is one less than the base.

Definition

Maximum Allowed Numeral in Base

The radix is the number of numerals allowed in a base, and it includes zero. The maximum numeral in a base is one less than the base.

Example (Allowed Numbers in Different Bases)

Definition

Maximum Allowed Numeral in Base

The radix is the number of numerals allowed in a base, and it includes zero. The maximum numeral in a base is one less than the base.

Example (Allowed Numbers in Different Bases)

125_{10}

Definition

Maximum Allowed Numeral in Base

The radix is the number of numerals allowed in a base, and it includes zero. The maximum numeral in a base is one less than the base.

Example (Allowed Numbers in Different Bases)

125_{10}

888_9

Definition

Maximum Allowed Numeral in Base

The radix is the number of numerals allowed in a base, and it includes zero. The maximum numeral in a base is one less than the base.

Example (Allowed Numbers in Different Bases)

125_{10}

888_9

48.5_8

Definition

Maximum Allowed Numeral in Base

The radix is the number of numerals allowed in a base, and it includes zero. The maximum numeral in a base is one less than the base.

Example (Allowed Numbers in Different Bases)

125_{10}

888_9

48.5_9

Radix Point

A point used to separate the integer part of a number from the fractional part.

Definition

Radix Point

A point used to separate the integer part of a number from the fractional part.

Example (Radix Point Expressions)

Definition

Radix Point

A point used to separate the integer part of a number from the fractional part.

Example (Radix Point Expressions)

10.5_{10}

Definition

Radix Point

A point used to separate the integer part of a number from the fractional part.

Example (Radix Point Expressions)

10.5_{10}

$A5.E_{16}$

Definition

Radix Point

A point used to separate the integer part of a number from the fractional part.

Example (Radix Point Expressions)

10.5_{10}

$A5.E_{16}$

1.1_2

Beta Expansion

A means of rewriting a number as the digits multiplied by the base which is raised to the power of position of the digit.

Definition

Beta Expansion

A means of rewriting a number as the digits multiplied by the base which is raised to the power of position of the digit.

Example (Example of Beta Expansions)

Definition

Beta Expansion

A means of rewriting a number as the digits multiplied by the base which is raised to the power of position of the digit.

Example (Example of Beta Expansions)

$$125_{10} = 1 \times 10^2 + 2 \times 10^1 + 5 \times 10^0$$

Definition

Beta Expansion

A means of rewriting a number as the digits multiplied by the base which is raised to the power of position of the digit.

Example (Example of Beta Expansions)

$$125_{10} = 1 \times 10^2 + 2 \times 10^1 + 5 \times 10^0$$

$$A5.E_{16} = 10 \times 16^1 + 5 \times 16^0 + 14 \times 16^{-1}$$

Definition

Beta Expansion

A means of rewriting a number as the digits multiplied by the base which is raised to the power of position of the digit.

Example (Example of Beta Expansions)

$$125_{10} = 1 \times 10^2 + 2 \times 10^1 + 5 \times 10^0$$

$$A5.E_{16} = 10 \times 16^1 + 5 \times 16^0 + 14 \times 16^{-1}$$

$$10_2 = 1 \times 2^1 + 0 \times 2^0$$

Overview

1 Whole Numbers

- Definitions
- **Examples**
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- Examples
- Properties of Non-Integer Bases
- Language Generated by Non-Integer Bases

3 Questions and Answers

Let's Play a Game

- Suppose that you're given the interval $[0, 1]$, with a point somewhere on it

Let's Play a Game

- Suppose that you're given the interval $[0, 1]$, with a point somewhere on it



Let's Play a Game

- Suppose that you're given the interval $[0, 1]$, with a point somewhere on it



- You're allowed one operation
 - Multiplication by a whole number, followed by modulo one

Let's Play a Game

- Suppose that you're given the interval $[0, 1]$, with a point somewhere on it



- You're allowed one operation
 - Multiplication by a whole number, followed by modulo one
 - Once you choose a multiplier, you must use it for the duration

Let's Play a Game

- Suppose that you're given the interval $[0, 1]$, with a point somewhere on it



- You're allowed one operation
 - Multiplication by a whole number, followed by modulo one
 - Once you choose a multiplier, you must use it for the duration
- We keep track of the interval that it falls in after the operation

Dealing with Radix Point Expressions (Geometrically)


Example (Suppose you have a point...)



original

Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



How Partitioning Works

- We partition before we multiply

How Partitioning Works


- We partition before we multiply
- We multiply so that the first partition becomes the unit interval

How Partitioning Works

- We partition before we multiply
- We multiply so that the first partition becomes the unit interval
 - The number we multiply by is the base that we are converting to

How Partitioning Works

- We partition before we multiply
- We multiply so that the first partition becomes the unit interval
 - The number we multiply by is the base that we are converting to
- So for binary, this is what we were doing:



What this Tells Us

- Where the point was originally

What this Tells Us

- Where the point was originally
 - The conversion of 0.25_{10} to a number in binary: 0.01_2
 - We can check this with a beta expansion:
$$0.01_2 = 0 \times 2^0 + 0 \times 2^{-1} + 1 \times 2^{-2} = 1/4 = 0.25_{10}$$

What this Tells Us

- Where the point was originally
 - The conversion of 0.25_{10} to a number in binary: 0.01_2
 - We can check this with a beta expansion:
$$0.01_2 = 0 \times 2^0 + 0 \times 2^{-1} + 1 \times 2^{-2} = 1/4 = 0.25_{10}$$
- That there is a finite radix point expression in a given base

What this Tells Us

- Where the point was originally
 - The conversion of 0.25_{10} to a number in binary: 0.01_2
 - We can check this with a beta expansion:
$$0.01_2 = 0 \times 2^0 + 0 \times 2^{-1} + 1 \times 2^{-2} = 1/4 = 0.25_{10}$$
- That there is a finite radix point expression in a given base
- That our choice of partitioning and multiplication matter... this is what happens geometrically when we change bases

What this Tells Us

- Where the point was originally
 - The conversion of 0.25_{10} to a number in binary: 0.01_2
 - We can check this with a beta expansion:
$$0.01_2 = 0 \times 2^0 + 0 \times 2^{-1} + 1 \times 2^{-2} = 1/4 = 0.25_{10}$$
- That there is a finite radix point expression in a given base
- That our choice of partitioning and multiplication matter... this is what happens geometrically when we change bases
- Let's do another example

Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



original


Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



What this Tells Us

- Getting stuck in this cycle means this number is not representable as a finite radix point expression in base 3

What this Tells Us

- Getting stuck in this cycle means this number is not representable as a finite radix point expression in base 3
- In general, this can and does happen in any base

What this Tells Us

- Getting stuck in this cycle means this number is not representable as a finite radix point expression in base 3
- In general, this can and does happen in any base
- We can explain this phenomenon arithmetically

Theorem

Finite Radix Point Expression

Theorem

Finite Radix Point Expression

A number p is said to have a finite radix point expression if

$$p = \frac{n}{m_1 m_2 \dots m_k}$$

where $n \in \mathbb{N}$, and $m_1 m_2 \dots m_k$ are all the prime factors of the base raised to some power.

<http://mathworld.wolfram.com/DecimalExpansion.html>

Theorem

Finite Radix Point Expression

A number p is said to have a finite radix point expression if

$$p = \frac{n}{m_1 m_2 \dots m_k}$$

where $n \in \mathbb{N}$, and $m_1 m_2 \dots m_k$ are all the prime factors of the base raised to some power.

<http://mathworld.wolfram.com/DecimalExpansion.html>

Example (1/3 in Base 10)

$$\frac{1}{3} \neq \frac{n}{2^\alpha 5^\beta}$$

where $n, \alpha, \beta \in \mathbb{N}$.

Dealing with Radix Point Expressions (Arithmetically)

- We can do this operation with arithmetic

Dealing with Radix Point Expressions (Arithmetically)

- We can do this operation with arithmetic
 - We must first qualify what we were doing in our “game” previously

Beta Map

The transformation $T : [0, 1) \rightarrow [0, 1)$ defined by $T = (x \cdot \beta) \pmod{1}$.

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

$$0.1 * 2 = 0.2$$

$$0.2 \bmod (1) = 0.2$$

0.0

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

$0.1 * 2 = 0.2$	$0.2 \bmod (1) = 0.2$	0.0
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$	0.00

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

$0.1 * 2 = 0.2$	$0.2 \bmod (1) = 0.2$	0.0
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$	0.00
$0.4 * 2 = 0.8$	$0.8 \bmod (1) = 0.8$	0.000

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

$0.1 * 2 = 0.2$	$0.2 \bmod (1) = 0.2$	0.0
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$	0.00
$0.4 * 2 = 0.8$	$0.8 \bmod (1) = 0.8$	0.000
$0.8 * 2 = 1.6$	$1.6 \bmod (1) = 0.6$	0.0001

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

$0.1 * 2 = 0.2$	$0.2 \bmod (1) = 0.2$	0.0
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$	0.00
$0.4 * 2 = 0.8$	$0.8 \bmod (1) = 0.8$	0.000
$0.8 * 2 = 1.6$	$1.6 \bmod (1) = 0.6$	0.0001
$0.6 * 2 = 1.2$	$1.2 \bmod (1) = 0.2$	0.00011

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

$0.1 * 2 = 0.2$	$0.2 \bmod (1) = 0.2$	0.0
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$	0.00
$0.4 * 2 = 0.8$	$0.8 \bmod (1) = 0.8$	0.000
$0.8 * 2 = 1.6$	$1.6 \bmod (1) = 0.6$	0.0001
$0.6 * 2 = 1.2$	$1.2 \bmod (1) = 0.2$	0.00011
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$...

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert 0.1_{10} to binary.)

$0.1 * 2 = 0.2$	$0.2 \bmod (1) = 0.2$	0.0
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$	0.00
$0.4 * 2 = 0.8$	$0.8 \bmod (1) = 0.8$	0.000
$0.8 * 2 = 1.6$	$1.6 \bmod (1) = 0.6$	0.0001
$0.6 * 2 = 1.2$	$1.2 \bmod (1) = 0.2$	0.00011
$0.2 * 2 = 0.4$	$0.4 \bmod (1) = 0.4$...
		$0.0001\overline{11}_2$

1 Whole Numbers

- Definitions
- Examples
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- Examples
- Properties of Non-Integer Bases
- Language Generated by Non-Integer Bases

3 Questions and Answers

Definition

Word

A word is a sequence of characters from an alphabet.

Definition

Word

A word is a sequence of characters from an alphabet.

Example (Examples of Words)

Definition

Word

A word is a sequence of characters from an alphabet.

Example (Examples of Words)

101 (Binary)

Definition

Word

A word is a sequence of characters from an alphabet.

Example (Examples of Words)

101 (Binary)

99.9 (Decimal)

Definition

Language

A language is the set of all allowed words.

Definition

Language

A language is the set of all allowed words.

Example (Examples of Languages)

Definition

Language

A language is the set of all allowed words.

Example (Examples of Languages)

Binary

Definition

Language

A language is the set of all allowed words.

Example (Examples of Languages)

Binary

Ternary

Definition

Language

A language is the set of all allowed words.

Example (Examples of Languages)

Binary

Ternary

Numerical Bases

Language Generated by Whole Number Bases

- A word is not allowed in a language if it contains a forbidden subword

Language Generated by Whole Number Bases

- A word is not allowed in a language if it contains a forbidden subword
 - Acceptable words in base 10 include 9, or 5.5 – but not CAT or 5Z: they contain forbidden subwords

Language Generated by Whole Number Bases

- A word is not allowed in a language if it contains a forbidden subword
 - Acceptable words in base 10 include 9, or 5.5 – but not CAT or 5Z: they contain forbidden subwords
 - Acceptable words in base 2 include 1, or 0.110000 – but not 3, or 4: they contain forbidden subwords


Language Generated by Whole Number Bases

- A word is not allowed in a language if it contains a forbidden subword
 - Acceptable words in base 10 include 9, or 5.5 – but not CAT or 5Z: they contain forbidden subwords
 - Acceptable words in base 2 include 1, or 0.110000 – but not 3, or 4: they contain forbidden subwords
- How can we represent the language generated by these bases?


Finite State Machine

A finite state machine (FSM) is a mathematical structure that allows us to imagine a machine with no memory.


FSM for Binary




FSM for Ternary



Simplified FSM for Ternary



FSM for Whole Number Bases



- $n \in \mathbb{N}, n > 1$

Overview

1 Whole Numbers

- Definitions
- Examples
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- Examples
- Properties of Non-Integer Bases
- Language Generated by Non-Integer Bases

3 Questions and Answers

Maximum Allowed Numeral in Base

If the radix is a whole number, the previous definition holds.

If the radix is not a whole number, then the maximum allowed numeral is the floor function of the base.

Definition

Maximum Allowed Numeral in Base

If the radix is a whole number, the previous definition holds.

If the radix is not a whole number, then the maximum allowed numeral is the floor function of the base.

Example (Allowed Numbers in Different Bases)

Definition

Maximum Allowed Numeral in Base

If the radix is a whole number, the previous definition holds.

If the radix is not a whole number, then the maximum allowed numeral is the floor function of the base.

Example (Allowed Numbers in Different Bases)

$\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618$, maximum numeral is 1

Definition

Maximum Allowed Numeral in Base

If the radix is a whole number, the previous definition holds.

If the radix is not a whole number, then the maximum allowed numeral is the floor function of the base.

Example (Allowed Numbers in Different Bases)

$\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618$, maximum numeral is 1

$\alpha = 1 + \sqrt{2} \approx 2.14$, maximum numeral is 2

Definition

Algebraic Integer

A number x is an algebraic integer if it is the root of a monic polynomial $f(x) = x^n + a_{n-1}x^{n-1} + \cdots + a_1x + a_0$, where $a_{n-1}, \dots, a_0 \in \mathbb{Z}$.

Example (Examples of Algebraic Polynomials)

Definition

Algebraic Integer

A number x is an algebraic integer if it is the root of a monic polynomial $f(x) = x^n + a_{n-1}x^{n-1} + \cdots + a_1x + a_0$, where $a_{n-1}, \dots, a_0 \in \mathbb{Z}$.

Example (Examples of Algebraic Polynomials)

$$\varphi^2 - \varphi - 1 = 0, \quad x = \frac{1 \pm \sqrt{5}}{2}$$

Definition

Algebraic Integer

A number x is an algebraic integer if it is the root of a monic polynomial $f(x) = x^n + a_{n-1}x^{n-1} + \cdots + a_1x + a_0$, where $a_{n-1}, \dots, a_0 \in \mathbb{Z}$.

Example (Examples of Algebraic Polynomials)

$$\varphi^2 - \varphi - 1 = 0, \quad x = \frac{1 \pm \sqrt{5}}{2}$$

$$\alpha^2 - 2\alpha - 1 = 0, \quad x = 1 \pm \sqrt{2}$$

Definition

Algebraic Integer

A number x is an algebraic integer if it is the root of a monic polynomial $f(x) = x^n + a_{n-1}x^{n-1} + \cdots + a_1x + a_0$, where $a_{n-1}, \dots, a_0 \in \mathbb{Z}$.

Example (Examples of Algebraic Polynomials)

$$\varphi^2 - \varphi - 1 = 0, \quad x = \frac{1 \pm \sqrt{5}}{2}$$

$$\alpha^2 - 2\alpha - 1 = 0, \quad x = 1 \pm \sqrt{2}$$

$$x^3 - x - 1 = 0, \quad x = \frac{1}{3} \sqrt[3]{\frac{27 - 3\sqrt{69}}{2}} + \frac{\sqrt[3]{\frac{1}{2}(9 + \sqrt{69})}}{3^{2/3}}$$

Partitioning Differently


- We don't have to use a whole number

Partitioning Differently

- We don't have to use a whole number
- The partitions won't be even, but that's not a problem


Partitioning Differently

- We don't have to use a whole number
- The partitions won't be even, but that's not a problem



Partitioning Differently

- We don't have to use a whole number
- The partitions won't be even, but that's not a problem



Overview

1 Whole Numbers

- Definitions
- Examples
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- **Examples**
- Properties of Non-Integer Bases
- Language Generated by Non-Integer Bases

3 Questions and Answers

Examples of Non-Integer Bases

- Base Golden Ratio
- Base Silver Ratio


Golden Ratio

The positive solution to the equation $\varphi^2 = \varphi + 1$.

$$\varphi = \frac{1+\sqrt{5}}{2} \approx 1.618.$$

Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



original


Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



What this Tells Us

- Getting stuck in this cycle means this number is not representable as a finite radix point expression in base golden ratio

What this Tells Us

- Getting stuck in this cycle means this number is not representable as a finite radix point expression in base golden ratio
 - $0.5_{10} = 0.\overline{010}_\varphi$

What this Tells Us

- Getting stuck in this cycle means this number is not representable as a finite radix point expression in base golden ratio
 - $0.5_{10} = 0.\overline{010}_\varphi$
- In general, whole numbers (excluding the base and its multiples) are typically the only numbers with finite representations in non-integer, algebraic bases

What this Tells Us

- Getting stuck in this cycle means this number is not representable as a finite radix point expression in base golden ratio
 - $0.5_{10} = 0.\overline{010}_\varphi$
- In general, whole numbers (excluding the base and its multiples) are typically the only numbers with finite representations in non-integer, algebraic bases
- Now, an arithmetic example

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert $(\varphi/2)_{10}$ to base golden ratio.)

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert $(\varphi/2)_{10}$ to base golden ratio.)

$$\frac{\varphi}{2} \cdot \varphi = \frac{\varphi^2}{2} \quad \frac{\varphi^2}{2} \bmod (1) = \frac{\varphi^2}{2} - 1 \quad 0.1$$

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert $(\varphi/2)_{10}$ to base golden ratio.)

$$\frac{\varphi}{2} \cdot \varphi = \frac{\varphi^2}{2} \quad \frac{\varphi^2}{2} \bmod (1) = \frac{\varphi^2}{2} - 1 \quad 0.1$$

$$\left(\frac{\varphi^2}{2} - 1\right) \cdot \varphi = \frac{\varphi^3}{2} - \varphi \quad \frac{\varphi^3}{2} - \varphi \bmod (1) = \frac{\varphi^3}{2} - \varphi \quad 0.10$$

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert $(\varphi/2)_{10}$ to base golden ratio.)

$$\frac{\varphi}{2} \cdot \varphi = \frac{\varphi^2}{2} \quad \frac{\varphi^2}{2} \bmod (1) = \frac{\varphi^2}{2} - 1 \quad 0.1$$

$$\left(\frac{\varphi^2}{2} - 1\right) \cdot \varphi = \frac{\varphi^3}{2} - \varphi \quad \frac{\varphi^3}{2} - \varphi \bmod (1) = \frac{\varphi^3}{2} - \varphi \quad 0.10$$

$$\left(\frac{\varphi^3}{2} - \varphi\right) \cdot \varphi = \frac{\varphi^4}{2} - \varphi^2 \quad \frac{\varphi^4}{2} - \varphi^2 \bmod (1) = \frac{\varphi^4}{2} - \varphi^2 \quad 0.100$$

Dealing with Radix Point Expressions (Arithmetically)

Example (Convert $(\varphi/2)_{10}$ to base golden ratio.)

$$\frac{\varphi}{2} \cdot \varphi = \frac{\varphi^2}{2} \quad \frac{\varphi^2}{2} \bmod (1) = \frac{\varphi^2}{2} - 1 \quad 0.1$$

$$\left(\frac{\varphi^2}{2} - 1\right) \cdot \varphi = \frac{\varphi^3}{2} - \varphi \quad \frac{\varphi^3}{2} - \varphi \bmod (1) = \frac{\varphi^3}{2} - \varphi \quad 0.10$$

$$\left(\frac{\varphi^3}{2} - \varphi\right) \cdot \varphi = \frac{\varphi^4}{2} - \varphi^2 \quad \frac{\varphi^4}{2} - \varphi^2 \bmod (1) = \frac{\varphi^4}{2} - \varphi^2 \quad 0.100$$

$$\frac{\varphi^4}{2} - \varphi^2 = \frac{\varphi^2(\varphi^2 - 2)}{2} = \frac{(\varphi + 1)(\varphi - 1)}{2} = \frac{\varphi^2 - 1}{2}$$


$$0.\overline{100}_{\varphi}$$

Silver Ratio

The positive solution to the equation $\alpha^2 = 2\alpha + 1$.
 $\alpha = 1 + \sqrt{2} \approx 2.414$.

Dealing with Radix Point Expressions (Geometrically)


Example (Suppose you have a point...)



original


Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)




Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



Dealing with Radix Point Expressions (Geometrically)

Example (Suppose you have a point...)



Overview

1 Whole Numbers

- Definitions
- Examples
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- Examples
- **Properties of Non-Integer Bases**
- Language Generated by Non-Integer Bases

3 Questions and Answers

Properties of Non-Integer Bases

- There will be forbidden words

Properties of Non-Integer Bases

- There will be forbidden words
- It is easy/interesting to represent certain numbers in these bases

Properties of Non-Integer Bases

- There will be forbidden words
- It is easy/interesting to represent certain numbers in these bases
- Arithmetic is more of a pain than usual

Simplest Form

A number is in simplest form if we are unable to rewrite the beta expansion such that it includes a higher power.

Example of Simplest Form

Simplest Form

Let

$$\varphi^2 = \varphi + 1.$$

Example of Simplest Form

Simplest Form

Let

$$\varphi^2 = \varphi + 1.$$

Using the positive solution of that equation, $\varphi \approx 1.618$, we can show that $0.11_\varphi = 1.00_\varphi$ by using a beta expansion.

Example of Simplest Form

Simplest Form

Let

$$\varphi^2 = \varphi + 1.$$

Using the positive solution of that equation, $\varphi \approx 1.618$, we can show that $0.11_\varphi = 1.00_\varphi$ by using a beta expansion.

$$0.11_\varphi = 1 \times \varphi^{-1} + 1 \times \varphi^{-2} = \frac{1}{\varphi} + \frac{1}{\varphi^2}.$$

Example of Simplest Form

Simplest Form

Let

$$\varphi^2 = \varphi + 1.$$

Using the positive solution of that equation, $\varphi \approx 1.618$, we can show that $0.11_\varphi = 1.00_\varphi$ by using a beta expansion.

$$0.11_\varphi = 1 \times \varphi^{-1} + 1 \times \varphi^{-2} = \frac{1}{\varphi} + \frac{1}{\varphi^2}.$$

Then,

$$\frac{\varphi}{\varphi^2} + \frac{1}{\varphi^2} = \frac{\varphi + 1}{\varphi^2},$$

Example of Simplest Form

Simplest Form

Let

$$\varphi^2 = \varphi + 1.$$

Using the positive solution of that equation, $\varphi \approx 1.618$, we can show that $0.11_\varphi = 1.00_\varphi$ by using a beta expansion.

$$0.11_\varphi = 1 \times \varphi^{-1} + 1 \times \varphi^{-2} = \frac{1}{\varphi} + \frac{1}{\varphi^2}.$$

Then,

$$\frac{\varphi}{\varphi^2} + \frac{1}{\varphi^2} = \frac{\varphi + 1}{\varphi^2},$$

and by the definition of the base, we can rewrite the numerator:

$$\frac{\varphi^2}{\varphi^2} = 1_\varphi$$

Golden Ratio Addition

Example $(1_\varphi + 1_\varphi)$

Golden Ratio Addition

Example ($1_\varphi + 1_\varphi$)

$$1_\varphi = 0.11_\varphi.$$

Golden Ratio Addition

Example ($1_\varphi + 1_\varphi$)

$$1_\varphi = 0.11_\varphi.$$

$$1_\varphi + 0.11_\varphi = 1.11_\varphi = 10.01_\varphi.$$

Example ($1_\varphi + 1_\varphi$)

$$1_\varphi = 0.11_\varphi.$$

$$1_\varphi + 0.11_\varphi = 1.11_\varphi = 10.01_\varphi.$$

We can double check this operation by doing a beta expansion and using the definition of the base.

Example ($1_\varphi + 1_\varphi$)

$$1_\varphi = 0.11_\varphi.$$

$$1_\varphi + 0.11_\varphi = 1.11_\varphi = 10.01_\varphi.$$

We can double check this operation by doing a beta expansion and using the definition of the base.

$$10.01_\varphi = 1 \times \varphi^1 + 0 \times \varphi^0 + 0 \times \varphi^{-1} + 1 \times \varphi^{-2} = \varphi + 1/\varphi^2 = (\varphi^2 + 1)/\varphi^2 = 2_{10}.$$

Example ($1_\varphi + 1_\varphi$)

$$1_\varphi = 0.11_\varphi.$$

$$1_\varphi + 0.11_\varphi = 1.11_\varphi = 10.01_\varphi.$$

We can double check this operation by doing a beta expansion and using the definition of the base.

$$10.01_\varphi = 1 \times \varphi^1 + 0 \times \varphi^0 + 0 \times \varphi^{-1} + 1 \times \varphi^{-2} = \varphi + 1/\varphi^2 = (\varphi^2 + 1)/\varphi^2 = 2_{10}.$$

We can calculate any whole number through this method.

Overview

1 Whole Numbers

- Definitions
- Examples
- Language Generated by Whole Number Bases

2 Non-Integer Algebraic Numbers

- Definitions
- Examples
- Properties of Non-Integer Bases
- Language Generated by Non-Integer Bases

3 Questions and Answers

Example of Simplest Form

Simplest Form

Let

$$\varphi^2 = \varphi + 1.$$

Using the positive solution of that equation, $\varphi \approx 1.618$, we can show that $0.11_\varphi = 1.00_\varphi$ by using a beta expansion.

$$0.11_\varphi = 1 \times \varphi^{-1} + 1 \times \varphi^{-2} = \frac{1}{\varphi} + \frac{1}{\varphi^2}.$$

Then,

$$\frac{\varphi}{\varphi^2} + \frac{1}{\varphi^2} = \frac{\varphi + 1}{\varphi^2},$$

and by the definition of the base, we can rewrite the numerator:

$$\frac{\varphi^2}{\varphi^2} = 1_\varphi$$

What Makes a Word Forbidden in Non-Integer Bases?

- In general, a word is forbidden if it is not in simplest form

What Makes a Word Forbidden in Non-Integer Bases?

- In general, a word is forbidden if it is not in simplest form
- In actuality, it is because it escapes the range of the beta map

Escaping the Beta Map

Example (Golden Base Ratio: $T(0.01101_\varphi)$)

Using our previous example:

$$T(0.01101_\varphi) = 0.1101_\varphi = 1.0001_\varphi.$$

Escaping the Beta Map

Example (Golden Base Ratio: $T(0.01101_\varphi)$)

Using our previous example:

$$T(0.01101_\varphi) = 0.1101_\varphi = 1.0001_\varphi.$$

But that can't happen! By definition, $T : [0, 1) \rightarrow [0, 1)$. As such, we say that this is a forbidden word.

Escaping the Beta Map

Example (Golden Base Ratio: $T(0.01101_\varphi)$)

Using our previous example:

$$T(0.01101_\varphi) = 0.1101_\varphi = 1.0001_\varphi.$$

But that can't happen! By definition, $T : [0, 1) \rightarrow [0, 1)$. As such, we say that this is a forbidden word.

Geometrically, this can't happen. If we multiply a number in $[0, 1)$ by any number and perform (mod 1), we will never get a number larger than one. It's the arithmetic equivalent of having $x \cdot \beta > \beta$, $x < 1$.

Orbit

The unique, non-terminating radix point representation of a number in any base.

Definition

Orbit

The unique, non-terminating radix point representation of a number in any base.

Example (Orbit of One in Different Bases)

Definition

Orbit

The unique, non-terminating radix point representation of a number in any base.

Example (Orbit of One in Different Bases)

$$1_{10} = 0.\overline{99}_{10}$$

Definition

Orbit

The unique, non-terminating radix point representation of a number in any base.

Example (Orbit of One in Different Bases)

$$1_{10} = 0.\overline{99}_{10}$$

$$1_9 = 0.\overline{88}_9$$

Definition

Orbit

The unique, non-terminating radix point representation of a number in any base.

Example (Orbit of One in Different Bases)

$$1_{10} = 0.\overline{99}_{10}$$

$$1_9 = 0.\overline{88}_9$$

$$1_2 = 0.\overline{11}_2$$

Finding Forbidden Words

- Take the orbit of one in the base

Finding Forbidden Words

- Take the orbit of one in the base
- Find the period of the orbit and take one segment

Finding Forbidden Words

- Take the orbit of one in the base
- Find the period of the orbit and take one segment
 - Any sequence of numerals greater than the numerals of the period is not allowed

Finding Forbidden Words

- The orbit of one is composed of the maximum allowed numeral in a language

Finding Forbidden Words

- The orbit of one is composed of the maximum allowed numeral in a language
- All the forbidden words then will subwords of the orbit, with one added until the word is the maximum numeral possible

Language Generated by base Golden Ratio

- $0.\overline{10}_\varphi$ is the orbit of one in base golden ratio

Language Generated by base Golden Ratio

- $0.\overline{10}_\varphi$ is the orbit of one in base golden ratio
- Since the maximum allowed numeral in the language is 10, we add one and find that the forbidden word is 11


Language Generated by base Golden Ratio

- $0.\overline{10}_\varphi$ is the orbit of one in base golden ratio
- Since the maximum allowed numeral in the language is 10, we add one and find that the forbidden word is 11
- There can be more than one forbidden word, but since the maximum allowed numeral in base golden ratio is 1, we can't add one again

Language Generated by base Golden Ratio

- $0.\overline{10}_\varphi$ is the orbit of one in base golden ratio
- Since the maximum allowed numeral in the language is 10, we add one and find that the forbidden word is 11
- There can be more than one forbidden word, but since the maximum allowed numeral in base golden ratio is 1, we can't add one again
- With this in mind, we can now create a finite state machine

FSM For Language Generated by Base Golden Ratio




Language Generated by base Silver Ratio

- $0.\overline{20}_\alpha$ is the orbit of one in base silver ratio

Language Generated by base Silver Ratio

- $0.\overline{20}_\alpha$ is the orbit of one in base silver ratio
- Therefore, 21 and 22 are forbidden words in the language

FSM For Language Generated by Base Silver Ratio



- Positive solution of the monic polynomial $x^2 = 5x + 5$

Language Generated by Base $\frac{5+(3\sqrt{5})}{2}$

- Positive solution of the monic polynomial $x^2 = 5x + 5$
- Base is ≈ 5.85 , so it's equivalent to base 5


Language Generated by Base $\frac{5+(3\sqrt{5})}{2}$

- Positive solution of the monic polynomial $x^2 = 5x + 5$
- Base is ≈ 5.85 , so it's equivalent to base 5
- $0.\overline{54}_\beta$ is the orbit of one

Language Generated by Base $\frac{5+(3\sqrt{5})}{2}$

- Positive solution of the monic polynomial $x^2 = 5x + 5$
- Base is ≈ 5.85 , so it's equivalent to base 5
- $0.\overline{54}_\beta$ is the orbit of one
- Therefore, 55 is the forbidden word in the language

FSM For Language Generated by Base $\frac{5+(3\sqrt{5})}{2}$



Questions?