

## Homework 5

### Connor Baker, March 2017

1. Prove that if the real-valued function  $f$  is strictly increasing or strictly decreasing on  $\mathbb{R}$ , then  $f$  is one-to-one (Note: You cannot assume  $f$  is differentiable).

*Proof.* Case 1:  $f$  is strictly decreasing.

If  $f$  is strictly decreasing, then  $\forall x, a \in \text{dom}(f)$ ,  $x \leq a$ ,  $f(x) \geq f(a)$ . If it is the case that  $f(x) = f(a)$ , then it must also be the case that  $x = a$ . Then,  $f(x) = f(a)$  if and only if  $x = a$ , and  $f(x) > f(a) \forall x, a \in \text{dom}(f)$ ,  $x < a$ , the function  $f$  must be one-to-one.

Case 2:  $f$  is strictly increasing.

If  $f$  is strictly increasing, then  $\forall x, a \in \text{dom}(f)$ ,  $x \leq a$ ,  $f(x) \leq f(a)$ . If it is the case that  $f(x) = f(a)$ , then it must also be the case that  $x = a$ . Then,  $f(x) = f(a)$  if and only if  $x = a$ , and  $f(x) < f(a) \forall x, a \in \text{dom}(f)$ ,  $x < a$ , the function  $f$  must be one-to-one.  $\square$

2. Prove the following are metrics:

(a)  $X = \mathbb{R}, d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$

(b)  $X = \mathbb{R} \times \mathbb{R}, d((x, y), (z, w)) = \sqrt{(x - z)^2 + (y - w)^2}$

**Definition 1** (Metric). A metric on a set  $X$  is a function  $d : X \times X \rightarrow \mathbb{R}$  such that for all  $x, y, z \in X$ ,

(a)  $d(x, y) \geq 0$

(b)  $d(x, y) = 0$  if and only if  $x = y$

(c)  $d(x, y) = d(y, x)$ ,

(d)  $d(x, y) + d(y, z) \geq d(x, z)$ .

*Proof.* We begin by proving that the first function is a metric.

1. The  $\text{rang}(d) = \{0, 1\}$  so the function is definitely greater than zero.
2. By the definition of  $d$ , if and only if  $x = y$  does  $d(x, y) = 0$ .
3. Since the equals relationship is symmetric,  $x = y \implies y = x$ . As such,  $d(x, y) = d(y, x)$ .
4. Proving that  $d(x, y) + d(y, z) \geq d(x, z)$  must be done with five cases:
  - (a) Case 1:  $x = y = z$
  - (b) Case 2:  $x = y, y \neq z$
  - (c) Case 3:  $x \neq y, x = z$
  - (d) Case 4:  $x \neq y, y = z$
  - (e) Case 5:  $x \neq y, y \neq z$

□

3. Let  $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  be given by  $f(m, n) = 2^{m-1}(2n - 1)$ . Prove that  $f$  is one-to-one and onto.

*Proof.* Since

□

4. Let  $f : A \rightarrow B$  be a function from a nonempty set  $A$ . Prove that the set  $\mathcal{C} = \{f^{-1}(b) : b \in \text{rang}(f)\}$  is a partition of  $A$ . Note:  $\mathcal{C}$  is a subset of  $\mathcal{P}(A)$ .

*Proof.* Since

□

5. Let  $f : A \rightarrow B$  be a function from a nonempty set  $A$  which is surjective. Find a new function  $g : C \rightarrow B$  which is one-to-one such that  $C \subseteq A$ ,  $\text{rang}(g) = \text{rang}(f)$ , and for every  $x \in C$ ,  $f(x) = g(x)$ . Explain why  $g$  has an inverse function,  $g^{-1}$ . Then, compute  $f(g^{-1}(x))$  for all  $x \in B$ .

*Proof.* Since

□