

MTH 295: Homework 1

Connor Baker

January 2018

1. Prove by contradiction that if $a - b$ is odd, then $a + b$ is odd.

Proof. Assume that both a and b are integers. We proceed using proof by contradiction, and assume that $a - b$ is even implies that $a + b$ is odd.

Since $a - b$ is even, it can be written as

$$a - b = 2k \tag{1}$$

for some integer k , and since $a + b$ is odd, it can be written as

$$a + b = 2j + 1 \tag{2}$$

for some integer j . Solving both equations for a yields

$$a = 2k + b \tag{3a}$$

$$a = 2j + 1 - b \tag{3b}$$

and setting them equal to the other (by transitivity) gives:

$$2k + b = 2j + 1 - b. \tag{3c}$$

Simplifying by means of collecting b on the left hand side and factoring out a two gives

$$2(k + b) = 2j + 1. \tag{3d}$$

Since an even number can never be equal to an odd number (by definition), we have arrived at a contradiction, and our original assumption that $a - b$ is even implies that $a + b$ is odd must be incorrect.

Therefore, by means of proof by contradiction, if $a - b$ is odd, then $a + b$ is odd. \square

2. Write a proof by contrapositive to show that if xy is odd, then both x and y are odd.

Proof. We will prove that if x or y is even, then the product xy is even. Assume that x is even, and that $x, y \in \mathbb{Z}$. By definition, $x = 2k$ for all $k \in \mathbb{Z}$. Then $xy = 2ky$ is even. So, regardless of the parity of y , the product xy will be even so long as at least one is even. If x was odd, and y was even, then the above would still hold, due to multiplication being commutative.

Since either x or y is even, and xy is even, we can infer by the contrapositive that if xy is odd, then both x and y are odd. \square

3. Prove that there do not exist integers m and n such that $12m + 15n = 1$.

Proof. The equation $12m + 15n = 1$ is equivalent to $3(4m + 5n) = 1$. For this statement to be true, $4m + 5n$ must be the multiplicative inverse of 3, which is not in the set of natural numbers. Therefore, there do not exist integers m and n such that $12m + 15n = 1$. \square

4. Prove there is a natural number M such that for every natural number n , $\frac{1}{n} < M$.

Proof. Because $n \in \mathbb{N}$, $n \geq 1$ for all choices of n . As a result, $1/n \leq 1$, for all choices of n . Therefore, M can be any number such that $M \geq 1$. \square

5. Prove that if $-2 < x < 1$ or $x > 3$, then $\frac{(x-1)(x+2)}{(x-3)(x+4)} > 0$.

Proof. Let the function $f(x)$ be as follows:

$$f(x) = \frac{(x-1)(x+2)}{(x-3)(x+4)}$$

Then $f(x)$ has two x -intercepts at $x = -2, 1$, and two vertical asymptotes at $x = -4, 3$. By the Intermediate Value Theorem, those four x -values are the only places that the function can change the sign of its output. As such, it has been established that $f(x)$ does not change sign over the intervals $(-\infty, -4)$, $(-4, -2)$, $(-2, 1)$, $(1, 3)$, $(3, \infty)$.

By picking a point on the intervals $(-2, 1)$ and $(3, \infty)$ and verifying the sign, then by the Intermediate Value Theorem proves, the function value has the same sign on the entirety of the interval.

Let $x = 0$. Then, on the interval $(-2, 1)$, the function is positive.

Let $x = 4$. Then, on the interval $(3, \infty)$, the function is positive.

Therefore, by the Intermediate Value Theorem, if $-2 < x < 1$ or $x > 3$, then $\frac{(x-1)(x+2)}{(x-3)(x+4)} > 0$. □