


Rectangular domain



Plane strain
no f.

Try: $\phi(x, y) = \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} a_{mn} x^n y^m$

Note: 1.- Lower order terms ($m+n \leq 3$) satisfy biharmonic automatically


2.- Terms $a_{00} + a_{01}x + a_{10}y$ do not produce stress.

e.g. (a) $\phi = Axy \rightarrow \sigma_{xx} = \sigma_{yy} = 0$
 $\sigma_{xy} = -A$ (pure shear)

(b) $\phi = Ax^3 \rightarrow \sigma_{xx} = 6Axy$

$\sigma_{yy} = 0$

$\sigma_{xy} = -3Ay^2$



]

• Use S.O.M. and symmetry to guess m, n .

• From symmetry here: $\sigma_x = \frac{\partial^2 \phi}{\partial y^2}$ must be even in x

$$\sigma_y = \frac{\partial^2 \phi}{\partial x^2} \quad " \quad " \quad " \quad " \quad x$$

$$\sigma_{xy} = -\frac{\partial^2 \phi}{\partial x \partial y} \quad " \quad " \quad \text{odd} \quad " \quad x$$

try: $\phi(x, y) = ax^2 + bx^2y + cy^3 + dx^4 + eyx^4$
 $+ fx^2y^3 + gy^5 + \dots$

Biharmonic:

$$\nabla^4 \phi = 0 \Rightarrow 24d + 24ey + 24fy + 120gy = 0$$

Must hold $\forall (x, y) \Rightarrow \boxed{d=0}$
 and

$$e + f + 5g = 0$$

$$\Rightarrow \boxed{f = -e - 5g}$$

$$\therefore \phi(x, y) = ax^2 + bx^2y + cy^3 + e(x^4y - x^2y^3) + g(y^5 - 5x^2y^3)$$

Stresses become:

$$\sigma_x = 6cy - 6ex^2y + g(20y^3 - 30x^2y)$$

$$\sigma_y = 2a + 2by + e(12yx^2 - 2y^3) + g(-10y^3)$$

$$-\sigma_{xy} = 2bx + e(4x^3 - 6xy^2) + g(-30xy^2)$$

Boundary conditions:

On surface ③: $y = \frac{h}{2}$

$$\sigma_y(x, \frac{h}{2}) = -P_0 = 2a + 2b\frac{h}{2} + e(12\frac{h}{2}x^2 - 2\frac{h^3}{8}) - 10g\frac{h^3}{8}$$

$$\sigma_{xy}(x, \frac{h}{2}) = 0 = 2bx + e[4x^3 - 6x(\frac{h}{2})^2] - \frac{15}{2}h^2gx$$

wt hold $\forall x \Rightarrow \boxed{e=0}$

$$\therefore \left. \begin{aligned} -P_0 &= 2a + bh - \frac{10}{8}gh^3 \\ 0 &= 2b - \frac{15}{2}h^2g \end{aligned} \right\} \text{--- (1)}$$

On surface ④: $y = -\frac{h}{2}$

$$\sigma_y(\neq 0, -\frac{h}{2}) = 0 = 2a - bh + 10g \left(\frac{h}{2}\right)^3$$

$$\sigma_{xy}(x, -\frac{h}{2}) = 0 = \sigma(x, +\frac{h}{2}) \quad \text{same as above}$$

$$\Rightarrow 2a - bh + \frac{10}{8} h^3 g = 0 \quad \text{--- (2)}$$

$$(1), (2) \Rightarrow \boxed{g = -\frac{P_0}{5h^3}}$$

$$\boxed{a = -\frac{1}{4} P_0}$$

$$\boxed{b = -\frac{3}{4} \frac{P_0}{h}}$$

check dimensions
to be sure.

Unfortunately we can only satisfy the b.c. on
①, ② faces on average

On surface ①

$$\sigma_x \left(\frac{l}{2}, y \right) = 6 c y + g \left(20 y^3 - \frac{15}{2} l^2 y \right)$$

$$-\sigma_{xy} \left(\frac{l}{2}, y \right) = 6l + g(-15ly^2)$$

Use S.V. for justification of average b.c.

To satisfy "pinned" condition require no moment on end faces:



$$\int_{-h/2}^{h/2} y \sigma_x dy = 0$$

$$\Rightarrow \boxed{c = \frac{P_0}{10h} \left[1 - \frac{5}{2} \left(\frac{l}{h} \right)^2 \right]}$$

∴ Solution becomes:

$$\frac{\sigma_x}{P_0} = - \underbrace{\frac{3}{2} \left\{ \left(\frac{l}{h} \right)^2 - 4 \left(\frac{x}{l} \right)^2 \right\}}_{\text{S.O.M. solution}} \frac{y}{l} + \left\{ \frac{3}{5} - 4 \left(\frac{y}{h} \right)^2 \right\} \frac{y}{l}$$

$$\frac{\sigma_{xy}}{P_0} = \underbrace{\frac{3}{2} \left\{ 1 - 4 \left(\frac{y}{h} \right)^2 \right\} \frac{x}{h}}_{\text{S.O.M. solution}}$$

$$\frac{\sigma_y}{P_0} = -\frac{1}{2} - \frac{3}{2} \frac{y}{h} + 2 \left(\frac{y}{h} \right)^3$$

Note: Reaction forces on ① and ② come in through a net shear force S^*

$$S^* = \int_{-h/2}^{h/2} \sigma_{xy} dy$$

$$\Rightarrow S^* = \frac{3P_0}{2} \int_{-h/2}^{h/2} \left\{ 1 - 4 \left(\frac{y}{h} \right)^2 \right\} \cdot \frac{1}{2h} dy$$

$$\Rightarrow \boxed{S^* = \frac{1}{2} P_0 l}$$