

Functional Analysis — Homework 1

Connor Mooneyhan

Problem 1. For $p \in [1, \infty)$, give an example of a sequence which is in ℓ^q for all $q > p$, but not in ℓ^p .

Problem 2. If a subspace S of a normed space V contains a nonempty open set, then $S = V$.

Problem 3. Show that in any normed space V , the operations

$$+ : V \times V \rightarrow V \quad \cdot : \mathbb{R} \times V \rightarrow V$$

of addition and scalar multiplication are continuous.

Problem 4. Let V be a normed space. Show that a linear map $f : V \rightarrow \mathbb{R}$ is continuous if and only if $\ker(f)$ is closed.

Problem 5. Let $\rho : V \rightarrow [0, +\infty)$ be a seminorm. Prove that if there exists a positive number $\delta > 0$ such that the preimage $\rho^{-1}[0, \delta]$ contains an open ball, then ρ is continuous.

Problem 6. Let V be a normed space. Show that V is Banach if and only if every absolutely convergent series is convergent.

Problem 7. In this problem, you will show that for any number $p > 1$, the function

$$\|x\|_p = \|(x_1, \dots, x_n)\|_p = (|x_1|^p + \dots + |x_n|^p)^{1/p}$$

is a norm on $\mathbb{R}^n$. The extension to ℓ^p follows easily from that. It will be helpful to define a number q , called the conjugate exponent to p , by the equation: $\frac{1}{p} + \frac{1}{q} = 1$.

- (a) (Young's inequality) Use the fact that $\ln(x)$ is a concave function to prove that if a, b are positive numbers, then

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}.$$

- (b) (Hölder's inequality) Given any $x, y \in \mathbb{R}^n$, use Young's inequality to prove that

$$|\langle x, y \rangle| \leq \|x\|_p \|y\|_q$$

where $\langle x, y \rangle$ denotes the standard dot product in $\mathbb{R}^n$.

- (c) (Minkowski's inequality) Given any $x, y \in \mathbb{R}^n$, use Hölder's inequality to prove

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p,$$

which is the triangle inequality for $\|\cdot\|_p$.