

Functional Analysis — Homework 6

Connor Mooneyhan

Problem 1. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be an analytic function, and let

$$f(x, y) = u(x, y) + iv(x, y)$$

be its decomposition into real and imaginary parts. Consider the limit definition of the complex derivative at a point $z = x + iy$:

$$f'(z) = \lim_{w \rightarrow 0} \frac{f(z + w) - f(z)}{w}.$$

- (a) Evaluate this limit by letting $w \rightarrow 0$ strictly along the real axis. Express the result in terms of the partial derivatives of u and v .
- (b) Now evaluate the same limit by letting $w \rightarrow 0$ strictly along the imaginary axis.

Since f is analytic, the limit must be independent of the path. Equate your results from (a) and (b) to prove the Cauchy-Riemann equations:

$$u_x = v_y, \quad u_y = -v_x.$$

Proof.

- (a) Let $z = (x, y)$ and $w = (w_x, w_y)$, where in this case $w_y = 0$ because we are taking the limit along the real line. Then

$$\begin{aligned} f'(z) &= \lim_{w \rightarrow 0} \frac{f((x, y) + (w_x, w_y)) - f(x, y)}{w_x + iw_y} \\ &= \lim_{w_x \rightarrow 0} \frac{f(x + w_x, y) - f(x, y)}{w_x} \\ &= \lim_{w_x \rightarrow 0} \frac{u(x + w_x, y) + iv(x + w_x, y) - u(x, y) - iv(x, y)}{w_x} \\ &= \lim_{w_x \rightarrow 0} \frac{u(x + w_x, y) - u(x, y)}{w_x} + i \lim_{w_x \rightarrow 0} \frac{v(x + w_x, y) - v(x, y)}{w_x} \\ &= u_x + iv_x. \end{aligned}$$

(b) Now if $w_x = 0 \neq w_y$, then

$$\begin{aligned}
f'(z) &= \lim_{w \rightarrow 0} \frac{f((x, y) + (w_x, w_y)) - f(x, y)}{iw_y} \\
&= \lim_{w_y \rightarrow 0} \frac{f(x, y + w_y) - f(x, y)}{iw_y} \\
&= \lim_{w_y \rightarrow 0} \frac{u(x, y + w_y) + iv(x, y + w_y) - u(x, y) - iv(x, y)}{iw_y} \\
&= \lim_{w_y \rightarrow 0} \frac{u(x, y + w_y) - u(x, y)}{iw_y} + i \lim_{w_y \rightarrow 0} \frac{v(x, y + w_y) - v(x, y)}{iw_y} \\
&= -i \lim_{w_y \rightarrow 0} \frac{u(x, y + w_y) - u(x, y)}{w_y} + \lim_{w_y \rightarrow 0} \frac{v(x, y + w_y) - v(x, y)}{w_y} \\
&= -iu_y + v_y.
\end{aligned}$$

Setting the two real and imaginary parts of the equations equal (respectively), we get $u_x = v_y$ and $v_x = -u_y$ as desired. $\square$

Problem 2. Use the limit definition to show that $f(z) = \bar{z}$ is not complex differentiable anywhere, and $g(z) = |z|^2$ is not complex differentiable at any $z \neq 0$, but is complex differentiable at $z = 0$.

Proof. Taking the derivative along the real axis, we get

$$\begin{aligned}
f'(z) &= \lim_{w_x \rightarrow 0} \frac{f(x + w_x, y) - f(x, y)}{w_x} \\
&= \lim_{w_x \rightarrow 0} \frac{x + w_x - yi - (x - yi)}{w_x} \\
&= \lim_{w_x \rightarrow 0} \frac{x + w_x - yi - x + yi}{w_x} \\
&= \lim_{w_x \rightarrow 0} \frac{w_x}{w_x} \\
&= \lim_{w_x \rightarrow 0} 1 \\
&= 1,
\end{aligned}$$

and taking the derivative along the imaginary axis, we get

$$\begin{aligned}
 f'(z) &= \lim_{w_y \rightarrow 0} \frac{f(x, y + w_y) - f(x, y)}{w_y} \\
 &= \lim_{w_y \rightarrow 0} \frac{x - i(y + w_y) - (x - iy)}{w_y} \\
 &= \lim_{w_y \rightarrow 0} \frac{x - iy - iw_y - x + iy}{w_y} \\
 &= \lim_{w_y \rightarrow 0} \frac{-iw_y}{w_y} \\
 &= \lim_{w_y \rightarrow 0} -i \\
 &= -i.
 \end{aligned}$$

Since $1 \neq -i$, f is not differentiable. □

Problem 3. Fix a small, positive radius $r < 1$ and let γ_1 and γ_2 be the positively oriented circles of radius r centered at $z = 1$ and $z = 2$, respectively:

$$\gamma_1 = 1 + re^{it}, \quad \gamma_2(t) = 2 + re^{it}, \quad t \in [0, 2\pi].$$

Compute the following line integrals for $j = 1, 2$:

$$(a) \int_{\gamma_j} \frac{dz}{1-z} \quad (b) \int_{\gamma_j} \frac{dz}{2-z} \quad \int_{\gamma_j} \frac{dz}{(1-z)^2} \quad (d) \int_{\gamma_j} \frac{dz}{(2-z)(1-z)}.$$

Hint: for item (d), partial fractions can immediately reduce it to (a) and (b).

Problem 4. Consider the linear operator $A : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ represented in the standard basis by the matrix

$$A = \begin{pmatrix} 2 & 1 & -1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

Note that the spectrum of A is $\sigma(A) = \{1, 2\}$.

- (a) Find the kernel of $(A-1)^2$ and the kernel of $A-2$. These are the generalized eigenspaces of A for the eigenvalues 1 and 2.
- (b) Compute the resolvent matrix $R(z) = (A - z)^{-1}$ for $z \notin \sigma(A)$. Simplify the rational functions in each entry completely.
- (c) For each eigenvalue $\lambda_j = j, j = 1, 2$ compute the operator-valued integral

$$P_j = -\frac{1}{2\pi i} \int_{\gamma_j} R(z) dz,$$

where γ_j is a small positively oriented circle enclosing only λ_j .

- (d) Show that P_1 and P_2 are projections, that their images are the generalized eigenspaces of A ,

$$\operatorname{Im}(P_1) = \ker(A - 1)^2, \quad \operatorname{Im}(P_2) = \ker(A - 2),$$

and that they complement each other, meaning:

$$P_1 + P_2 = 1, \quad P_1 P_2 = 0.$$

P_1 and P_2 are called *Riesz projections* associated with A .