

Polynomial Methods in Combinatorics

Conrad Crowley

Supervisor: Marco Vitturi

March 2022

What are Polynomial Methods?

A collection of techniques in Combinatorics which use polynomials and their rigidity properties to argue a

What are Polynomial Methods?

A collection of techniques in Combinatorics which use polynomials and their rigidity properties to argue a

Example (Rigidity and Interpolation)

- Rigidity: If $P \in \mathbb{R}[X_1, \dots, X_n]$ has degree D and a line ℓ intersects P in more than D points then $\ell \subset Z(P)$.

What are Polynomial Methods?

A collection of techniques in Combinatorics which use polynomials and their rigidity properties to argue a

Example (Rigidity and Interpolation)

- Rigidity: If $P \in \mathbb{R}[X_1, \dots, X_n]$ has degree D and a line ℓ intersects P in more than D points then $\ell \subset Z(P)$.
- Interpolation: We can do parameter-counting arguments using the fact that $\dim \mathbb{R}_{\deg \leq D}[X_1, \dots, X_n] \sim D^n$.

Theorems proven using polynomial methods

We examined proofs of these theorems using the polynomial method:

- Kakeya Conjecture in Finite Fields: If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.

Theorems proven using polynomial methods

We examined proofs of these theorems using the polynomial method:

- Kakeya Conjecture in Finite Fields: If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- Cauchy-Davenport Theorem: $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.

Theorems proven using polynomial methods

We examined proofs of these theorems using the polynomial method:

- Kakeya Conjecture in Finite Fields: If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- Cauchy-Davenport Theorem: $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.
- Joints Problem: A collection of N lines in \mathbb{R}^3 can form at most $N^{3/2}$ joints.

Theorems proven using polynomial methods

We examined proofs of these theorems using the polynomial method:

- Kakeya Conjecture in Finite Fields: If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- Cauchy-Davenport Theorem: $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.
- Joints Problem: A collection of N lines in \mathbb{R}^3 can form at most $N^{3/2}$ joints.
- Szemerédi-Trotter Theorem: Given S points and L lines, there are $\lesssim (SL)^{2/3} + S + L$ point-line incidences. (i.e. (p, ℓ) s.t. $p \in \ell$)

Theorems proven using polynomial methods


We examined proofs of these theorems using the polynomial method:

- Kakeya Conjecture in Finite Fields: If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- Cauchy-Davenport Theorem: $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.
- Joints Problem: A collection of N lines in \mathbb{R}^3 can form at most $N^{3/2}$ joints.
- Szemerédi-Trotter Theorem: Given S points and L lines, there are $\lesssim (SL)^{2/3} + S + L$ point-line incidences. (i.e. (p, ℓ) s.t. $p \in \ell$)
- Circle Tangencies: Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies.


Circle Tangencies

Theorem

Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies.



Circle Tangencies: What's degenerate?



Circle Tangencies: Lifting into \mathbb{R}^3

Theorem

Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies.


We now present a sketch of a recent proof due to Ellenberg, Solymosi, and Zahl. [1]

Sketch Proof: Assume that there are $\gtrsim N^{3/2}$ tangencies with each circle tangent to $\gtrsim N^{1/2}$ other circles in our collection.


For each circle γ in our collection, we define the curve $\beta(\gamma) \subset \mathbb{R}^3$ as:

$$\beta(\gamma) := \left\{ (x, y, z) \mid (x, y) \in \gamma, z = \frac{-1}{\text{Slope of tangent at } (x, y)} \right\}$$


Circle Tangencies: Lifting into \mathbb{R}^3




Circle Tangencies: Lifting into \mathbb{R}^3




Circle Tangencies: Lifting into \mathbb{R}^3



Circle Tangencies: Lifting into \mathbb{R}^3



Circle Tangencies: Tangencies to Incidences




Circle Tangencies: Tangencies to Incidences



Tangency problem in $\mathbb{R}^2 \rightarrow$ Incidences problem in \mathbb{R}^3 .

Circle Tangencies: Tangencies to Incidences



Tangency problem in $\mathbb{R}^2 \rightarrow$ Incidences problem in \mathbb{R}^3 .

Circle Tangencies: Tangent Vectors at Incidences

Let us examine the tangent vectors at a point of incidence between two curves:



Circle Tangencies: Tangent Vectors at Incidences




Circle Tangencies: Tangent Vectors at Incidences



At every point of intersection, the tangent vectors span a vertical plane.


Circle Tangencies: Tangent Vectors at Incidences



At every point of intersection, the tangent vectors span a vertical plane.


Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.




Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.




Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.




Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.



Circle Tangencies: Interpolation


Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.



- Recall that $\dim \mathbb{R}_{\deg \leq D}[X, Y, Z] \sim D^3$.

Circle Tangencies: Interpolation


Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.



- Recall that $\dim \mathbb{R}_{\deg \leq D}[X, Y, Z] \sim D^3$.
- Each incidence point gives one linear equation for the coefficients.


Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.




- Recall that $\dim \mathbb{R}_{\deg \leq D}[X, Y, Z] \sim D^3$.
- Each incidence point gives one linear equation for the coefficients.
- $D^3 \sim N^{3/2} \implies D \sim N^{1/2}$.

Circle Tangencies: Interpolation




Circle Tangencies: Interpolation




- Each curve $\beta(\gamma)$ contains $\gtrsim N^{1/2}$ points of intersection with $Z(P)$.

Circle Tangencies: Interpolation




- Each curve $\beta(\gamma)$ contains $\gtrsim N^{1/2}$ points of intersection with $Z(P)$.
- $\deg \beta(\gamma) = O(1)$ and $\deg P \sim N^{1/2}$

Circle Tangencies: Interpolation




- Each curve $\beta(\gamma)$ contains $\gtrsim N^{1/2}$ points of intersection with $Z(P)$.
- $\deg \beta(\gamma) = O(1)$ and $\deg P \sim N^{1/2}$
- $\implies \beta(\gamma) \subset Z(P)$ by Bézout's Theorem.

Circle Tangencies: Interpolation




- Each curve $\beta(\gamma)$ contains $\gtrsim N^{1/2}$ points of intersection with $Z(P)$.
- $\deg \beta(\gamma) = O(1)$ and $\deg P \sim N^{1/2}$
- $\implies \beta(\gamma) \subset Z(P)$ by Bézout's Theorem.

Circle Tangencies: Interpolation




Circle Tangencies: Interpolation



- Tangent vectors span the z -axis at tangencies, so $\partial_z P = 0$ on all incidences!
- $\implies Z(\partial_z P)$ also contains all incidences.
- $\implies P(X, Y, Z) = Q(X, Y)$.

Circle Tangencies: Interpolation



$\deg Q \sim N^{1/2}$, but $Z(Q)$ contains N circles. Contradiction!

Circle Tangencies: Recap of Argument

- 1 Assume there are $\gtrsim N^{3/2}$ tangencies in our collection.
- 2 Lift curves into \mathbb{R}^3 and change into an incidence problem.
- 3 Use a low degree polynomial P to interpolate these points. (parameter-counting)
- 4 Argue that if $Z(P)$ contains $\gtrsim N^{1/2}$ points of $\beta(\gamma)$ then $\beta(\gamma) \subset Z(P)$. (rigidity)
- 5 Use structure of the objects to argue $P(X, Y, Z) = Q(X, Y)$
- 6 Contradiction as degree of Q is $N^{1/2}$ but contains N circles.



Jordan S. Ellenberg, Jozsef Solymosi, and Joshua Zahl.

New bounds on curve tangencies and orthogonalities, 2016.