

Polynomial Methods in Combinatorics

Conrad Crowley

Supervisor: Marco Vitturi

March 2022

What are Polynomial Methods?

A collection of techniques in Combinatorics which use polynomial interpolation and rigidity properties of polynomials to control the size of collections of objects with a certain structure.

What are Polynomial Methods?

A collection of techniques in Combinatorics which use polynomial interpolation and rigidity properties of polynomials to control the size of collections of objects with a certain structure.

Example (Rigidity and Interpolation)

- **Rigidity:** If $P \in \mathbb{R}[X_1, \dots, X_n]$ has degree D and a line ℓ intersects $Z(P)$ in more than D points then $\ell \subset Z(P)$.

What are Polynomial Methods?

A collection of techniques in Combinatorics which use polynomial interpolation and rigidity properties of polynomials to control the size of collections of objects with a certain structure.

Example (Rigidity and Interpolation)

- **Rigidity:** If $P \in \mathbb{R}[X_1, \dots, X_n]$ has degree D and a line ℓ intersects $Z(P)$ in more than D points then $\ell \subset Z(P)$.
- **Interpolation:** We can do parameter-counting arguments using the fact that $\dim \mathbb{R}_{\deg \leq D}[X_1, \dots, X_n] \sim D^n$.

Theorems proven using polynomial methods

Below lists results that can be proved using the polynomial method:

- **Kakeya Conjecture in Finite Fields:** If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.

Theorems proven using polynomial methods

Below lists results that can be proved using the polynomial method:

- **Kakeya Conjecture in Finite Fields:** If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- **Cauchy-Davenport Theorem:** $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.

Theorems proven using polynomial methods

Below lists results that can be proved using the polynomial method:

- **Kakeya Conjecture in Finite Fields:** If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- **Cauchy-Davenport Theorem:** $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.
- **Joints Problem:** A collection of N lines in \mathbb{R}^3 can form at most $N^{3/2}$ joints. A joint is a point which lies in three non-coplanar lines.

Theorems proven using polynomial methods

Below lists results that can be proved using the polynomial method:

- **Kakeya Conjecture in Finite Fields:** If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- **Cauchy-Davenport Theorem:** $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.
- **Joints Problem:** A collection of N lines in \mathbb{R}^3 can form at most $N^{3/2}$ joints. A joint is a point which lies in three non-coplanar lines.
- **Szemerédi-Trotter Theorem:** Given S points and L lines, there are $\lesssim (SL)^{2/3} + S + L$ point-line incidences. (i.e. (p, ℓ) s.t. $p \in \ell$)

Theorems proven using polynomial methods

Below lists results that can be proved using the polynomial method:


- **Kakeya Conjecture in Finite Fields:** If $A \subset \mathbb{F}^n$ contains a line in every direction then $|A| \gtrsim |\mathbb{F}|^n$.
- **Cauchy-Davenport Theorem:** $|A + B| \geq \min\{p, |A| + |B| - 1\}$ where $A, B \subset \mathbb{Z}_p$ and $A + B := \{a + b \mid a \in A, b \in B\}$.
- **Joints Problem:** A collection of N lines in \mathbb{R}^3 can form at most $N^{3/2}$ joints. A joint is a point which lies in three non-coplanar lines.
- **Szemerédi-Trotter Theorem:** Given S points and L lines, there are $\lesssim (SL)^{2/3} + S + L$ point-line incidences. (i.e. (p, ℓ) s.t. $p \in \ell$)
- **Circle Tangencies:** Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies.

Circle Tangencies

Theorem


Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies^a.

^aA tangency is a pair of circles (γ, γ') that are tangent.



Circle Tangencies: What's degenerate?

Collection of N circles with $\sim N^2$ tangencies:



Circle Tangencies: Lifting into \mathbb{R}^3

Theorem

Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies.

We now present a sketch of a recent proof due to Ellenberg, Solymosi, and Zahl. [1]

Circle Tangencies: Lifting into \mathbb{R}^3

Theorem

Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies.

We now present a sketch of a recent proof due to Ellenberg, Solymosi, and Zahl. [1]


Sketch Proof: Assume:

- There are $\gtrsim N^{3/2}$ tangencies.
- Collection is uniform: each circle tangent to $\gtrsim N^{1/2}$ other circles.


For each circle γ in our collection, we define the curve $\beta(\gamma) \subset \mathbb{R}^3$ as:

$$\beta(\gamma) := \left\{ (x, y, z) \mid (x, y) \in \gamma, z = \frac{1}{\text{Slope of tangent at } (x, y)} \right\}$$

Circle Tangencies: Lifting into \mathbb{R}^3




Circle Tangencies: Lifting into \mathbb{R}^3




Circle Tangencies: Lifting into \mathbb{R}^3




Circle Tangencies: Lifting into \mathbb{R}^3



Circle Tangencies: Tangencies to Incidences




Circle Tangencies: Tangencies to Incidences



Intersection \implies z co-ords equal \implies circles are tangent.

Tangency problem in $\mathbb{R}^2 \iff$ Incidences problem in \mathbb{R}^3 .

Circle Tangencies: Tangencies to Incidences



Intersection \implies z co-ords equal \implies circles are tangent.

Tangency problem in $\mathbb{R}^2 \iff$ Incidences problem in \mathbb{R}^3 .

Circle Tangencies: Tangent Vectors at Incidences

Let us examine the tangent vectors at a point of incidence between two curves:



Circle Tangencies: Tangent Vectors at Incidences




Circle Tangencies: Tangent Vectors at Incidences



At every point of intersection, the tangent vectors span a vertical plane.


Circle Tangencies: Tangent Vectors at Incidences



At every point of intersection, the tangent vectors span a vertical plane.


Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.




Circle Tangencies: Interpolation

We can find a polynomial P such that all $N^{3/2}$ incidences are contained in $Z(P)$.




Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.



Circle Tangencies: Interpolation


Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.



- Recall that $\dim \mathbb{R}_{\deg \leq D}[X, Y, Z] \sim D^3$.

Circle Tangencies: Interpolation


Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.



- Recall that $\dim \mathbb{R}_{\deg \leq D}[X, Y, Z] \sim D^3$.
- Each incidence point gives one linear equation for the coefficients.


Circle Tangencies: Interpolation

Let P be a polynomial such that all $N^{3/2}$ incidences are contained in $Z(P)$.




- Recall that $\dim \mathbb{R}_{\deg \leq D}[X, Y, Z] \sim D^3$.
- Each incidence point gives one linear equation for the coefficients.
- $D^3 \sim N^{3/2} \implies D \sim N^{1/2}$.

Circle Tangencies: $\beta(\gamma) \subset Z(P)$




Circle Tangencies: $\beta(\gamma) \subset Z(P)$




- Each curve $\beta(\gamma)$ intersects $Z(P)$ at $\gtrsim N^{1/2}$ points.

Circle Tangencies: $\beta(\gamma) \subset Z(P)$




- Each curve $\beta(\gamma)$ intersects $Z(P)$ at $\gtrsim N^{1/2}$ points.
- But $\deg \beta(\gamma) = O(1)$ and $\deg P \sim N^{1/2}$

Circle Tangencies: $\beta(\gamma) \subset Z(P)$




- Each curve $\beta(\gamma)$ intersects $Z(P)$ at $\gtrsim N^{1/2}$ points.
- But $\deg \beta(\gamma) = O(1)$ and $\deg P \sim N^{1/2}$
- $\implies \beta(\gamma) \subset Z(P)$ by Bézout's Theorem! (rigidity)

Circle Tangencies: $\beta(\gamma) \subset Z(P)$




- Each curve $\beta(\gamma)$ contains $\gtrsim N^{1/2}$ points of intersection with $Z(P)$.
- $\deg \beta(\gamma) = O(1)$ and $\deg P \sim N^{1/2}$
- $\implies \beta(\gamma) \subset Z(P)$ by Bézout's Theorem.

Circle Tangencies: Tangent Vectors




Circle Tangencies: Tangent Vectors




- Before we showed tangent space at incidences is vertical, so $\partial_z P = 0$ on all incidences!

Circle Tangencies: Tangent Vectors




- Before we showed tangent space at incidences is vertical, so $\partial_z P = 0$ on all incidences!
- $\implies Z(\partial_z P)$ also contains all incidences.

Circle Tangencies: Tangent Vectors



- Before we showed tangent space at incidences is vertical, so $\partial_z P = 0$ on all incidences!
- $\implies Z(\partial_z P)$ also contains all incidences.
- If $\deg P$ minimal $\implies P(X, Y, Z) = Q(X, Y)$.

Circle Tangencies: Contradiction



Recall that $\deg P = \deg Q \sim N^{1/2}$, but $Z(Q)$ contains N circles.
Contradiction!


Circle Tangencies: Recap of Argument

Theorem

Given a (suitably non-degenerate) collection of N circles in \mathbb{R}^2 , they determine $\lesssim N^{3/2}$ tangencies.


- 1 Assume there are $\gtrsim N^{3/2}$ tangencies.
- 2 Lift curves into \mathbb{R}^3 and change into an incidence problem.
- 3 Use a low degree polynomial P to interpolate these points.
(parameter-counting)
- 4 Argue that if $Z(P)$ contains $\gtrsim N^{1/2}$ points of $\beta(\gamma)$ then $\beta(\gamma) \subset Z(P)$. (rigidity)
- 5 Use structure of the objects to argue $P(X, Y, Z) = Q(X, Y)$.
- 6 Contradiction as degree of Q is $\sim N^{1/2}$ but contains N circles.

Circle Tangencies: New Proof




Polynomial Partitioning: We can find a polynomial P of degree D such that $Z(P)$ partitions \mathbb{R}^3 into $\sim D^3$ cells. (parameter-counting + Borsuk-Ulam)

Circle Tangencies: New Proof



Polynomial Partitioning: We can find a polynomial P of degree D such that $Z(P)$ partitions \mathbb{R}^3 into $\sim D^3$ cells. (parameter-counting + Borsuk-Ulam)

Circle Tangencies: New Proof



Polynomial Partitioning: We can find a polynomial P of degree D such that $Z(P)$ partitions \mathbb{R}^3 into $\sim D^3$ cells. (parameter-counting + Borsuk-Ulam)

Polynomial Partitioning:

- We can find a polynomial P of degree D such that $Z(P)$ partitions \mathbb{R}^3 into $\sim D^3$ cells. (parameter-counting + Borsuk-Ulam)
- Each cell intersects $\lesssim \frac{N}{D^2}$ curves $\beta(\gamma)$.

Set of $\beta(\gamma)$'s can be partitioned:

$$C_1 = \{\beta(\gamma) \not\subset Z(P)\} \text{ and } C_2 = \{\beta(\gamma) \subset Z(P)\}$$

Total incidences of $\beta(\gamma)$'s = $I(C_1, C_1) + I(C_1, C_2) + I(C_2, C_2)$

Circle Tangencies: New Proof

$I(C_1, C_1)$ = incidences between curves inside the cells. Each cell contains $\lesssim \frac{N}{D^2}$ curves so:

$$\begin{aligned} I(C_1, C_1) &\leq \sum_{\text{cells}} \left(\frac{N}{D^2} \right)^2 \\ &= D^3(N^2 D^{-4}) = N^2 D^{-1} \end{aligned}$$

Circle Tangencies: New Proof

$I(C_1, C_2)$ = incidences between a curve β' in $Z(P)$ and a curve β not in $Z(P)$.

$$I(C_1, C_2) = \sum_{\beta \in C_1} \sum_{\beta' \in C_2} \mathbb{1}[\beta \cap \beta' \neq \emptyset]$$

Each $\beta(\gamma) \in C_1$ can intersect $Z(P)$ at most $\lesssim D$ times. (Bézout)

$$\lesssim \sum_{\beta \in C_1} D \lesssim ND$$

Circle Tangencies: New Proof

$I(C_2, C_2)$ = incidences between $\beta(\gamma)$'s both in $Z(P)$. We can assume P was of minimal degree for the partitioning. We consider $\partial_Z P$:

Case 1: $Z(\partial_Z P)$ contains all the curves of C_2 . Proceed by same argument in previous proof to yield a bound of $\lesssim ND$.

Circle Tangencies: New Proof

$I(C_2, C_2) =$ incidences between $\beta(\gamma)$'s both in $Z(P)$. We can assume P was of minimal degree for the partitioning. We consider $\partial_z P$:

Case 1: $Z(\partial_z P)$ contains all the curves of C_2 . Proceed by same argument in previous proof to yield a bound of $\lesssim ND$.

Case 2: $C_2 \setminus Z(\partial_z P)$ is non-empty. There are $\lesssim D$ incidences between curves in $C_2 \setminus Z(\partial_z P)$ and the zero set. We repeat the process up to D times until we are in case 1, accumulating $\lesssim \sum D = D^2$ incidences.

Circle Tangencies: New Proof

Adding these up we get the number of tangencies to be:

$$I(C_1, C_1) + I(C_1, C_2) + I(C_2, C_2) \lesssim N^2 D^{-1} + ND + D^2$$

Circle Tangencies: New Proof

Adding these up we get the number of tangencies to be:

$$I(C_1, C_1) + I(C_1, C_2) + I(C_2, C_2) \lesssim N^2 D^{-1} + ND + D^2$$

We optimize D now by setting $N^2 D^{-1} \sim ND \implies D \sim N^{1/2}$.

We achieve:

$$\lesssim N^{3/2}$$

Circle Tangencies: Recap of New Proof

- ① Lift curves into \mathbb{R}^3 and change into an incidence problem.
- ② Use a low degree polynomial P to partition \mathbb{R}^3 into D^3 cells, each intersecting $\lesssim ND^{-2}$ curves. (parameter-counting + Borsuk-Ulam)
- ③ Use a trivial bound in each cell and sum over all cells.
- ④ Deal with curves contained in the zero set using algebraic tools.
- ⑤ Choose D to optimise inequality and achieve $\lesssim N^{3/2}$ bound.

Thank you for your attention.
Any questions?



Jordan S. Ellenberg, Jozsef Solymosi, and Joshua Zahl.

New bounds on curve tangencies and orthogonalities, 2016.