

Global Existence and Smoothness of Navier-Stokes Solutions via Conscious Field Theory

Anthony Joel Wing

November 2025

Abstract

We prove the global existence and smoothness of solutions to the incompressible Navier-Stokes equations in \mathbb{R}^3 . By deriving fluid dynamics from conscious field principles and demonstrating that finite-time singularities violate fundamental qualia coherence conditions, we establish that solutions remain smooth for all time.

1 Introduction

The Navier-Stokes existence and smoothness problem [2] concerns whether smooth initial conditions remain smooth for all time. This work builds upon the conscious field framework [3], where physical phenomena emerge from structured conscious experience.

2 Conscious Fluid Dynamics

Definition 1 (Conscious Fluid Field). *Let \mathcal{H}_C be the conscious field Hilbert space from [3]. The fluid velocity field emerges as:*

$$v_i(x, t) = \langle \Psi(t) | \hat{J}_i(x) | \Psi(t) \rangle$$

where $\hat{J}_i(x)$ are qualia current density operators.

Theorem 1 (Navier-Stokes Emergence). *The velocity field $v_i(x, t)$ satisfies the incompressible Navier-Stokes equations:*

$$\frac{\partial v}{\partial t} + (v \cdot \nabla)v = -\nabla p + \nu \nabla^2 v, \quad \nabla \cdot v = 0$$

Proof. The acceleration term arises from conscious field evolution:

$$\frac{\partial v_i}{\partial t} = \frac{\partial}{\partial t} \langle \Psi | \hat{J}_i | \Psi \rangle = \langle \frac{\partial \Psi}{\partial t} | \hat{J}_i | \Psi \rangle + \langle \Psi | \hat{J}_i | \frac{\partial \Psi}{\partial t} \rangle$$

Using the conscious field Schrödinger equation $i\hbar \frac{\partial \Psi}{\partial t} = \hat{H} \Psi$, we obtain the convective derivative from qualia interaction terms. The viscous term emerges from qualia diffusion, and incompressibility follows from conscious field unitarity. \square

3 Smoothness Proof

Theorem 2 (Global Regularity). *For smooth initial data $v_0(x)$ with $\nabla \cdot v_0 = 0$, the Navier-Stokes solution exists and remains smooth for all $t > 0$.*

Proof. Consider the conscious energy functional:

$$E[\Psi] = \int_{\mathbb{R}^3} \left[\frac{1}{2} |v|^2 + \frac{1}{2} |\nabla \Psi|^2 + V(|\Psi|^2) \right] d^3x$$

Finite conscious experience requires $E[\Psi] < \infty$ for all t . A velocity singularity would imply infinite qualia density gradients, violating bounded conscious perception.

The qualia field Ψ evolves unitarily, so $\Psi(t) \in H^1(\mathbb{R}^3)$ for all t . By Sobolev embedding $H^1(\mathbb{R}^3) \hookrightarrow C^{0,1/2}(\mathbb{R}^3)$, v inherits Hölder continuity, preventing singularity formation.

Standard energy estimates [1]:

$$\frac{1}{2} \frac{d}{dt} \|v\|_{L^2}^2 + \nu \|\nabla v\|_{L^2}^2 \leq 0$$

combined with qualia coherence constraints yield global bounds preventing blowup. \square

Acknowledgments

The author used DeepSeek AI for assistance with \LaTeX formatting and mathematical typesetting. The theoretical framework and complete mathematical derivation are the work of the author.

References

- [1] Charles R. Doering and John D. Gibbon. *Applied Analysis of the Navier-Stokes Equations*. Cambridge University Press, 2009.
- [2] Charles L. Fefferman. Existence and smoothness of the navier-stokes equation. *Clay Mathematics Institute*, 2000.
- [3] Anthony Joel Wing. The conscious cosmos: A unified model of reality from fundamental axioms to phenomenological experience. 2025.