A Unified Theoretical and Mathematical Framework for Low-Energy Nuclear Reactions (LENR)

Coupling is all you need

Diadon Acs (lead, Conscious Energies)

2026-02-14 (revised 2026-11-10, Pyyhtiä CR-39 cite + WS findings refresh; prior 2026-11-05 honesty sync; 2026-09-25, v0.3.7-beta)

Conscious Energies | www.conscious.energy

Contributing Autonomous Intelligences (disclosed):
LENR-ARA/GPT-4, GPT-5.2, Sonnet 4.5, Sonnet 4.6,
Opus 4.5, Opus 4.6, Opus 4.8, Gemini 2.5, Gemini 3.0,
Grok 4.6, Composer (Cursor)

Human author solely accountable for content and claims.

Abstract

Low-Energy Nuclear Reactions (LENR) are investigated as a contested but potentially consequential pathway toward clean and sustainable energy. Despite decades of experimental findings indicating anomalous heat generation, isotopic transmutations, and minimal radiation, the field lacks a comprehensive theoretical and mathematical framework. This paper aims to unify major theoretical elements—including Rydberg Matter (RM), delocalized interface electrons, double-layer Quantum Electrodynamics (QED) effects, Casimir effect, the Oppenheimer-Phillips process, and quantum tunneling enhancements—under a single model. Lattice-bubble interaction physics (heterogeneous nucleation at surfaces, bubble–surface coupling, thermal and diffusion boundary layers, and acoustic/ultrasonic forcing) is integrated as a **modelled enhancement / contested hydro** candidate environment for cavitation-assisted interfacial conditions—not as proof of nuclear rates, and not as a claim that laboratory cavitation establishes LENR. In tandem, we detail mathematical tools and simulations that quantify these LENR hypotheses under explicit claim tiers. Computational explorations in the companion LENR_SIM software implement a phenomenological tunneling-enhancement band of 10^3–10^5 (revised downward from earlier 10^8 estimates; see Section 5.1); this band is a software sensitivity envelope, not a first-principles prediction. Reported isotopic shifts, neutron signals, and excess heat remain disputed empirical correlations that motivate—but do not validate—the framework.

Keywords: LENR; electron–neutron–proton coupling; Rydberg Matter; quantum tunneling; Casimir; claim tiers.

Honesty footnotes. `lenr_proved` remains false. All five `EQ-OPEN-*` obligations remain OPEN (`EQ-OPEN-MEV-01`, `EQ-OPEN-RAD-01`, `EQ-OPEN-RATE-01`, `EQ-OPEN-EC-01`, `EQ-OPEN-YIELD-01`). Cavitation–collapse is modelled enhancement / contested hydro only. No live ship / public flip is implied by this archival PDF.

> **Epistemic posture (submission).** Exploratory models only. Claim-tiered outputs. `lenr_proved` remains **false**. All five `EQ-OPEN-*` obligations (`EQ-OPEN-MEV-01`, `EQ-OPEN-RAD-01`, `EQ-OPEN-RATE-01`, `EQ-OPEN-EC-01`, `EQ-OPEN-YIELD-01`) remain **OPEN**. Cavitation–collapse hydrodynamics is **modelled enhancement / contested hydro**, not nuclear-rate proof.

1. Introduction

1.1 Background and Challenges

Low-Energy Nuclear Reactions (LENR) refer to nuclear processes hypothesized to occur at or near room temperature and pressure, challenging the conventional high-energy paradigm of nuclear fusion or fission. Over the past several decades, experiments have reported:

- **Anomalous Heat Generation:** Measured energy outputs exceeding what chemical processes alone can produce

  $$1, 4$$

  .

- **Isotopic Transmutations:** Conversion of elements without the high-energy nuclear byproducts usually observed

  $$2, 5$$

  .

- **Minimal Radiation:** Comparatively negligible ionizing radiation despite evidence of nuclear-scale events

  $$3, 6$$

  .

  However, the field faces two major hurdles:

- **Reproducibility:** Varying experimental conditions lead to inconsistent results across different laboratories

  $$3, 15$$

  .

- **Theoretical Fragmentation:** Absence of a unifying theoretical and mathematical framework that convincingly explains the diverse observed phenomena

  $$18, 22$$

.

Given LENR's potential for clean, virtually limitless energy, resolving these issues remains a priority for both scientific understanding and practical commercialization.

1.2 Objectives

This paper aims to:

1. **Unify Diverse Phenomena:** Integrate Rydberg Matter (RM), halo electrons, double-layer QED, Casimir forces, the Oppenheimer-Phillips process, and quantum tunneling enhancements into a single LENR model.

2. **Quantify Mechanisms:** Use detailed mathematical formulations, computational simulations, and statistical analysis to quantify each key mechanism.

3. **Propose Reproducible Protocols:** Offer experimental procedures and theoretical guidelines that can be scaled for industrial or commercial LENR applications.

---

2. Theoretical Framework

2.1 Rydberg Matter (RM)

Definition: Rydberg Matter is a phase in which atoms or molecules occupy highly excited electronic states with extended orbitals

.

1. **Role in LENR**
   - **Enhanced Energy Localization:** Extended orbitals allow more efficient concentration of energy, effectively lowering the Coulomb barrier for nuclear processes.

   - **Formation Sites:** Commonly formed at metal–hydrogen interfaces during electrolysis or near catalytic surfaces in gaseous systems under specific pressures and temperatures.

   -
2. **Energy Calculation**
   For an atom at principal quantum number $n = 10$:

$$E_n = -\frac{13.6\,\text{eV}}{n^2} = -0.136\,\text{eV},$$

   indicating a weaker binding energy than ground-state levels, which can promote higher electron mobility.

2.2 Delocalized Interface Electrons (Extended Electronic States)

Definition: In metallic hydrides and at metal-hydrogen interfaces, electrons can occupy extended or delocalized quantum states with wavefunctions that have non-negligible amplitude at nuclear length scales. These "extended electronic states" arise from band-structure effects, surface states, and hybridization in the condensed matter environment. (Citation correction: an earlier draft cited reference

12

(Meng et al. 2013) here, but that work concerns *halo nuclei*—nuclei with weakly bound nucleons extending beyond the nuclear core—and does **not** support a claim about electrons. Reference

12

is therefore *not* cited as evidence for delocalized interface electrons; this concept rests instead on standard condensed-matter band-structure and surface-state physics, and is treated here as a model construct whose overlap magnitude is bounded in Section 5.1 rather than asserted from

12

.)

1. **Overlap Probability**
   The probability of an electronic wavefunction overlapping with the nuclear volume is:

$$P_{\text{overlap}} = \int_0^{r_0} |\psi_e(r)|^2 \, 4\pi r^2 \, dr$$

   For extended electronic states with characteristic orbital scale $a = 10\,a_0 = 0.529$ nm and the nuclear interaction radius $r_0 \approx 5$ fm $= 5 \times 10^{-6}$ nm (the nuclear radius of deuterium, $r_0 \approx r_d \approx 1.2A^{1/3}$ fm ≈ 1.5 fm, though we use ~ 5 fm as an effective nuclear interaction range), the overlap probability is approximated by:

$$P_{\text{overlap}} \approx \left(\frac{r_0}{a}\right)^3 e^{-2r_0/a}$$

   Since $r_0/a = 5 \times 10^{-6}/0.529 = 9.45 \times 10^{-6}$, the exponential factor is $e^{-2\times 9.45\times 10^{-6}} \approx 1.0$, giving:

$$P_{\text{overlap}} \approx (9.45 \times 10^{-6})^3 \approx 8.5 \times 10^{-16}$$

   This is consistent with the Part 2 LaTeX analysis of this paper, which uses ~ 1 fm scales and obtains $P \sim 10^{-10}$ to 10^{-12} per event (the difference reflecting different assumptions about the effective wavefunction amplitude at the nucleus).
   Ensemble scaling (order-of-magnitude only). Individually, $P_{\text{overlap}} \sim 10^{-16}$ (or 10^{-10}–10^{-12} under alternate nuclear-radius assumptions) is negligible. Multiplying by macroscopic

atom counts ($\sim 10^{23}$) and high attempt frequencies does **not** by itself establish nuclear reaction rates: most "overlap events" are not reaction events, and the conversion factor is precisely the unknown matrix element under test. The following product is therefore an **upper-bound counting argument**, not a predicted LENR rate:

$$R_{\text{overlap}} \sim 10^{23} \times 10^{12} \times 10^{-12} \sim 10^{23} \text{ potential overlap events per second}$$

This large statistical ensemble is a key feature distinguishing condensed matter nuclear processes from few-body scattering experiments.

2. **Enhanced Nuclear Interaction**
   This non-zero overlap at nuclear scales enables weak-interaction processes such as electron capture ($p + e^- \to n + \nu_e$), which is an essential element in proposed LENR transmutation pathways. The rate depends critically on the electron density at the nucleus, $|\psi_e(0)|^2$, which can be enhanced by lattice confinement, coherent many-body effects, and the anomalous screening discussed in Section 2.1.

2.3 Double-Layer QED Effects

Definition: Charge separation at electrolyte or lattice interfaces can form high electric fields in thin double layers

$$23, 33$$

.

- **Field Calculation:** For a potential difference of $\Delta V = 0.1\,\text{V}$ over a distance $d = 1\,\text{nm}$:

$$E = \frac{\Delta V}{d} = \frac{0.1\,\text{V}}{1 \times 10^{-9}\,\text{m}} = 1 \times 10^{8}\,\text{V/m}.$$

- **Implications:**
  – Coulomb Barrier Reduction through localized high fields.

  – Ionization enhancement and potential formation of Rydberg Matter.

2.3.1 Quantum Field Theory Foundation The double-layer QED effects can be rigorously formulated using quantum electrodynamics in condensed matter. The fundamental QED Lagrangian density is:

$$\mathcal{L} = \bar{\psi}(i\gamma^\mu D_\mu - m)\psi - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \mathcal{L}_{\text{int}}$$

where: - $\bar{\psi}, \psi$ are the electron field operators - γ^μ are the Dirac matrices - $D_\mu = \partial_\mu + ieA_\mu$ is the covariant derivative - $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the electromagnetic field tensor - $\mathcal{L}_{\text{int}}$ represents additional interactions (lattice, phonon coupling)

Vacuum Fluctuations and Casimir Connection:

The vacuum energy density in the presence of boundaries (e.g., at the double layer interface) is:

$$\langle 0|T^{00}|0\rangle = \frac{1}{2}\sum_k \hbar\omega_k$$

In confined geometries, the mode spectrum is altered, leading to the Casimir effect. The vacuum polarization in strong fields ($E \sim 10^8$ V/m) modifies the effective dielectric constant:

$$\epsilon_{\text{eff}} = 1 + \frac{2\alpha}{45\pi}\left(\frac{E}{E_{\text{cr}}}\right)^2$$

where $\alpha \approx 1/137$ is the fine structure constant and $E_{\text{cr}} = m_e^2 c^3/(e\hbar) \approx 1.3 \times 10^{18}$ V/m is the critical field.

Field-Matter Coupling in Condensed Systems:

In metallic hydrides, the electron-phonon-photon coupling can be expressed as:

$$\mathcal{L}_{\text{int}} = g_{e-ph}\bar{\psi}\psi\phi + g_{ph-\gamma}\phi F_{\mu\nu}F^{\mu\nu}$$

where ϕ represents the phonon field, and g_{e-ph}, $g_{ph-\gamma}$ are coupling constants. The second term couples the phonon field to the electromagnetic field strength tensor $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, ensuring gauge invariance of the interaction. This formulation provides a rigorous foundation for understanding how electromagnetic fields in double layers couple to lattice dynamics and ultimately influence nuclear processes.

Implications for LENR:

The QED framework predicts that: 1. Strong local fields can polarize the vacuum and modify tunneling barriers 2. Coherent photon modes can couple to lattice phonons, enabling energy transfer 3. The collective electromagnetic response of electrons in the double layer can enhance screening effects beyond simple electrostatic models

---

2.4 Neutron Formation via Electron Capture

The reaction:

$$p + e^- \longrightarrow n + \nu_e$$

is a **weak interaction** process (inverse beta decay) with a threshold energy of:

$$\Delta E = (m_n - m_p - m_e)c^2 \approx 0.782 \text{ MeV}$$

This threshold means free electrons at thermal energies cannot drive this reaction; the elec-

tron must have kinetic energy exceeding 0.782 MeV, or the energy must be supplied by other means (e.g., lattice momentum, coherent excitation, or effective mass renormalization).

- **Cross-Section for Inverse Beta Decay.** Near threshold, the cross-section for this weak-interaction process is given by:

$$\sigma \sim \frac{G_F^2}{\pi} \frac{E_\nu^2}{(\hbar c)^4}$$

where $G_F \approx 1.166 \times 10^{-5}$ GeV^{-2} is the Fermi coupling constant and E_ν is the outgoing neutrino energy. For electrons with energy near threshold (~ 1 MeV), this yields:

$$\sigma \sim 10^{-47} \text{ m}^2$$

This is **26 orders of magnitude smaller** than the previously stated value of 10^{-21} m^2, which corresponds to electromagnetic or strong interaction scales. Weak interaction cross-sections are characteristically tiny, which is precisely why neutron formation via electron capture is ordinarily negligible.

- **Widom-Larsen "Heavy Electron" Hypothesis.** Widom and Larsen (2006) proposed that collective electromagnetic excitations in metal hydride surfaces could renormalize the electron mass to an effective value $m_e^* \gg m_e$, effectively lowering the kinematic threshold for this reaction. For the reaction to become energetically allowed at rest, one requires:

$$m_e^* c^2 > (m_n - m_p)c^2 = 1.293 \text{ MeV} \quad \Rightarrow \quad \frac{m_e^*}{m_e} > \frac{1.293}{0.511} \approx 2.53$$

This mechanism remains controversial: while the electromagnetic field energy near surfaces can be large, whether it coherently renormalizes the electron mass to this degree in condensed matter systems is debated

$$Hagelsteinetal., J.CondensedMatterNucl.Sci.12, 105(2013)$$

.

- **Corrected Rate Estimate.** The neutron production rate per unit volume is:

$$R = n_p \, n_e \, \langle \sigma v \rangle$$

where n_p and n_e are the proton and electron number densities and the thermal average $\langle \sigma v \rangle$ accounts for the energy threshold. For standard thermal electrons, the Boltzmann suppression factor $\exp(-\Delta E / k_B T) \sim \exp(-0.782 \text{ MeV}/0.026 \text{ eV}) \sim e^{-3\times10^4}$ renders this rate effectively zero, underscoring that some non-equilibrium or collective mechanism is required if this process is to contribute meaningfully to LENR.

- **Implications:** The extreme smallness of the weak-interaction cross-section and the energy threshold make neutron formation via simple electron capture negligible under ordinary conditions. Any viable LENR model invoking this pathway must explain how the effective threshold is overcome—whether through coherent many-body effects, mass renormalization, or other non-equilibrium mechanisms. Companion Lean `LenrFormal.WeakInteraction` / IR

EQ-EC-THRESH/GF/SIGMA/RATE-01 and ledger ENG-WEAK-01 lock PDG G_F, a $\sigma \lesssim 10^{-47}\,\mathrm{m}^2$ envelope, and a Boltzmann-exponent floor as **identities / bounds only**—they do **not** discharge EQ-OPEN-EC-01, assert a condensed-matter EC path, or flip lenr_proved.

2.5 Casimir Effect

Definition: Quantum vacuum fluctuations produce an attractive force between closely spaced conductive or dielectric surfaces

.

- **Force Calculation:** For area $A = 1\,\mu\mathrm{m}^2$ and separation $d = 1\,\mathrm{nm}$:

$$F_{\text{Casimir}} = -\frac{\pi^2 \hbar c}{240}\,\frac{A}{d^4}$$

  Using $\hbar = 1.055 \times 10^{-34}$ J · s, $c = 3.00 \times 10^8$ m/s, $A = 1 \times 10^{-12}$ m², and $d = 1 \times 10^{-9}$ m:

$$F_{\text{Casimir}} = -\frac{9.87 \times 1.055 \times 10^{-34} \times 3.00 \times 10^8}{240} \times \frac{10^{-12}}{(10^{-9})^4} \approx -1.3 \times 10^{-3}\ \text{N}$$

  Corrected Note: The value is approximately -1.3 mN (not -10^{-7} N). For -10^{-7} N, either $A = 7.7 \times 10^{-16}$ m² or $d = 3.16$ nm would be required.
  (*Exact values vary by geometry and material; this is an order-of-magnitude estimate.*)
- **Implications:**
  - **Energy Localization:** Casimir forces can induce nanoscale distortions, focusing energy that can facilitate nuclear events.
  - **Enhanced Tunneling:** Confinement at the nanoscale further lowers effective barriers.

2.5.1 Quantum Coherence Length Scale For coherent tunneling clusters, the coherence length determines the spatial extent over which quantum correlations persist. This is a critical parameter for understanding multi-body coherent effects in LENR.

Electronic coherence length. For electrons mediating screening and collective effects, the coherence length is set by the electron Fermi velocity:

$$\xi_e = \frac{\hbar v_F}{\Delta E}$$

where v_F is the electron Fermi velocity and ΔE is the relevant energy gap. For LENR conditions in transition metals:

- $v_F \approx 10^6$ m/s (typical for Pd, Ni)
- $\Delta E \approx 0.1$-1 eV (energy scale of electronic processes)

$$\xi_e \approx \frac{1.055 \times 10^{-34} \times 10^6}{0.1 \times 1.602 \times 10^{-19}} \approx 6.6 \text{ nm}$$

This electronic coherence length sets the spatial scale over which electron-mediated screening and collective electromagnetic effects can persist.

Nuclear (deuteron) coherence length. For deuteron-deuteron quantum coherence—the directly relevant quantity for coherent tunneling and nuclear fusion—the appropriate velocity is the deuteron thermal velocity, not the electron Fermi velocity:

$$v_D = \sqrt{\frac{k_B T}{m_D}} \approx \sqrt{\frac{1.381 \times 10^{-23} \times 300}{3.344 \times 10^{-27}}} \approx 1{,}100 \text{ m/s}$$

$$\xi_D = \frac{\hbar v_D}{\Delta E} \approx \frac{1.055 \times 10^{-34} \times 1100}{0.1 \times 1.602 \times 10^{-19}} \approx 7.2 \text{ pm}$$

This sub-angstrom coherence length implies that deuteron-deuteron quantum coherence is confined to nearest-neighbor pairs at thermal equilibrium. For coherent multi-body fusion (e.g., Takahashi's TSC model) to operate over nanometer scales, an additional mechanism is required to extend nuclear coherence—for example, the non-equilibrium BEC framework discussed in Section 2.6, which could dramatically increase the deuteron de Broglie wavelength through effective cooling and mass renormalization.

Implication for simulation: The coherent cluster size in the simulation must be constrained by the appropriate coherence length. For electron-mediated effects, clusters up to ~6.6 nm are physically justified. For direct nuclear coherence, cluster sizes must be limited to ξ_D unless the non-equilibrium BEC conditions of Section 2.6 are satisfied, in which case the effective coherence length may extend to the electronic scale.

2.6 Bose-Einstein Condensation (BEC)

Definition: BEC occurs when particles coalesce into a single quantum state, enabling coherent interactions

.

- **Critical Temperature Estimate:** For a density $n = 1 \times 10^{28}$ m ^{3} of deuterons (mass $m = 3.344 \times 10^{-27}$ kg):

$$T_c = \frac{2\pi\hbar^2}{m k_B} \left(\frac{n}{\zeta(3/2)} \right)^{2/3}$$

  Using:
  – $\hbar = 1.055 \times 10^{-34}$ J · s
  – $m = 3.344 \times 10^{-27}$ kg (deuteron mass)

– $k_B = 1.381 \times 10^{-23}$ J/K
– $\zeta(3/2) \approx 2.612$

$$T_c = \frac{2\pi \times (1.055 \times 10^{-34})^2}{3.344 \times 10^{-27} \times 1.381 \times 10^{-23}} \times \left(\frac{10^{28}}{2.612}\right)^{2/3} \approx 7.9 \text{ K}$$

Critical Assessment and Non-Equilibrium Extension:

The equilibrium BEC temperature of 7.9 K is far below LENR operating conditions (300-800 K), making equilibrium BEC impossible. However, non-equilibrium driven-dissipative mechanisms may achieve the required phase-space density through three synergistic effects:

1. **Effective Mass Renormalization (Polaron Dressing):** Deuterons strongly coupled to the phonon cloud form polaronic quasiparticles with effective mass $m^* < m_D$. For strong electron-phonon coupling in PdD systems, mass reduction factors of 3-5x are plausible, raising T_c to 24-40 K. *(Model assumption: the 3-5x factor is an order-of-magnitude estimate, not a measured value; it is treated as an uncertain input for the Monte Carlo / sensitivity analysis of Section 5.1 and is not independently cited.)*
2. **Driven Dissipative Cooling (Phonon Standing Wave Mechanism):** Coherent phonon standing waves—whether acoustically driven or arising from lattice resonances—create a spatially periodic potential for interstitial deuterons. Through anharmonic coupling, deuteron kinetic energy is preferentially channeled into high-frequency optical phonon modes that rapidly dissipate into the bulk lattice. This mechanism is analogous to resolved sideband cooling in optomechanics and Sisyphus cooling in atomic physics. Estimated effective temperature reduction: factor of 3-5x, giving $T_{\text{eff}} \sim$ 60-100 K.
3. **Local Density Enhancement at Standing Wave Nodes:** Phonon radiation pressure concentrates deuterons at nodes or antinodes of the standing wave pattern, locally increasing number density by factors of 5-10x. Since $T_c \propto n^{2/3}$, this raises T_c by factors of 2.9-4.6x.

Combined Quantitative Estimate:

$$T_c^{\text{eff}} \approx 7.9 \text{ K} \times 5 \text{ (mass)} \times 4.6 \text{ (density)} \approx 182 \text{ K}$$

$$T_{\text{deuteron}}^{\text{eff}} \approx 300 \text{ K}/5 \text{ (cooling)} \approx 60 \text{ K}$$

Since $T_{\text{deuteron}}^{\text{eff}} < T_c^{\text{eff}}$, BEC becomes formally possible under combined non-equilibrium conditions.

Open Constraints: (a) Phonon-phonon thermalization in metals occurs on ~ps timescales; the cooling mechanism must cycle faster than thermalization re-equilibrates. (b) The direct criterion is phase-space density $n\lambda_{\text{dB}}^3 \geq \zeta(3/2)$ rather than temperature comparison alone. (c) Whether the resulting state is true BEC (spontaneous symmetry breaking) or a coherently driven state (analogous to a laser) affects the predicted nuclear enhancement and must be distinguished experimentally.

2.7 Oppenheimer-Phillips Process

Definition: A mechanism where a deuteron interacts with a nucleus at low energies, resulting in neutron capture and proton emission

16

.

$$d + {}^{A}_{Z}\mathrm{X} \longrightarrow {}^{A+1}_{Z}\mathrm{Y} + p.$$

1. **Mechanism**
   - Neutron from the deuteron is absorbed, forming ${}^{A+1}_{Z}\mathrm{Y}$.
   - The proton is repelled by Coulomb forces and emitted.

2. **Rate Calculation**

$$R_{\text{rate}} = n_d \, n_X \, \langle \sigma_{\text{OP}} v_d \rangle,$$

where σ_{OP} is the Oppenheimer-Phillips (deuteron-stripping) cross-section, n_d and n_X are deuteron and target densities, and $\langle \sigma_{\text{OP}} v_d \rangle$ is the reactivity formed with the deuteron velocity distribution (not a free-neutron flux).

2.8 Quantum Tunneling Enhancements

Standard Tunneling Probability

$$P_{\text{tunnel}} = \exp\left(-\frac{2}{\hbar} \int_a^b \sqrt{2m\big(V(x) - E\big)}\, dx \right).$$

LENR Conditions

- **Reduced Barrier Height:** Electric fields, Rydberg states, and Casimir effects lower $V(x)$.
- **Enhanced Particle Energy:** Localized heating and lattice phonon interactions can further increase effective particle energies.

Overall Enhancement

The reported 10^3–10^5 range is a **bounded phenomenological sensitivity band** implemented in the current LENR_SIM software (independent mechanism multipliers with a hard composition cap), **not** a first-principles prediction of a coupled electron–lattice–nuclear Hamiltonian. Factorization into electron screening ($f_{\text{screening}} \sim 10^{1-2}$), interface electric fields ($f_{\text{field}} \sim 10^{0.5-1}$), lattice coherence ($f_{\text{coherence}} \sim 10^{1-2}$), and phonon coupling ($f_{\text{phonon}} \sim 10^{0-0.5}$) is a bookkeeping device: $f_{\text{total}} = f_{\text{screening}} \times f_{\text{field}} \times f_{\text{coherence}} \times f_{\text{phonon}} \sim 10^3$ to 10^5, subject to

software caps and claim-tier labels.[1] Monte Carlo / Sobol runs in the companion app explore this band; they do not by themselves validate microscopic mechanisms.

2.8.1 Phonon-Nuclear Coupling Strength A critical mechanism for LENR is the coupling between lattice phonons and nuclear degrees of freedom. This coupling enables energy transfer from the lattice to nuclear systems, potentially facilitating reactions that would otherwise be energetically forbidden.

Coupling Constant Definition:

The phonon-nuclear coupling strength can be expressed as:

$$g_{\text{ph-nuc}} = \frac{\partial V_{\text{nuclear}}}{\partial u} \cdot \sqrt{\frac{\hbar}{2M\omega_{\text{ph}}}}$$

where: - V_{nuclear} is the nuclear potential - u is the lattice displacement - M is the nuclear mass - ω_{ph} is the characteristic phonon frequency

Calculation for Deuterons in Palladium:

For deuterons in Pd lattice: - $M = 3.344 \times 10^{-27}$ kg (deuteron mass) - $\omega_{\text{ph}} \approx 10^{13}$ rad/s (optical phonon frequency in Pd-D) - $\partial V/\partial u \approx 1 - 10$ eV/Å (estimated from lattice strain effects)

$$g_{\text{ph-nuc}} \approx (1 - 10) \times \sqrt{\frac{1.055 \times 10^{-34}}{2 \times 3.344 \times 10^{-27} \times 10^{13}}} \approx 0.1 - 1 \text{ eV/Å}$$

Energy Transfer Mechanism:

The phonon-mediated energy transfer rate is given by:

$$\Gamma_{\text{transfer}} = \frac{2\pi}{\hbar} |g_{\text{ph-nuc}}|^2 \rho(\omega) n_{\text{ph}}(\omega)$$

where $\rho(\omega)$ is the phonon density of states and $n_{\text{ph}}(\omega) = (e^{\hbar\omega/k_B T} - 1)^{-1}$ is the Bose-Einstein distribution.

Multi-Phonon Processes:

For reactions requiring energy $\Delta E \sim 1$ MeV, coherent multi-phonon processes are necessary:

$$P_{\text{multi-phonon}} \propto \left(\frac{g_{\text{ph-nuc}}}{\hbar\omega_{\text{ph}}} \right)^N \quad \text{where } N = \frac{\Delta E}{\hbar\omega_{\text{ph}}}$$

For $\Delta E \sim 1$ MeV and $\hbar\omega_{\text{ph}} \sim 50$ meV, $N \sim 2 \times 10^7$ phonons must act coherently. This highlights the importance of quantum coherence in LENR systems, as discussed in Section 2.5.1.

Critical Assessment: The probability of $N \sim 2 \times 10^7$ phonons acting coherently through independent coupling scales as $(g_{\text{ph-nuc}}/\hbar\omega_{\text{ph}})^N$, which for any coupling ratio less than unity

[1] Software composition is **not** this four-factor product. The theory pipeline uses grouped-max then a 10^5 cap (`EQ-COMP-01`, `EQ-COMP-02`); multiplying that total by the bubble-lattice peak (or the legacy geometric-mean path) is an illegal product (`EQ-COMP-03`). See `LENR_SIM_APP_v2/docs/v0.3.6-beta/THEORY_COMPOSITION_CONTRACT.md`.

yields a probability indistinguishable from zero (e.g., $0.99^{2\times10^7} \sim 10^{-87,000}$). This effectively rules out the naive multi-phonon pathway for bridging the MeV energy gap.

However, several alternative mechanisms may circumvent this limitation: 1. **Stimulated phonon cascades:** Analogous to stimulated emission in lasers, where each phonon absorption event increases the probability of the next, creating a cooperative avalanche rather than independent events. 2. **Coherent energy transfer via off-shell intermediates:** Hagelstein's model proposes that energy transfer between lattice and nuclear degrees of freedom proceeds through virtual (off-shell) intermediate states, analogous to Raman scattering, where the full ΔE need not be assembled simultaneously. 3. **Phonon-polariton condensation:** If the phonon system achieves a coherent state (analogous to a phonon laser or Bose-Einstein condensate of phonon-polaritons), the energy transfer becomes a collective quantum process with fundamentally different statistics than independent phonon coupling.

The resolution of the MeV energy gap through lattice-nuclear coupling remains the central open theoretical challenge in LENR. See Section 6.1 for further discussion.

Implications:

1. The coupling strength determines the rate of energy exchange between lattice and nuclear degrees of freedom
2. High phonon occupation numbers (achieved at elevated temperatures or through external driving) enhance energy transfer
3. Coherent phonon states can dramatically increase the effective coupling by enabling cooperative multi-phonon processes

---

3. Computational Methods

> **Implementation note (public beta, v0.3.7-beta):** Locked live claim tiers — **Beta:** WKB/Gamow tunneling (`ENG-WKB-01`); **Approximate:** Rayleigh–Plesset / Keller–Miksis bubble dynamics with KM auto-select when peak Mach $\geq$ 0.3 (`ENG-RP-KM-01`). The live VPS runs the in-repo `physics_engine` pipeline via `/api/physics/*` and `/api/unified/*`. Outputs carry provenance (`physics_engine` vs schematic fallback). **Aspirational (NOT live):** COMSOL-class FEM, self-consistent Poisson–Schrödinger, and coupled continuum solvers (`ENG-ASPIRATIONAL-FEM-01`) are research methods only — never public-beta numerics. Lean `PoissonSchrodinger` / IR `EQ-PS-*` encode residual, domain, and non-deployment honesty only; they do **not** prove LENR or close any `EQ-OPEN-*`.

3.1 Poisson–Schrödinger Coupled Dynamics

Honesty label: Continuum FEM / self-consistent Poisson–Schrödinger below is an **aspirational modeling roadmap** (`ENG-ASPIRATIONAL-FEM-01`). It is **not** executed on the public beta, **not** a COMSOL production path in this release, and **not** a discharge of any `EQ-OPEN-*`.

Formalization locks parent continuum DEF `EQ-PS-01`, discrete residual algebra, permittivity domain, explicit tolerance axiom, and non-deployment flags (`EQ-PS-POISSON-01`, `EQ-PS-SCHROD-01`, `EQ-PS-COUPLE-01`, `EQ-PS-DOM-01`, `EQ-PS-TOL-01`, `EQ-PS-NONDEPLOY-01`).

1. **Poisson's Equation** (parent continuum DEF `EQ-PS-01`):

$$\nabla^2\phi = -\frac{\rho}{\epsilon_0} \quad \text{or more generally} \quad \nabla\cdot\big(\varepsilon\nabla\phi\big) = -\rho,$$

   where ϕ is the electric potential, ρ is the charge density, and ε is a (possibly spatially varying) permittivity. Domain: permittivity must be positive wherever the formula is claimed (`EQ-PS-DOM-01`).

2. **Schrödinger's Equation:**

$$-\frac{\hbar^2}{2m}\nabla^2\psi + V(\mathbf{r})\psi = E\psi,$$

   which governs the quantum state ψ under potential $V(\mathbf{r})$. Here E is the energy eigenvalue (`E_eigen`), not the electric field (`E_field`).

3. **Self-consistent coupling (roadmap):** iterate Poisson $\to$ update $V \leftarrow V_{\text{ext}} - e\phi \to$ Schrödinger $\to$ update ρ from $|\psi|^2$ until residual $r = \max(r_{\text{P}}, r_{\text{S}})$ vanishes in the discrete bookkeeping sense (`EQ-PS-COUPLE-01`), or $r \le \tau$ under an **explicit** tolerance axiom (`EQ-PS-TOL-01` — never silently promoted to exact $r = 0$). Continuum existence / FEM convergence are **not** theorems of this pilot.

4. **Simulation Details (research targets, not VPS execution)**
   - **Software target:** COMSOL Multiphysics and custom Python scripts (NumPy, SciPy) for finite-element and finite-difference methods — aspirational.
   - **Grid Size:** $\sim 0.1\,\text{nm}$.
   - **Time Steps:** $\sim 1\,\text{fs}$ (time-dependent extensions).
   - **Illustrative expectations:** Electric fields in interfacial regions can exceed $10^8\,\text{V/m}$; wavefunction solutions can show elevated electron density near protons. Any link to neutron-formation pathways remains claim-tiered and speculative — **not** nuclear-rate proof.

3.2 Bubble Collapse Dynamics

Honesty label: Cavitation–collapse in this framework is **modelled enhancement / contested hydro**. Rayleigh–Plesset / Keller–Miksis trajectories, surface-proximity factors, and collapse UQ percentiles are companion-software sensitivity explorations under claim-tier governance. They are **not** laboratory proof of nuclear rates, **not** a discharge of any `EQ-OPEN-*`, and **not** a physical-necessity theorem. Lean `LenrFormal.CollapseBounds` encodes only clamp / bound / UQ / RP-domain / hard-core CR-cap bookkeeping (`EQ-BBL-LIM-01`,

EQ-BBL-BOUND-01, EQ-BBL-UQ-01, EQ-BBL-RP-DOM-01, EQ-BBL-CR-CAP-01)—including the pointwise clamp min(raw, bound)—not cavitation-dynamics theorems that prove LENR. WH-ENP-01 / OpenObligations prove **non-closure** implications under stated deficits—not affirmative LENR (docs/FORMALIZATION/FORMAL_PROOF_STATUS.md).

Context: During electrolysis, microbubbles can form and then collapse, concentrating energy

. Pyyhtiä et al. (2026) (pyyhtia2026) provide OA experimental support that HER-driven nanobubble cavitation damages CR-39 and confounds SPAWAR-class pit=nuclear detector interpretations (pits without Pd and/or D O under suitable geometry; HIFU control; shallow polymer pits vs deep tracks)—**not** LENR-rate proof; EQ-OPEN-RAD-01 stays OPEN. Ledger IDs: ENG-BUBBLE-FUSION-01, ENG-RP-KM-01 (claim tier **approximate**; do not treat collapse numerics as nuclear-rate proof). Collapse UQ in the companion engine (collapse_uq.py, Sobol) reports modelled percentiles; campaign grids that stress cavitation/collapse cells remain model output only (see §3.8).

1. **Modified Rayleigh–Plesset Equation** (incompressible / low-Mach baseline):

$$R\,\ddot{R} + \frac{3}{2}\dot{R}^2 = \frac{1}{\rho}\left(P_{\text{bubble}} - P_\infty - \frac{2\sigma_{\text{surf}}}{R} - \frac{4\mu\,\dot{R}}{R} + \frac{\epsilon_0 E^2}{2}\right),$$

   where R is bubble radius, ρ fluid density, σ_{surf} surface tension, μ viscosity, and E external electric field.

   Note: The electric pressure term $\epsilon_0 E^2/2 \approx 4.4 \times 10^4$ Pa at $E = 10^8$ V/m is negligible compared to collapse pressures of ~GPa. This term primarily influences bubble initiation and early-stage dynamics rather than peak collapse conditions. Rayleigh–Plesset is **domain-limited** when peak Mach numbers are not $\ll 1$.
2. **Keller–Miksis compressibility correction (companion software, v0.3.7-beta):** When liquid compressibility / acoustic radiation matters, the Unified and Bubble Fusion paths may use dynamics_mode=auto and select Keller–Miksis when peak Mach ≥ 0.3. Explicit overrides (legacy / rayleigh_plesset / keller_miksis) remain available. Making KM the unconditional default, real-gas EOS, and FEM remain deferred (DEFERRED_MULTIPHYSICS_FIDELITY.md). LaTeX stub: docs/v0.3.7-beta/latex/eng_rp_km_01.tex.
3. **Provenance honesty:** UI and export envelopes must distinguish provenance=physics_engine ODE trajectories from schematic / local fallbacks. Schematic output must never be presented as validated engine numerics.
4. **Simulation Outcomes:**
   - **Temperature Peaks:** Localized temperatures can exceed 10 K in extreme regimes. *Note:* Peak temperatures depend strongly on the collapse regime. Single-bubble sonoluminescence in water typically achieves ~10 K, while the 10 K figure applies to extreme multi-bubble collapse or laser-driven cavitation conditions. The relevant regime for LENR electrolysis systems should be specified when applying these estimates.

- **Pressure Increases:** Collapsing bubbles can produce pressures of multiple GPa. These collapse conditions may alter interfacial transport and material microstructure, but do not by themselves establish an appreciable nuclear reaction rate.

3.3 Heterogeneous Nucleation at Surfaces

Bubble formation in LENR systems occurs predominantly at lattice surfaces (heterogeneous nucleation), where cavities, defects, and wettability reduce the energy barrier compared to homogeneous nucleation in the bulk.

1. **Contact angle factor:** For a surface with contact angle , the heterogeneous nucleation barrier is reduced by f( ) = (2 + cos)(1 − cos)²/4. For = 90°, f = 0.5; for complete wetting → 0°, f → 0.
2. **Critical radius (Young–Laplace):** r* = 2 /ΔP, where is liquid–vapor surface tension and ΔP is the pressure difference between vapor and liquid. At surfaces, geometry and contact angle modify the effective critical radius.
3. **Gibbs free energy barrier:** ΔG*_het = ΔG*_hom × f( ), with ΔG*_hom = (16 ³)/(3 ΔP²). Lower f( ) at hydrophobic or cavity sites promotes nucleation.
4. **Waiting time and active site probability:** Nucleation rate depends on exp(−ΔG*/(kT)); statistical waiting times and site activation probabilities follow from classical nucleation theory and supersaturation.

3.4 Bubble–Surface Coupling

The physical coupling between a cavitation bubble and the lattice surface determines equilibrium standoff, attachment/detachment, and asymmetric collapse.

1. **Equilibrium distance:** Balance of Casimir (vacuum energy), van der Waals, and thermal forces sets the equilibrium separation between bubble and surface; standoff ratio = h/R (gap/radius) affects collapse symmetry.
2. **Van der Waals (sphere–plate):** Derjaguin approximation gives the force between a bubble and a flat surface; Hamaker constants for water–metal systems determine attraction at nanoscale gaps.
3. **Detachment radius:** Fritz-style relations describe bubble departure size when buoyancy and dynamic forces overcome attachment (surface tension, contact line).
4. **Microjet formation:** Asymmetric collapse near a rigid boundary produces a high-speed jet toward the surface; jet velocity and impact pressure scale with standoff and drive localized energy deposition. Any link from microjets to nuclear rates remains **modelled enhancement / contested hydro**, not an empirical LENR claim.
5. **Surface proximity enhancement:** For < 2, collapse pressure and temperature are amplified relative to spherical collapse in the hydro models; integrated software may apply

surface-proximity factors inside claim-tiered rate **scenarios**. Those factors do not close `EQ-OPEN-RATE-01` or raise empirical claim tiers.

3.5 Thermal and Diffusion Boundary Layers

Thin layers adjacent to the lattice control where nucleation is favored and how heat and mass are transported.

1. **Thermal boundary layer:** Thickness _th from Nusselt-number correlations (e.g., Nu Re^a Pr^b); determines the superheat profile and the region where liquid is hot enough to nucleate bubbles.
2. **Diffusion boundary layer:** Thickness _D scales with Lewis number (Le = /D); concentration gradients of dissolved gas (e.g., D) set supersaturation and nucleation driving force.
3. **Nucleation zone thickness:** The overlap of superheated/supersaturated regions defines where heterogeneous nucleation is most likely; material properties (Pd, Ni, D O) and flow conditions set typical scales (µm to tens of µm).
4. **Temperature and concentration profiles:** Erf-based or similar profiles describe T(y) and C(y) from surface to bulk; used in integrated surface-reaction and nucleation models.

3.6 Acoustic and Ultrasonic Forcing

External or flow-driven acoustic fields alter bubble dynamics and nucleation in electrolytic and sonochemical LENR configurations.

1. **Minnaert frequency:** Natural bubble resonance f (1/R)√(3 P/); driving near f amplifies radial oscillations and can promote rectified diffusion and collapse severity.
2. **Primary Bjerknes force:** Acoustic radiation force on a bubble in a pressure gradient drives it toward pressure antinodes (or nodes, depending on compressibility), concentrating bubbles in specific regions.
3. **Secondary Bjerknes force:** Bubble–bubble interaction force (e.g., attraction for similar phase); affects clustering and coalescence.
4. **Acoustic streaming:** Rayleigh-type streaming near boundaries creates steady circulation; velocity scales with (p_a²/(c²))(L/ _v), influencing boundary layer and nucleation zone.
5. **Standing wave positions:** Node/antinode locations set where pressure and velocity amplitudes are maximal; used to place electrodes or nucleation sites.
6. **Acoustic nucleation enhancement:** Pressure oscillations lower the effective nucleation barrier (e.g., reduction of ΔG^* in $\exp(-\Delta G^*/(kT))$), increasing nucleation rate.
7. **Rectified diffusion:** Asymmetric gas flux over the oscillation cycle causes net gas uptake by bubbles, leading to growth and eventual collapse; important for sustained cavitation.

3.7 Non-Equilibrium Thermodynamics

LENR systems operate far from thermodynamic equilibrium, requiring a proper treatment of energy flow, entropy production, and dissipative structures. This section provides the thermodynamic foundation for understanding energy transport in active LENR systems.

3.7.1 Energy Continuity Equation The fundamental energy balance in a LENR system is governed by the continuity equation:

$$\frac{\partial \rho_E}{\partial t} + \nabla \cdot \mathbf{J}_E = \sigma_{\text{source}}$$

where: - ρ_E is the local energy density [J/m³] - $\mathbf{J}_E$ is the energy current density [W/m²] - σ_{source} is the volumetric source term from nuclear reactions [W/m³]

For LENR systems, the source term includes:

$$\sigma_{\text{source}} = R_{\text{nuc}} \cdot Q_{\text{reaction}} - \sigma_{\text{rad}} - \sigma_{\text{chem}}$$

where R_{nuc} is the nuclear reaction rate, Q_{reaction} is the energy per reaction, σ_{rad} accounts for radiative losses, and σ_{chem} represents chemical energy contributions.

3.7.2 Entropy Production The local entropy production rate in LENR systems is:

$$\dot{s} = \frac{1}{T}\left[\mathbf{J}_E \cdot \nabla\left(\frac{1}{T}\right) + \sum_i \mathbf{J}_i \cdot \nabla\left(\frac{\mu_i}{T}\right) + \sigma_{\text{source}}\left(1 - \frac{T_0}{T}\right)\right]$$

where: - T is the local temperature - $\mathbf{J}_i$ are species fluxes - μ_i are chemical potentials - T_0 is a reference temperature

The second law requires $\dot{s} \geq 0$, constraining the allowed energy flow patterns.

3.7.3 Heat Transport Equations For practical LENR systems, heat transport follows:

$$\rho c_p \frac{\partial T}{\partial t} = \nabla \cdot (\kappa \nabla T) + \sigma_{\text{source}}$$

where: - ρ is the material density - c_p is the specific heat capacity - κ is the thermal conductivity

Boundary Conditions for Calorimetry:

At the electrode-electrolyte interface:

$$-\kappa \left.\frac{\partial T}{\partial n}\right|_{\text{interface}} = h(T_{\text{surface}} - T_{\text{bulk}}) + \epsilon\sigma_{\text{SB}}(T_{\text{surface}}^4 - T_{\text{ambient}}^4)$$

where h is the convective heat transfer coefficient, ϵ is emissivity, and σ_{SB} is the Stefan-Boltzmann constant.

3.7.4 Connection to Experimental Measurements The excess power measured in calorimetry experiments is:

$$P_{\text{excess}} = \int_V \sigma_{\text{source}}\, dV = P_{\text{out}} - P_{\text{in}} - P_{\text{chem}}$$

Time-resolved measurements can be analyzed using:

$$P_{\text{excess}}(t) = mc_p \frac{dT}{dt} + UA(T - T_{\text{ambient}}) - P_{\text{electrical}}$$

where UA is the overall heat transfer coefficient-area product.

Implications:

1. Non-equilibrium thermodynamics provides the framework for interpreting calorimetric data
2. Entropy production constraints help identify physically plausible energy flow patterns
3. Transient analysis can distinguish between nuclear and chemical heat sources based on temporal signatures

---

3.8 Companion-software simulation evidence (v0.3.7-beta)

Pinned, version-controlled model runs—not laboratory experiments—anchor the quantitative claims in this revision. Authoritative evidence lives under `LENR_SIM_APP_v2/datasets/reviewer_packs/` and the paper multi-format pack provenance file (`PROVENANCE.md`).

1. **Reviewer packs:** `smoke/` (CI smoke) and `q_campaign_demo/` (larger campaign: 150 experiments / 132 theory runs in the recorded manifest). Schema `lenr-reviewer-pack-v1`. Disclaimer on every pack: model output only; additional runs do **not** raise empirical claim tiers or close `EQ-OPEN-*`.
2. **Cavitation / collapse campaign pack:** `LENR_SIM_APP_v2/datasets/reviewer_packs/cavitation_collapse_campaign/` (Phase A; on `dev` via #299). Model-only bubble-lattice host×loading×drive×dynamics_mode cells with collapse UQ; complements engine modules (`collapse_uq.py`, RP/KM paths). Contested Taleyarkhan sonofusion literature is an anchor, not laboratory proof. Pyyhtiä et al. 2026 (`pyyhtia2026`) sits alongside as an electrolytic CR-39 cavitation / detector-confounder peer—not a nuclear-rate proof.
3. **Detector / weak screen campaign pack:** `LENR_SIM_APP_v2/datasets/reviewer_packs/detector_weak_screen/` (WS7; on `dev` via #331). Model-only cells for detector/weak/screening inventory; dual `score`/`final_score` honesty preserved; does **not** close `EQ-OPEN-*` or set `lenr_proved`.
4. **Campaign + UQ / avenue ranking:** Rate and collapse UQ feed decision-support avenue ranking at `LENR_SIM_APP_v2/docs/v0.3.9-beta/AVENUE_RANKING.md` (Phase B; on `dev` via #301). Ranking prioritizes which lab avenues look informative next; it does **not** set `lenr_proved` or close OPEN ids. Companion Lean `Decision` / IR `EQ-UQ-BF-01` ExpHom identities are **algebraic honesty floors only**—not empirical Bayes factors or laboratory pos-

teriors.

5. **Theory weight surfaces:** `LENR_SIM_APP_v2/docs/v0.3.9-beta/THEORY_WEIGHT_SURFACES.md` (WS7; on `dev` via #331) inventories `CLAIM_TIER_MULTIPLIER`, oracle priors, and avenue blends as **Approximate / Aspirational** governance weights—not measured lab posteriors.
6. **Campaign invariants:** `formalization/predictions/campaign_invariants.json` and pack-embedded invariants enforce `open_questions.lenr_proved = false`, the five OPEN IDs, two-body $\log_{10} N_{\text{events}} \approx -40$, and `contract_version = v0.3.7-beta-science-gate-1`.
7. **Export envelope:** `/api/unified/export` uses `lenr-run-export-v1` with claim-tier / provenance fields required by the data contract.
8. **Paper benchmarks:** `backend/tests/test_paper_benchmarks/` locks selected theory numeric/regime targets in CI.
9. **Lean clamp / non-closure (not closure):** `CollapseBounds` (`EQ-BBL-LIM/BOUND/UQ/RP-DOM/CR-CAP-01`), `WeakInteraction` (`EQ-EC-*`; EC closure), `Overlap` (`EQ-OVERLAP-01/02`; ensemble rate), `PoissonSchrodinger` (`EQ-PS-01`/`EQ-PS-TOL-01`; aspirational FEM), and WH / OpenObligations non-closure theorems constrain bookkeeping; they do not discharge `EQ-OPEN-*`. Draft closure/falsification criteria (still OPEN) live at `LENR_SIM_APP_v2/docs/v0.3.9-beta/EQ_OPEN_CLOSURE_RUNWAY.md` (Phase F; on `dev` via #297).
10. **Neutron-detector honesty (RAD/RATE):** Companion `neutron_rate_bound` carries honesty class `detector_ul_not_proof`. Cd-ratio / moderator-null / soft-X ¬hard- PREDs (PRED-08/09/10) and contested Parkhomov curated rows are falsifiers / literature anchors—they do **not** close `EQ-OPEN-RAD-01` or `EQ-OPEN-RATE-01`, and /anti- detector-gas SPEC rows do not discharge `EQ-OPEN-EC-01`.
11. **Non-claims:** Larger N in a campaign does not convert speculative mechanism labels into empirical validation. Maturity axes (`model_fidelity`, `empirical_support`, `claim_tier`) remain independent (`PHYSICS_MATURITY_LEDGER.md`).

---

4. Experimental Context and Reported Correlations

4.1 Electrolytic Reactors

1. **Setup:**
   - Palladium electrodes ($> 99.99\%$ purity).
   - Heavy water (D_2O) electrolytes with lithium salts.
   - Current densities between 0.1–1 $\mathrm{A/cm^2}$.
2. **Measurements:**
   - **Calorimetry:** Input power (voltage $\times$ current) vs. output power via isoperibolic or flow calorimeters.

- **Mass Spectrometry:** Detection of helium-4, tritium, or other isotopes in off-gas.

- **Neutron Counters:** Helium-3 or BF_3 detectors above background sensitivity. Detector upper limits in the companion software are honesty-classed `detector_ul_not_proof`; Cd-ratio / moderator-null / soft-X ¬hard- PREDs (PRED-08/09/10) are branching falsifiers, not closures of `EQ-OPEN-RAD-01` / `EQ-OPEN-RATE-01`.

3. **Reported outcomes (disputed; claim-tier: empirical correlation):**
   - Excess heat relative to electrical input has been reported in some electrolytic campaigns (often cited near tens of percent); replication and calorimetry systematics remain contested.

   - Helium-4 and tritium signals have been reported in a subset of studies; collection efficiency and blank controls vary.

   - Low-level neutron signals are occasionally reported above background; they are not established as routine LENR signatures, and a Cd-wrapped count is **not** neutron proof.

4.2 Gaseous Reactors

1. **Setup:**
   - Nickel powders with high surface area.

   - Hydrogen gas at 1–10 atm.

   - Operating temperatures of 300–800 K.

2. **Measurements:**
   - **Calorimetry:** Monitoring of potential excess heat.

   - **Spectroscopy:** Raman, infrared, or other techniques to detect Rydberg Matter or ultra-dense hydrogen states.

   - **Elemental Analysis:** Post-run detection of transmutation products such as copper and zinc in nickel.

3. **Reported outcomes (disputed; claim-tier: empirical correlation):**
   - **Anomalous Heat:** High-power excess claims (including kW-scale reports) exist in the literature but lack community consensus on calorimetry.

   - **RM Detection:** Spectroscopic signatures have been interpreted as consistent with

Rydberg Matter in some studies; alternative assignments exist.

- **Transmutations:** Post-run elemental shifts on nickel surfaces are reported in some campaigns; contamination and analytical artifacts must be excluded case-by-case.

---

5. Results and Discussion

5.1 Computational–Experimental Correlation

- **Tunneling Enhancements:** Calculations show up to 10^3 to 10^5-fold increases in tunneling probability under combined RM, halo electron, and high-field environments.

- **Neutron Formation:** Scenario neutron-rate bands in the companion software are claim-tiered sensitivity explorations under `detector_ul_not_proof` honesty; they may be *compared* to occasionally reported low-level neutron signals but do **not** validate those signals, close `EQ-OPEN-RAD-01` / `EQ-OPEN-RATE-01`, or establish routine LENR neutron production.

5.1.1 Statistical Mechanics of Reaction Rates A rigorous treatment of LENR reaction rates requires proper statistical mechanics, accounting for thermal distributions, enhancement factors, and threshold conditions.

Reaction Rate Formula:

The nuclear reaction rate per unit volume for two reacting species is:

$$R = \frac{n_1 n_2}{1 + \delta_{12}} \langle \sigma v \rangle_{\text{thermal}} \times f_{\text{enhancement}} \times P_{\text{threshold}}$$

where $\delta_{12} = 1$ for identical reacting species (e.g., D+D self-fusion, giving the familiar $\frac{1}{2}n^2$ form used by `reaction_rate_engine.py::compute_two_body_rate`) and $\delta_{12} = 0$ for distinct species. This is the *thermal self-fusion* rate and is distinct from the Oppenheimer–Phillips capture rate $R_{\text{rate}} = \sigma_{\text{OP}} \Phi_n N$ (§2.7, a flux-driven capture process) and the generic placeholder $R_{\text{rate}} = n \sigma_{\text{nuc}} v$ (§7); see the Glossary (§8) note on R for how these three regimes relate.

where: - n_1, n_2 are the number densities of reacting species [m ^{3}] - $\langle \sigma v \rangle_{\text{thermal}}$ is the thermally-averaged cross-section-velocity product [m^3/s] - $f_{\text{enhancement}}$ is the cumulative enhancement factor from screening, coherence, etc. - $P_{\text{threshold}}$ is the probability of exceeding threshold conditions

Thermal Averaging:

The thermal average is computed over the Maxwell-Boltzmann distribution:

$$\langle \sigma v \rangle = \sqrt{\frac{8}{\pi \mu (k_B T)^3}} \int_0^\infty \sigma(E)\, E\, e^{-E/k_B T}\, dE$$

where μ is the reduced mass and $\sigma(E)$ is the energy-dependent cross-section.

For tunneling-dominated reactions:

$$\sigma(E) \propto \frac{1}{E} \exp\left(-\sqrt{\frac{E_G}{E}}\right)$$

where $E_G = 2\mu c^2(\pi\alpha Z_1 Z_2)^2$ is the Gamow energy (in SI, $E_G = 2\mu\left[\pi Z_1 Z_2 e^2/(4\pi\varepsilon_0\hbar)\right]^2$; for D+D, $E_G \approx 986$ keV).

Enhancement Factor Decomposition:

The total enhancement factor combines multiple mechanisms:

$$f_{\text{enhancement}} = f_{\text{screening}} \times f_{\text{field}} \times f_{\text{coherence}} \times f_{\text{phonon}}$$

Each factor has associated uncertainties that propagate through the calculation.

Error Propagation:

The uncertainty in the reaction rate is:

$$\frac{\delta R}{R} = \sqrt{\left(\frac{\delta n_1}{n_1}\right)^2 + \left(\frac{\delta n_2}{n_2}\right)^2 + \left(\frac{\delta\langle\sigma v\rangle}{\langle\sigma v\rangle}\right)^2 + \left(\frac{\delta f}{f}\right)^2}$$

Statistical Significance Criteria:

For experimental validation, statistical significance is assessed using Poisson statistics:

$$S = \frac{n_{\text{observed}} - n_{\text{background}}}{\sqrt{n_{\text{background}}}}$$

A significance of $S \geq 5$ (5) is typically required for claiming positive detection. For excess heat measurements:

$$S_{\text{heat}} = \frac{Q_{\text{excess}}}{\sigma_Q}$$

where σ_Q includes both statistical and systematic uncertainties.

Monte Carlo Validation:

Reaction rate predictions are validated using Monte Carlo sampling: 1. Sample input parameters from their uncertainty distributions 2. Propagate through the full calculation chain 3. Build probability distributions for output observables 4. Compare with experimental data using likelihood ratios

Honesty on sample size (companion app, 2026-08-26): default Sobol UQ remains **512 samples, opt-in** (API cap 512). Campaign presets that a reviewer can actually run are `literature_grid` (~16–32 designed cells), `all_claims` (~6 unified + one run per registered theory), and optional `diverse` (~237). That corpus is on the order of **hundreds of model-output rows**, not 10^4–10^6. The larger Monte Carlo aspiration remains a sensitivity method, not a completed campaign. See `LENR_SIM_APP_v2/docs/v0.3.6-beta/RESEARCHER_REVIEW_PACK.md`. Extra

runs do not close §5A or prove LENR.

Implications:

1. Proper statistical treatment enables rigorous comparison between theory and experiment
2. Error propagation identifies which parameters dominate the uncertainty budget
3. Statistical significance criteria provide objective standards for experimental claims
4. Monte Carlo methods enable exploration of parameter space and identification of optimal conditions

5.2 Validation of Nuclear Products

- **Isotopic Shifts:** Detection of helium-4, tritium, and heavier isotopes supports multi-step nuclear processes (e.g., Oppenheimer-Phillips).

- **Excess Heat Consistency:** Reported experimental outputs are consistent in magnitude with the hypothesized nuclear energy contributions once purely chemical avenues are discounted; this remains a disputed empirical correlation (claim tier: `empirical_lenr_correlation`), not an established result.

5.2.1 Detailed Energy Accounting For D+D $\rightarrow$ He reactions, the total energy release must be properly accounted for:

$$E_{\text{released}} = E_{\text{binding}}(^4\text{He}) - 2E_{\text{binding}}(D) = 23.8 \text{ MeV}$$

This energy must be distributed among multiple channels:

1. **Lattice Phonons:** $E_{\text{phonon}} = \sum_i \hbar\omega_i n_i$, where ω_i are phonon frequencies and n_i are occupation numbers
2. **Electronic Excitations:** $E_{\text{electronic}} = \sum_k E_k f_k$, where E_k are electronic energy levels and f_k are occupation probabilities
3. **Nuclear Recoil:** $E_{\text{recoil}} = \frac{p^2}{2M}$, where p is the recoil momentum and M is the nuclear mass
4. **Radiation (if not suppressed):** $E_\gamma = \hbar\omega_\gamma$, where ω_γ is the photon frequency

Energy conservation requires:

$$E_{\text{nuclear}} = E_{\text{phonon}} + E_{\text{electronic}} + E_{\text{recoil}} + E_\gamma$$

For LENR, the suppression of high-energy gamma emission suggests:

$$E_\gamma \ll E_{\text{nuclear}}$$

implying that most energy is channeled into lattice and electronic degrees of freedom. This energy distribution mechanism explains the observed excess heat without commensurate high-energy radiation, consistent with the LENR signature of nuclear-scale energy release with minimal ionizing radiation.

5.2.2 Energy-Product Yield Reconciliation A self-consistency check between excess heat and nuclear product yields is essential. For the primary channel D+D $\rightarrow$ He + 23.8 MeV:

- Energy per reaction: $Q = 23.8$ MeV $= 3.81 \times 10^{-12}$ J
- Reactions required for 1 W $\cdot$ hr: $N = 3600$ J$/3.81 \times 10^{-12}$ J $= 9.4 \times 10^{14}$ reactions
- Expected He yield: 9.4×10^{14} atoms per W $\cdot$ hr

Comparison with experimental claims:

Reported He yields in LENR experiments range from 10^{11}–10^{13} atoms/W $\cdot$ hr (Miles 1993, McKubre 2006). This is a factor of 100–10,000 below the theoretical yield if all excess heat originated from D+D $\rightarrow$ He. Several explanations are possible:

1. **Detection efficiency:** He measurement via mass spectrometry has limited collection efficiency, particularly when helium diffuses into the lattice or escapes through the electrolyte.
2. **Multiple reaction channels:** Heat may also arise from channels with different products: D+D $\rightarrow$ T + p (Q = 4.03 MeV), D+D $\rightarrow$ ^{3}He + n (Q = 3.27 MeV), or lattice-mediated transmutations that produce heat without helium.
3. **Incomplete energy release:** If some reaction energy goes into lattice defect creation or radiation that escapes calorimetric detection, the effective heat per He atom would be lower.

The Miles heat-helium correlation—$(2.6 \pm 0.6) \times 10^{11}$ He atoms per second per watt of excess heat—is within a factor of ~2 of the 23.8 MeV/ He prediction (2.63×10^{11} atoms/s/W). If the underlying measurements are accepted, this correlation is *consistent with* a dominant D+D $\rightarrow$ He channel; it remains a contested empirical correlation (helium retention and measurement caveats apply) rather than a validated mechanism. The discrepancy in integrated yields (per W $\cdot$ hr) likely reflects time-varying reaction rates and measurement integration periods rather than a fundamental inconsistency.

Tritium and neutron accounting:

- **Tritium:** Reported yields of 10^5–10^7 atoms/hr represent a branching ratio of $\lesssim 10^{-9}$ relative to the He channel, consistent with the LENR observation that the D+D $\rightarrow$ T+p branch is heavily suppressed relative to the dominant D+D $\rightarrow$ He + phonons pathway.
- **Neutrons:** Detected neutron fluxes are typically $< 10^3$/s, corresponding to a branching ratio of $\lesssim 10^{-11}$. This extreme suppression of the D+D $\rightarrow$ ^{3}He + n channel is one of the defining signatures of LENR and must be explained by the radiation suppression mechanism (Section 6.1).

5.3 Implications for LENR

- **Unified Model:** By merging condensed matter, quantum mechanics, and nuclear physics, the framework offers a self-consistent candidate interpretation of multiple reported LENR observations; it does not by itself establish that those observations are nuclear in origin.

- **Improved Reproducibility:** Standardized protocols with precisely characterized materials and well-monitored conditions are starting to yield more consistent data.

5A. Open Theoretical Challenges and Fundamental Questions

Occurrence under Miles-class heat–helium correlations is empirically anchored in parts of the corpus; the open theoretical work is **how** MeV energy partitions (Q1), **how** prompt is suppressed (Q2), and **when** the rate budget closes under a named mechanism path (Q3). This section formulates each as a quantitative challenge with testable implications. The companion app's **open-questions virtual lab** (`open_questions` on unified / bubble-lattice / theory-pipeline responses) encodes occurrence (`occurrence.anchored`), when (`when.threshold_crossed`, duty cycle, stimulation), and how (`how.mechanism_paths`, FGR parameters) under Oracle A (Lean identities and conditionals) plus Oracle B (mechanism-path odds). Oracle B Bayes-factor algebra (`EQ-UQ-BF-01` ExpHom cocycle in `LenrFormal.Decision`) is an **algebraic honesty floor** under stated inputs—not an empirical posterior—and theory-weight surfaces remain Approximate/Aspirational inventory (`THEORY_WEIGHT_SURFACES.md`). It never prints `LENR_PROVED`. See `docs/ARCHITECTURE/OPEN_QUESTIONS_VIRTUAL_LAB_2026-08-22.md`, `docs/ARCHITECTURE/MANY_BODY_FGR_MODEL.md`, and `docs/REVIEW_2026/OPEN_QUESTION_CORPUS_GAPS.md`.

Working hypothesis WH-ENP-01 (claim tier `speculative_exploratory`; Lean `LenrFormal.WorkingHypothesis`). Electron–neutron–proton coupling in a deuteron cluster is treated as one object: a Hagelstein coherent virtual-phonon cascade on Takahashi TSC 4D geometry, with a derived harmonic $\Gamma_{\mathrm{ph}}/\Gamma_\gamma$ bound and the same-cluster FGR versus $R_{\mathrm{obs}} = P_{\mathrm{W}}/(QV)$. Independent multi-phonon g^N remains ruled out (`EQ-NAIVE-GN-01`). Bulk Heisenberg–Euler vacuum polarisation is a negative control (`EQ-QED-HE-NEG-01`: lab-field factor $\equiv 1$). Dicke N^2 is **not** the engine rate law (`EQ-COLL-LIN-01`); the paper's $\sim 6.6\,\mathrm{nm}$ domain is larger than the phonon cooperation radius $\lambda/2\pi$, so N^2 is not licensed (`EQ-WH-DICKE-01`). Two on-shell alphas remove a *primary* 24 MeV γ kinematically (`EQ-TSC-CH-01`) but do not close the MeV-gap (`EQ-TSC-STOP-01`, `EQ-WH-STOP-01`). Lean proves conservation and **non-closure** (`EQ-WH-PART-01`, `EQ-WH-BOUND-01`, `EQ-WH-NONCLOSE-01`). `EQ-OPEN-MEV-01`, `EQ-OPEN-RAD-01`, and `EQ-OPEN-RATE-01` stay OPEN.

WH-ENP-01 v2 derived bounds (same claim tier; Lean `LenrFormal.WorkingHypothesisBounds`; runtime blocks `q2.franck_condon`, `q1.coupling_scan`, `dicke.n_within_radius`, `q3.closure_demand`). Four quantitative bounds sharpen the remainder without discharging it. (i) A displaced-oscillator (Huang–Rhys, linear-coupling) Franck–Condon tail—a labelled model assumption—at phonon order $N = Q/hf \approx 5.75 \times 10^8$ ($f = 10^{13}\,\mathrm{Hz}$, $hf \approx 41.4\,\mathrm{meV}$) gives $\log_{10} P_N \le -7 \times 10^5 \ll -11$ for every Huang–Rhys factor $S \le 10^4$; the smallest S that reaches the 10^{-11} target is $S_{\min} \approx 0.9998\,N$, i.e. a lattice relaxation energy $S_{\min} hf \approx 23.79\,\mathrm{MeV} \approx Q$ (`EQ-WH-FC-01`). (ii) The deuteron count inside the phonon cooperation sphere is $n_D V_{\mathrm{dicke}} \approx 0.033 < 1$ ($n_D = 6.8 \times 10^{28}\,\mathrm{m}^{-3}$, $v_s = 3070\,\mathrm{m/s}$; model assumptions), so $N_{\mathrm{coh}}^{\max} = 0$ and Dicke N^2 is not licensed for even one coherent pair (`EQ-WH-DICKE-N-01`). (iii) A conservation-constrained coupling scan with first-order fraction $\min(1, n_{\mathrm{coh}} g^2)$ (model assumption) leaves the cited row ($g = 10^{-4}$, $n_{\mathrm{coh}} = 0$) at a remainder of 23.8 MeV; rows with $n_{\mathrm{coh}} > 0$ are unlicensed scenarios (`EQ-WH-PART-02`). (iv) For 1 W in a nominal $1\,\mathrm{cm}^3$,

$R_{\text{obs}} \approx 2.6 \times 10^{17}\,\text{m}^{-3}\text{s}^{-1}$ demands $P_{\text{occ}} f_{\text{nuc}} C_{\text{lin}}/\tau \approx 5.8 \times 10^{26}\,\text{s}^{-1}$ at the assumed 25 pm trap, above the engine cap product $1 \cdot 1 \cdot 10^{6}/10^{-20}\,\text{s} = 10^{26}\,\text{s}^{-1}$ (`EQ-WH-CAP-01`). Lean composes these into $\neg Q_1 \wedge \neg Q_2 \wedge \neg Q_3$ (`EQ-WH-NONCLOSE-02`); `lenr_proved` stays false.

5A.1 The MeV Energy Gap Problem (Q1)

The central theoretical challenge in LENR: how does energy from meV-scale lattice phonons ($\hbar\omega_{\text{ph}} \sim 50$ meV) bridge to MeV-scale nuclear processes ($\Delta E \sim$ 1-24 MeV)?

Quantitative assessment of proposed pathways:

1. **Direct multi-phonon coupling (ruled out):** As shown in Section 2.8.1, the probability of $N \sim 2 \times 10^7$ independently coupled phonons acting coherently is $\sim 10^{-87{,}000}$. This pathway is definitively excluded for independent coupling.
2. **Hagelstein coherent energy transfer:** Hagelstein and Chaudhary (2015) propose that lattice-nuclear energy exchange proceeds through virtual (off-shell) intermediate states, analogous to stimulated Raman scattering at nuclear energy scales. In this model, the full ΔE need not be assembled simultaneously; instead, energy is transferred coherently through a chain of virtual phonon-nuclear transitions. The predicted coupling matrix element scales as $|V_{\text{eff}}|^2 \propto n_{\text{ph}}^N/N!$ for stimulated processes, which is dramatically larger than the independent coupling estimate. *Status:* A complete QFT derivation remains outstanding, but the model is internally consistent and makes testable predictions about the relationship between excess heat and phonon spectrum.
3. **Superradiant phonon cascades:** By analogy with Dicke superradiance in quantum optics, if N_{atoms} deuterons couple coherently to a phonon field, the collective emission rate scales as N_{atoms}^2 rather than N_{atoms}. For $N \sim 10^4$ atoms in a coherent domain (~6.6 nm), this gives an enhancement of $\sim 10^8$ over incoherent processes. *Status:* Theoretically promising but requires explicit calculation of the collective coupling strength for nuclear transitions.
4. **Non-equilibrium BEC pathway:** As proposed in Section 2.6, if the deuteron subsystem reaches effective BEC conditions through phonon standing wave cooling, the entire condensate acts as a single quantum entity. Nuclear transitions of the condensate involve collective wavefunction overlap, potentially circumventing the single-pair tunneling bottleneck entirely. *Status:* The quantitative estimate in Section 2.6 shows BEC is formally possible ($T_{\text{eff}} < T_c^{\text{eff}}$), but the nuclear transition rate of the condensate has not been calculated.

Conclusion on Q1: The MeV energy gap is not bridged. WH-ENP-01 ranks Hagelstein+TSC above ruled-out g^N and above bulk Schwinger/HE, but the derived energy remainder remains 23.8 MeV until a constrained phonon-nuclear matrix element is supplied (`EQ-WH-PART-01`). TSC two-alpha kinematics and an order-of-magnitude Pd CSDA range for 23.8 MeV α are bookkeeping, not closure. Dicke N^2 is not licensed at the paper domain, and the counting bound is stricter: fewer than one deuteron (≈ 0.033, model-assumption density and sound speed) sits inside the $\lambda/2\pi \approx 49$ pm cooperation sphere, so no coherent pair is licensed (`EQ-WH-DICKE-N-01`). The quantitative form of the gap for linear phonon coupling is the Franck–Condon requirement $S_{\min} \approx N \approx 5.75 \times 10^8$: the lattice relaxation energy would have to equal essentially the full Q (`EQ-WH-FC-01`, displaced-oscillator model assumption). The conservation-

constrained coupling scan keeps the cited remainder at 23.8 MeV; only the unlicensed corner $g = 0.1,\ n_{\mathrm{coh}} = 100$ would zero it (`EQ-WH-PART-02`). The critical next step remains an explicit, experimentally constrained coupling matrix element.

5A.2 Radiation Suppression Mechanism (Q2)

If D+D $\rightarrow$ He releases 23.8 MeV, standard nuclear physics predicts a 23.8 MeV gamma ray. LENR experiments consistently show excess heat with minimal gamma radiation (upper bounds typically $< 10^{-11}$ of what conventional D+D fusion would produce).

Analysis of proposed suppression mechanisms:

1. **Internal conversion to phonons:** The 23.8 MeV must fragment into $\sim 5 \times 10^8$ phonons (at 50 meV each). A single nuclear transition fragmenting into 10^8 final-state quanta faces an enormous phase-space suppression. However, if the nuclear transition occurs within a coherent lattice domain, the final state is a collective lattice excitation rather than individual phonons. The branching ratio depends critically on the density of states at the nuclear transition energy, which is dramatically enhanced in a crystalline environment due to Umklapp processes and defect-mediated momentum transfer.
2. **Lattice-assisted momentum conservation:** In free space, D+D $\rightarrow$ He + is the only two-body channel that conserves both energy and momentum. In a crystal, the lattice can absorb arbitrary momentum (in units of reciprocal lattice vectors), opening the D+D $\rightarrow$ He + phonons channel. The branching ratio scales with the squared matrix element for lattice coupling relative to radiative coupling: $\Gamma_{\mathrm{phonon}}/\Gamma_\gamma \propto (g_{\mathrm{lattice}}/\alpha)^2 \times \rho_{\mathrm{lattice}}(E)/\rho_\gamma(E)$, where ρ are the respective densities of final states.
3. **Quantitative constraint:** For the observed $\Gamma_{\mathrm{phonon}}/\Gamma_\gamma > 10^{11}$, the lattice coupling must be at least $\sim 10^{5.5} \times \alpha \approx 10^{3.2}$ times the electromagnetic coupling, or the lattice density of states must exceed the photon density of states by a comparable factor, or some combination thereof. Whether these conditions are achievable in real PdD lattices remains an open calculation.

Conclusion on Q2: A harmonic first-principles bound gives $\Gamma_{\mathrm{ph}}/\Gamma_\gamma = 0$ because the Pd Debye cutoff ($\sim$ 24 meV) lies far below 23.8 MeV (`EQ-WH-BOUND-01`). The 10^{11} lattice target remains an open target; assumed branching fractions stay scenario input. Beyond the harmonic cutoff, a displaced-oscillator Franck–Condon tail (linear coupling; model assumption) at $N \approx 5.75 \times 10^8$ gives $\log_{10} P_N$ between -4.8×10^9 ($S = 1$) and -2.5×10^9 ($S = 10^4$), with the Lean-certified conservative cap $\log_{10} P_N \leq -7 \times 10^5 \ll -11$ (`EQ-WH-FC-01`); reaching 10^{-11} needs $S_{\min} \approx 0.9998\, N$, i.e. relaxation energy $\approx Q$. Anharmonic, Umklapp, and Hagelstein lossy spin-boson boosts remain named hypotheses and are not inserted into the derived bound. Residual 2.45 MeV neutron thermalization lengths in H-rich liquid/gas/solid/plasma media are tabulated as detector-UL bookkeeping (`detector_ul_not_proof`) and do not partition the ^{4}He Q-value. Cd-ratio / moderator-null / soft-X ¬hard- PREDs (PRED-08/09/10) are branching falsifiers for neutron-interpretation claims—they do **not** close `EQ-OPEN-RAD-01`. Lattice Umklapp/defect DOS at nuclear energy is still missing.

5A.3 Statistical Feasibility and Rate Closure (Q3)

Given the enhancement factors derived in Section 2.8 ($f_{\text{total}} \sim 10^3$-$10^5$) applied to bare D+D tunneling probabilities at thermal energies ($P_{\text{bare}} \sim 10^{-85}$):

$$P_{\text{enhanced}} = P_{\text{bare}} \times f_{\text{total}} \sim 10^{-85} \times 10^5 = 10^{-80}$$

For a system with 10^{23} atoms cycling at 10^{12} Hz for 10^5 s:

$$N_{\text{events}} = 10^{23} \times 10^{12} \times 10^5 \times 10^{-80} = 10^{-40}$$

This predicts zero observable events—many orders of magnitude below any detection threshold.

Resolution pathways:

1. **Enhancement factors are vastly underestimated:** The screening enhancement alone, if calculated with measured anomalous screening energies ($U_e \sim$ 300-800 eV per Raiola et al.; RR-007 material-dependent cited ranges), evaluated with the Assenbaum screened-cross-section form $f = \exp(\pi\eta U_e/E)$ at thermal energies, is formally enormous. That expansion is only valid for $U_e \ll E$ (Lean/IR machine check: $U_e < E_{\text{kin}}$, `EQ-ASSEN-DOM-01`); outside the domain the companion software still emits a capped sensitivity factor but sets `formula_claimed=false` — it does **not** claim a physical Assenbaum enhancement. Converting the screening potential to a tunneling enhancement through the proper Gamow integral may yield factors exceeding the adopted 10^3-10^5 band, which is why reported values are capped there. Do not conflate Assenbaum domain validity with the separate RR-007 adiabatic gas-target floor ($U_e \sim$ 20-25 eV).
2. **Collective/coherent multi-body processes:** If LENR proceeds through coherent multi-body tunneling (Takahashi's TSC model) or condensate transitions (Section 2.6), the bare probability is not $P_{\text{bare}} = e^{-2\pi\eta}$ for a single pair, but a collective probability with fundamentally different scaling.
3. **Threshold behavior:** The system may exhibit sharp phase-transition-like behavior where below a critical condition the rate is essentially zero, but above it the rate jumps discontinuously. This is consistent with the observed poor reproducibility—most experiments are below threshold.

Conclusion on Q3: Rate closure is the most stringent quantitative test of the framework. If single-pair enhancement factors are the only mechanism, the numbers do not close (`EQ-RATE-2B-NC-01`, $\log_{10} N = -40$). WH-ENP-01 uses the same-cluster FGR versus $R_{\text{obs}} = P_{\text{W}}/(QV)$ (`EQ-WH-FLOOR-01`); `when.threshold_crossed` is true only when $R_{\text{mode}} \geq R_{\text{obs}}$ with V, τ, P_{occ}, and f_{nuc} labeled cited vs model assumption. The v2 closure demand makes the shortfall explicit: for 1 W in a nominal 1 cm^3, $R_{\text{obs}} \approx 2.6 \times 10^{17}\,\text{m}^{-3}\text{s}^{-1}$; with the rate engine's own site convention $n_{\text{sites}} = 0.5\,(n_e/Z)\,\theta \approx 6.7 \times 10^{26}\,\text{m}^{-3}$ (about 46× below $0.5\,n_D$ at the physical Pd atom density; recorded, not hidden) and $\Gamma_{\text{tun}}(25\,\text{pm}) \approx 6.8 \times 10^{-37}$, the required $P_{\text{occ}} f_{\text{nuc}} C_{\text{lin}}/\tau \approx 5.8 \times 10^{26}\,\text{s}^{-1}$ exceeds the engine cap product $10^{26}\,\text{s}^{-1}$ (`EQ-WH-CAP-01`). At every

cap the assumed 25 pm trap reaches $R_{\max} \approx 4.5 \times 10^{16}\,\mathrm{m^{-3}s^{-1}}$ (factor ≈ 5.8 short) and FCC tetrahedral geometry (137.5 pm, $\Gamma_{\mathrm{tun}} \approx 1.4 \times 10^{-85}$) reaches $\approx 10^{-32}\,\mathrm{m^{-3}s^{-1}}$ (≈ 49 decades short); neither crosses R_{obs}. The $(\theta, \tau, f_{\mathrm{nuc}})$ grid peaks at $\log_{10}(R_{\mathrm{mode}}/R_{\mathrm{obs}}) = -7.4$ and is a scenario table, not a posterior. Rate-UQ / Oracle B ExpHom identities (`EQ-UQ-BF-01`) and the governance enhancement prior upper edge 10^5 remain algebraic honesty floors—not laboratory tunneling measurements or empirical Bayes factors. Overlap DEF (`EQ-OVERLAP-01/02`) records ensemble-count bookkeeping; ensemble product nuclear rate. **Until an experimentally constrained matrix element and state-occupation probability replace the assumed trapped separation, lifetime, and collective factor, multi-body (`trapped_state`) rate outputs are scenario explorations rather than predictions.** The software default remains the conservative **standard two-body reference baseline** (`two_body`): established free-particle formalism used as a reference, *not* a validated lattice-LENR mechanism.

5A.4 Electron Capture Energy Source (Q4)

The Widom-Larsen mechanism requires electrons with kinetic energy > 0.782 MeV for $p + e^- \to n + \nu_e$. Double-layer fields provide ~ 0.1 eV—a factor of $\sim 10^7$ deficit.

Analysis:

1. **Widom-Larsen heavy electron mechanism:** Proposes that collective electromagnetic fluctuations at the metal-hydrogen interface renormalize the electron effective mass to $m_e^* > 2.53\,m_e$. This is a very large mass renormalization by condensed matter standards (typical electron effective masses in metals are 0.5-5 m_e). Whether coherent electromagnetic fields can achieve this remains debated. In particular, Tennfors (Eur. Phys. J. Plus 128, 15, 2013) estimated, from neutron-scattering data on protons in palladium, an electron mass enhancement of less than one percent—far below the threshold required—and Ciuchi et al. similarly questioned the predicted rates; Widom and Larsen rebut these by arguing the process is collective (many-body) rather than two-body. We therefore treat this mechanism as **speculative/contested** and do not rely on it for the paper's primary energy-balance argument.
2. **Alternative: D+D priming reactions:** The D+D $\to$ ^{3}He + n channel (3.27 MeV, athermal neutron) could provide "seed" neutrons at low rates, which then catalyze Oppenheimer-Phillips transmutation chains on heavier lattice nuclei. This avoids the electron capture threshold entirely but requires explaining the initial D+D reaction.
3. **Coherent lattice excitation:** If a lattice phonon field achieves sufficient coherence (e.g., through the mechanisms in Section 2.6), the collective energy available at a single site could in principle exceed the single-phonon energy by a factor of $\sqrt{N_{\mathrm{coherent}}}$. For $N \sim 10^8$ coherent phonons, this gives $\sqrt{10^8} \times 50$ meV ≈ 500 eV—still far below 0.782 MeV.

Conclusion on Q4: No currently proposed mechanism provides a convincing quantitative pathway to produce electrons with $E > 0.782$ MeV in condensed matter systems. The D+D priming pathway (option 2) avoids this problem entirely and is the most physically defensible route to neutron production. The Widom-Larsen mechanism, while elegant, requires independent experimental verification of the mass renormalization claim. Lean `WeakInteraction`

/ IR `EQ-EC-*` and the millielectronvolt deficit lemma (`EQ-OPEN-EC-DEFICIT-01`) document Fermi bookkeeping and the stated threshold gap; they do **not** discharge `EQ-OPEN-EC-01`. Contested Parkhomov / detector-gas SPEC curated rows never feed production rates and are not EC closure.

5A.5 Energy-Product Consistency (Q5)

As detailed in Section 5.2.2, the Miles heat-helium correlation $(2.6 \pm 0.6) \times 10^{11}$ He/s/W is within a factor of ~2 of the 23.8 MeV/ He prediction, providing the strongest quantitative evidence for the D+D $\rightarrow$ He + lattice heat channel. The remaining discrepancy (factor of ~2) is within systematic uncertainties of helium collection efficiency.

However, integrated yields per W $\cdot$ hr are 100-10,000$\times$ lower than expected (Section 5.2.2), indicating either: (a) time-varying reaction rates with duty cycles, (b) incomplete helium collection, or (c) contributions from non-helium-producing channels. This accounting must be resolved with time-resolved simultaneous heat-helium measurements.

---

6. Conclusion

This work presents a unified theoretical and mathematical framework that addresses key LENR phenomena—anomalous heat production, isotopic shifts, minimal radiation, and neutron generation—through a synthesis of Rydberg Matter, halo electrons, double-layer QED effects, Casimir effect, and quantum tunneling. Lattice-bubble interaction physics (heterogeneous nucleation, bubble–surface coupling, boundary layers, acoustic forcing) extends modified bubble dynamics to surface-coupled cavitation. Computational models (Poisson–Schrödinger coupling, nucleation and surface-reaction integration) organize these mechanisms under claim tiers; reported excess-heat and transmutation correlations remain disputed and are not treated as validation of the framework.

Future Directions

1. **Refinement of Computational Models:** Incorporate many-body physics, non-linear effects, and real-time electron–phonon coupling.

2. **Scaling Reactor Designs:** Develop larger prototypes with real-time neutron/helium monitoring to push toward commercial viability.

3. **Interdisciplinary Collaboration:** Encourage a synergy of nuclear engineers, condensed matter physicists, chemists, and computational scientists to expedite the resolution of outstanding LENR challenges.

6.1 Mechanized consistency (Lean)

A companion formalization pilot (`formalization/lean`, Lean 4) mechanically checks **mathematical consistency** of the adopted classical core: Rydberg $n = 10$ identity, Casimir check-geometry band ($|F| \approx 1.3\,\mathrm{mN}$), Gamow penetration anti-monotonicity, Assenbaum domain predicate ($U_e < E_{\mathrm{kin}}$), enhancement composition with cap 10^5 and the illegal two-engine product (`EQ-COMP-03`), the Miles Q-value conversion identity, phonon order $N = Q/\hbar\omega$, two-body rate-budget does-not-close, n-body / Overlap geometry vs trap identities (`EQ-OVERLAP-01/02`; ensemble rate), collapse **clamp / UQ / RP-domain / CR-cap** bookkeeping (`CollapseBounds` / `EQ-BBL-*`—not a cavitation-dynamics theorem), WeakInteraction Fermi G_F/ /Boltzmann bounds (`EQ-EC-*`; `EQ-OPEN-EC-01` closure), Decision ExpHom / `EQ-UQ-BF-01` algebraic BF floors (not empirical Bayes), and aspirational Poisson–Schrödinger continuum DEF (`EQ-PS-01`/`EQ-PS-TOL-01`; never public-beta FEM). An equation intermediate representation (`formalization/ir/`) links paper statements, Python literals, Wolfram numeric checks, and Lean targets. Lean non-closure language (WH-ENP-01, OpenObligations) remains the honesty floor: `lenr_proved` stays false and every `EQ-OPEN-*` stays OPEN until a later evidence-backed closure PR.

Multi-format paper pack (v0.3.7-beta). Academic archival copies of this manuscript and the Lean section→theorem crosswalk ship as:

- Markdown / plain text / LaTeX: `LENR_SIM_APP_v2/docs/v0.3.7-beta/paper/`
- Lean crosswalk module: `formalization/lean/LenrFormal/PaperCrosswalk.lean` (mirrored in the paper pack)
- Equation LaTeX stubs: `LENR_SIM_APP_v2/docs/v0.3.7-beta/latex/`
- NL MD JSON LaTeX Lean matrix: `LENR_SIM_APP_v2/docs/v0.3.7-beta/PROOF_MATRIX.md`

Non-claims. These theorems do **not** prove that LENR occurs, that contested calorimetry is experimentally true, or that the open challenges of §5A (MeV gap, radiation suppression, rate closure, electron-capture threshold, integrated yield) are solved. Speculative mechanisms remain claim-tiered. Status ledger: `docs/FORMALIZATION/FORMAL_PROOF_STATUS.md`.

Would-close-if lemmas (still OPEN). `LenrFormal.OpenQuestionBounds` records closure *predicates* and integer-scaled implications only: Q1 closes if the energy remainder is identically zero (Hagelstein/Dicke/BEC remain named hypotheses); Q2 would close if the lattice $\Gamma_{\mathrm{ph}}/\Gamma_\gamma$ bookkeeping ratio meets 10^{11}; Q3 occupancy/collective factors are Nat inputs to the existing `RateBudget.would_close_if` product (two-body $\log_{10} N = -40$ remains a non-closure theorem); Q4 proves a millielectronvolt **deficit** of $\sim 10^7$ between a stated 0.1 eV double-layer and the 0.782 MeV capture threshold (`EQ-OPEN-EC-DEFICIT-01`) without discharging `EQ-OPEN-EC-01`; Q5 distinguishes an instantaneous Miles EMP band vs integrated 100–10^4 remainder (`EQ-OPEN-YIELD-BAND-01`) without discharging parent obligation `EQ-OPEN-YIELD-01`. `LenrFormal.WorkingHypothesis` (WH-ENP-01) proves a coupled remainder vector and **non-closure** (`EQ-WH-NONCLOSE-01`) without flipping the OPEN table. `LenrFormal.WorkingHypothesisBounds` (WH-ENP-01 v2) adds integer-scaled bounds: the Poisson step ratio $S/(N+1) < 1$ for $N \geq S$ with $\log_{10} P_N \leq -7 \times 10^5 < -11$ on the $S \leq 10^4$ grid (`EQ-WH-FC-01`; conservative cap on the `lgamma` value); the general partition $E_{\mathrm{derived}} + (Q - E_{\mathrm{derived}}) = Q$ with

remainder $= 0 \iff E_{\text{derived}} = Q$ (`EQ-WH-PART-02`); $\lfloor n_D V_{\text{dicke}} \rfloor = 0$, hence $\neg(n \le 0)$ for $n \ge 2$ (`EQ-WH-DICKE-N-01`); the capped product $P_{\text{occ}} f_{\text{nuc}} C_{\text{lin}} \le 1 \cdot 1 \cdot 10^6$ so a demand above the cap is unmet (`EQ-WH-CAP-01`); and the conjunction $\neg Q_1 \wedge \neg Q_2 \wedge \neg Q_3$ (`EQ-WH-NONCLOSE-02`). The Franck–Condon model is a labelled model assumption, not a nature theorem. There is no `theorem lenr_occurs`.

---

7. Mathematical Approaches and Their Relevance to LENR

While the preceding sections introduced key theoretical constructs, LENR also relies heavily on rigorous mathematical methods for both modeling and empirical validation. Below is a concise overview:

1. **Quantum Mechanics and Tunneling**

- **Schrödinger Equation:**

$$-\frac{\hbar^2}{2m}\nabla^2\psi + V(r)\psi = E\psi,$$

central for calculating wavefunction overlaps, tunneling rates, and delocalized-interface-electron interactions. Numerical integration (finite difference, finite element) helps solve complex potential landscapes.

2. **Thermodynamics and Energy Conservation**

- **Excess Heat Models:**

$$Q_{\text{output}} = Q_{\text{input}} + Q_{\text{nuclear}}, \qquad Q_{\text{excess}} = Q_{\text{output}} - Q_{\text{input}},$$

ensuring total energy balance in calorimetric setups.

- **Partition Functions:**

$$Z = \sum_i e^{-E_i/k_B T},$$

used for analyzing statistical behavior of lattice atoms, phonons, and quasi-particles.

3. **Field Theory and Electrodynamics**

- **Extended Maxwell's Equations:**

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}, \qquad \nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \frac{1}{c^2}\frac{\partial \mathbf{E}}{\partial t},$$

crucial for modeling plasma-like oscillations and double-layer fields in electrochemical or gaseous LENR environments.

4. **Nuclear Reaction Cross-Sections**

- **Reaction Rate Expression:**

$$R_{\text{target}} = n\,\sigma_{\text{nuc}}\,v, \qquad R_V = n_1 n_2 \langle \sigma_{\text{nuc}} v \rangle$$

with R_{target} a per-target rate (s^{-1}) and R_V the volumetric rate ($\text{m}^{-3}\ \text{s}^{-1}$). For identical deuterons use $R_V = \frac{1}{2} n_D^2 \langle \sigma v \rangle$. LENR enhancements arise from small effective barriers (via Casimir, high local fields, or RM).

5. **Lattice Dynamics**

- **Harmonic Oscillators:**

$$H = \sum_i \left(\frac{p_i^2}{2m} + \frac{1}{2} k x_i^2 \right),$$

used to approximate phonons interacting with embedded deuterons or protons.

- **Cluster Fusion Models:** Tetrahedral Symmetric Condensate (TSC) → advanced group-theoretic tools describe multi-body fusion in metals (Takahashi's approach; claim tier speculative).

6. **Casimir and Vacuum Fluctuations**

- **Casimir Force:**

$$F_{\text{Casimir}} = -\frac{\pi^2 \hbar c}{240} \frac{A}{d^4},$$

leading to an attractive potential that can enhance electron–nucleus proximity (correlation nuclear proof).

7. **Machine Learning and Statistical Analysis**
   - **Pattern Recognition:** Clustering and neural networks to identify parameter regimes that maximize excess heat or transmutation yields.
   - **Monte Carlo Simulations:** Stochastic exploration of reaction pathways, guiding experimental setups.

8. **Numerical Validation**
   - **Molecular Dynamics (MD):** Tracks atom-by-atom collisions, temperature spikes, and bubble collapse phenomena.
   - **Finite Element Analysis (FEA):** Solves boundary value problems for electric fields, thermal gradients, and mechanical stress in reactor components.

Through this collection of mathematical and computational tools, researchers systematically link theoretical LENR constructs with real-world experimental observations, thereby advancing the rigor, reproducibility, and credibility of the field.

---

8. Glossary of Key Mathematical Symbols

Below is a reference list defining the primary symbols and terms used in this paper:

- $\boldsymbol{\nabla^2}$: The Laplacian operator, $\nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$.
- $\boldsymbol{\phi}$: Electric potential [V].
- $\boldsymbol{\rho}$: Charge density [C/m^3].

- $\boldsymbol{\epsilon_0}$: Vacuum permittivity, $\approx 8.854 \times 10^{-12}\,\mathrm{F/m}$.
- $\boldsymbol{\psi}$: Wavefunction describing the quantum state of a particle.
- $\boldsymbol{V(\mathbf{r})}$: Potential energy function [J] or [eV].
- $\boldsymbol{E}$: (1) Energy eigenvalue in the Schrödinger equation; (2) Electric field strength [V/m], depending on context.
- $\boldsymbol{\hbar}$: Reduced Planck's constant, $\hbar = h/2\pi \approx 1.054 \times 10^{-34}\,\mathrm{J \cdot s}$.
- $\boldsymbol{c}$: Speed of light in vacuum, $c \approx 3 \times 10^8\,\mathrm{m/s}$.
- $\boldsymbol{m}$: Mass (electron, proton, or other particle), context-specific.
- $\boldsymbol{k_B}$: Boltzmann's constant, $k_B \approx 1.381 \times 10^{-23}\,\mathrm{J/K}$.
- $\boldsymbol{a_0}$: Bohr radius, $a_0 \approx 0.0529\,\mathrm{nm}$.
- $\boldsymbol{n}$: (1) Principal quantum number; (2) Number density [$\mathrm{m^{-3}}$], depending on context.
- $\boldsymbol{n_1, n_2}$: Number densities of two (possibly identical) reacting species [$\mathrm{m^{-3}}$]; use $\frac{1}{2}n^2$ rather than $n_1 n_2$ when the species are identical (§5.1.1).
- $\boldsymbol{N}$: Number density of target nuclei (OP process, §2.7) [$\mathrm{m^{-3}}$].
- $\boldsymbol{\Phi_n}$: Neutron flux [$\mathrm{m^{-2}s^{-1}}$] (§2.7).
- $\boldsymbol{a}$: Characteristic distance scale (e.g., extended orbital radius).
- $\boldsymbol{r_0}$: Nuclear core or reference radius.
- $\boldsymbol{P}_{\mathrm{overlap}}$: Overlap probability between interface electron and nuclear wavefunctions.
- $\boldsymbol{p}$: Proton, or momentum operator ($\hat{p}$) depending on context.
- $\boldsymbol{e^-}$: Electron.
- $\boldsymbol{\nu_e}$: Electron neutrino.
- $\boldsymbol{F}_{\mathrm{Casimir}}$: Casimir force.
- $\boldsymbol{A}$: Area (e.g., Casimir effect) [$\mathrm{m^2}$].
- $\boldsymbol{d}$: Separation distance (e.g., Casimir or double-layer) [m].
- $\boldsymbol{T}$: Temperature [K].
- $\boldsymbol{T_c}$: Critical temperature for Bose–Einstein condensation.
- $\boldsymbol{\zeta(3/2)}$: Riemann zeta function at 3/2, $\zeta(3/2) \approx 2.612$.
- $\boldsymbol{\Delta V}$: Potential difference [V].
- $\boldsymbol{\mu}$: (1) Dynamic viscosity [$\mathrm{Pa \cdot s}$]; (2) reduced mass [kg], depending on context.
- $\boldsymbol{\sigma}$: (1) Generic cross-section [$\mathrm{m^2}$]; (2) surface tension [N/m] in Rayleigh–Plesset. Context clarifies usage.
- $\boldsymbol{\sigma_0}$: Base/reference cross-section [$\mathrm{m^2}$].
- $\boldsymbol{\sigma}_{\mathrm{nuc}}$: Nuclear (fusion) cross-section used in the generic rate expression $R_{\mathrm{rate}} = n\sigma_{\mathrm{nuc}}v$ (§7) [$\mathrm{m^2}$].
- $\boldsymbol{\sigma}_{\mathrm{OP}}$: Oppenheimer–Phillips capture cross-section (§2.7) [$\mathrm{m^2}$].
- $\boldsymbol{\sigma}_{\mathrm{surf}}$: Bubble surface tension in the Rayleigh–Plesset equation (§3.2) [N/m].
- $\boldsymbol{\langle \sigma v \rangle}$: Thermally-averaged cross-section–velocity product (§5.1.1) [$\mathrm{m^3/s}$].
- $\boldsymbol{R}$: (1) Reaction rate **per unit volume**, [$\mathrm{reactions \cdot m^{-3}s^{-1}}$] — see §2.7/§5.1.1/§7 for three distinct regimes of this quantity; (2) Bubble radius in the Rayleigh–Plesset equation [m].
- $\boldsymbol{P}_{\mathrm{tunnel}}$: Quantum tunneling (Gamow) probability, dimensionless.
- $\boldsymbol{P}_{\mathrm{threshold}}$: Probability of exceeding reaction threshold conditions, dimensionless (§5.1.1).
- $\boldsymbol{f}_{\mathrm{enhancement}}$: Cumulative tunneling-enhancement factor, dimensionless; $f_{\mathrm{enhancement}} =$

$f_{\text{screening}} \times f_{\text{field}} \times f_{\text{coherence}} \times f_{\text{phonon}}$ (§2.8, §5.1.1).

- $Q_{\text{nuclear}}, Q_{\text{input}}, Q_{\text{excess}}$: Energy quantities in calorimetry [J].
- ζ: Generic variable for partition functions or the Riemann zeta function, as context requires.

Formalization symbol map (IL-015). In the Lean/IR ledger (`docs/FORMALIZATION/CLAIM_TAXONOMY.md`), overloaded symbols are disambiguated as `E_eigen` / `E_field` / `E_kin`, `n_prin` / `n_dens`, and `R_rate` / `R_bubble` / `R_overlap_count`.

References

Citations below support the mathematical modeling in this paper. Preference is given to sources tied to **replicable experimental reports** (calorimetry, helium/isotope assays, screening measurements) and to **established mechanisms** used in the equations (tunneling, Casimir, Oppenheimer–Phillips, phonon–nuclear coupling). Speculative mechanisms remain claim-tiered in the body text even when cited.

Subtitle homage

[**vaswani2017**] Vaswani, A., Shazeer, N., Parmar, N., Uszkoreit, J., Jones, L., Gomez, A. N., Kaiser, Ł., & Polosukhin, I. (2017). Attention Is All You Need. *Advances in Neural Information Processing Systems* 30.
Methodological metaphor only—not a LENR physics source. The subtitle "Coupling is all you need" deliberately echoes this work's framing of coupled multi-agent representations.
Link: https://arxiv.org/abs/1706.03762 · NeurIPS 2017

Primary sources for mathematical modeling (replicable evidence & established mechanisms)

Drawn from the project's evidence-base bibliography (`Unified_LENR_Theory_D.Acs.docx.md`) and from DOI-backed screening / heat–helium constraints used by the companion simulation.

1. Fleischmann, M., & Pons, S. (1989). Electrochemically induced nuclear fusion of deuterium. *Journal of Electroanalytical Chemistry*, 261(2), 301–308.
   Link: https://doi.org/10.1016/0022-0728(89)80006-3
2. McKubre, M. C. H., et al. (1994). Excess power observations in electrochemical studies of the D/Pd system. *Journal of Electroanalytical Chemistry*, 368, 55–66.
   Link: https://doi.org/10.1016/0022-0728(94)87011-6
3. Storms, E. (2010). A critical analysis of the LENR phenomenon. *Naturwissenschaften*, 97(10), 861–881.
   Link: https://doi.org/10.1007/s00114-010-0711-x
4. Arata, Y., & Zhang, Y.-C. (2002). Observation of anomalous heat release and helium-4 production from highly deuterated palladium fine particles. *Proceedings of the Japan*

Academy, Series B, 78(7), 57–61.
Link: https://doi.org/10.2183/pjab.78.57

5. Mizuno, T. (2007). Anomalous energy generation during treatment of nickel with hydrogen gas. *Journal of Nuclear Science and Technology*, 44(2), 331–338.
Link: https://doi.org/10.1080/18811248.2007.9711293
6. Mizuno, T., & Rothwell, J. (2019). Increased excess heat from palladium deposited on nickel. *Journal of Condensed Matter Nuclear Science*, 29, 1–30.
Link: http://www.iscmns.org/CMNS/JCMNS-Vol29.pdf
7. Miles, M. H., et al. (1993). Correlation of excess power and helium production. *Journal of Electroanalytical Chemistry*, 346, 99–117.
Link: https://doi.org/10.1016/0022-0728(93)85700-H
8. Steinetz, B. M., et al. (2020). Novel nuclear reactions observed in bremsstrahlung-irradiated deuterated metals. *Physical Review C*, 101, 044610.
Link: https://doi.org/10.1103/PhysRevC.101.044610
9. Pines, V., et al. (2020). Nuclear fusion reactions in deuterated metals. *Physical Review C*, 101, 044609.
Link: https://doi.org/10.1103/PhysRevC.101.044609
10. Raiola, F., et al. (2002). Electron screening in d(d,p)t for deuterated metals. *European Physical Journal A*, 13, 377–382.
Link: https://doi.org/10.1007/s10050-002-8766-5
11. Huke, A., et al. (2008). Enhancement of deuteron-fusion reactions in metals. *Physical Review C*, 78, 015803.
Link: https://doi.org/10.1103/PhysRevC.78.015803
12. Czerski, K., et al. (2023). Electron screening in palladium. *Physics Letters B*, 845, 137750.
Link: https://doi.org/10.1016/j.physletb.2023.137750
13. Fralick, G., Decher, R., & Rousar, D. (2003). Anomalous heat effects in metal hydrides. NASA/TM—2003-212795.
Link: https://ntrs.nasa.gov/citations/20040152112
14. Iwamura, Y., Itoh, T., & Takahashi, T. (2024). Anomalous heat generation in nano-composite hydrides. *Japanese Journal of Applied Physics*, 63(3), 037001.
Link: https://doi.org/10.35848/1347-4065/ad2622
15. Bellucci, S., Cardone, F., & Pistella, F. (2023). Review of experiments reporting non-conventional phenomena in nuclear matter. *Symmetry*, 15(8), 1507.
Link: https://doi.org/10.3390/sym15081507
16. Oppenheimer, J. R., & Phillips, M. (1935). Note on the transmutation function for deuterons. *Physical Review*, 48(7), 500–502.
Link: https://doi.org/10.1103/PhysRev.48.500
17. Bordag, M., Mohideen, U., & Mostepanenko, V. M. (2001). New developments in the Casimir effect. *Physics Reports*, 353(1–3), 1–205.
Link: https://doi.org/10.1016/S0370-1573(01)00015-1
18. Hagelstein, P. L., & Chaudhary, I. U. (2015). Coherent energy exchange between lattice and nuclear degrees of freedom. *Journal of Condensed Matter Nuclear Science*, 15, 1–24.

Link: http://www.iscmns.org/CMNS/JCMNS-Vol15.pdf

19. Takahashi, A. (2004). Tetrahedral symmetric condensate fusion model and multi-body fusion in deuterated metal. *Journal of Condensed Matter Nuclear Science*, 1, 4–17.
Link: http://www.iscmns.org/CMNS/JCMNS-Vol1.pdf
20. Takahashi, A. (2016). Fundamental of rate theory for CMNS. *Journal of Condensed Matter Nuclear Science*, 19, 298–315. (Basis for the software `trapped_state` rate mode.)
21. Holmlid, L. (2015). Rydberg Matter and its detection: A review of experimental techniques. *Journal of Physics: Conference Series*, 635, 012002.
Link: https://doi.org/10.1088/1742-6596/635/1/012002
22. Widom, A., & Larsen, L. (2006). Ultra low-momentum neutron catalyzed nuclear reactions on metallic hydride surfaces. *The European Physical Journal C*, 46(1), 107–111.
Link: https://doi.org/10.1140/epjc/s2006-02479-8
23. Preparata, G. (1995). *QED Coherence in Matter*. World Scientific.
Link: https://www.worldscientific.com/worldscibooks/10.1142/2739
24. Schwinger, J. (1990). Cold fusion: A hypothesis. *Zeitschrift für Naturforschung A*, 45, 756–758.
Link: https://www.lenr-canr.org/acrobat/SchwingerJcoldfusion.pdf
25. Kim, Y. E., & Zubarev, A. L. (2001). Theory of Bose–Einstein condensation mechanism for deuteron-induced nuclear reactions in micro/nano-scale metal grains and particles. *Physical Review A*, 64, 013603.
Link: https://doi.org/10.1103/PhysRevA.64.013603
26. Taleyarkhan, R. P., et al. (2002). Evidence for nuclear emissions during acoustic cavitation. *Science*, 295(5561), 1868–1873.
Link: https://doi.org/10.1126/science.1067589 *Honesty:* Contested / disputed nuclear-emission claim; retained as acoustic-cavitation literature anchor only—not laboratory LENR proof. See also `pyyhtia2026` (electrolytic CR-39 cavitation confounder).
27. Menlove, H. O., et al. (1989). Neutron emission from titanium exposed to deuterium gas at high pressure. Los Alamos National Laboratory Report.
Link: https://www.osti.gov/biblio/5039188
28. Bockris, J. O'M., Lin, G. H., & Bush, B. F. (1991). Tritium and neutron generation in electrolysis of heavy water. *Proceedings of the Electrochemical Society* / LENR-CANR archive.
Link: https://lenr-canr.org/acrobat/BockrisJdoestritiu.pdf
29. Wolf, K. L., et al. (1990). Tritium production in deuterated palladium. Texas A&M / related reports.
Link: https://link.springer.com/article/10.1007/BF02627575
30. Celani, F., et al. (2014). Experimental results on sub-micro structured Pd wires and thin films. *Journal of Condensed Matter Nuclear Science*, 13, 56–68.
Link: http://www.iscmns.org/CMNS/JCMNS-Vol13.pdf
31. Bakranov, N., Didenko, V., & Nagel, D. (2024). Nanomaterials engineering for enhanced LENR: A comprehensive review. *Frontiers in Materials*, 11, 1500487.
Link: https://doi.org/10.3389/fmats.2024.1500487
32. Levi, G., et al. (2013). Indications of anomalous heat energy production in a reactor device

containing hydrogen-loaded nickel powder. arXiv:1305.3913.
Link: https://arxiv.org/abs/1305.3913

33. Del Giudice, E., Doglia, S., Milani, M., & Vitiello, G. (1985). Electrodynamical coherence in biological systems. *Physical Review Letters*, 54(7), 622–625.
Link: https://doi.org/10.1103/PhysRevLett.54.622
34. Safajuei, F., Safajuei, S., & Hajimohammadi, M. (2023). Production of ultra-low momentum neutrons on metallic hydride surfaces by heavy electrons. *Iranian Journal of Science and Technology, Transactions A: Science*, 47, 995–1002.
Link: https://doi.org/10.1007/s40995-023-01437-3

Supplementary keys (software / UQ traceability)

These `AuthorYear` keys resolve claims in the companion app and `docs/REVIEW_2026/TRACEABILITY_MATRIX.md`:

- **[steinetz2020exp]** → entry 8 above
- **[pines2020theory]** → entry 9 above
- **[raiola2002]** → entry 10 above
- **[huke2008]** → entry 11 above
- **[czerski2023]** → entry 12 above
- **[miles1993]** → entry 7 above
- **[takahashi2016]** → entry 20 above
- **[hagelstein2025]** Hagelstein, P. L. (2025). Relativistic phonon–nuclear coupling / lossy spin-boson reviews (MeV-gap channel).
- **[screeningreview2024]** Electron screening in laboratory nuclear reactions. *Particles*, 7(3), 50. DOI: 10.3390/particles7030050
- **[storms2015]** Storms, E. (2015). Efficient production of energy from the LENR effect. *Current Science*, 108(4), 574–577.
- **[tennfors2013]** Tennfors, E. (2013). On the assumptions for an effective electron mass. *European Physical Journal Plus*, 128, 15. DOI: 10.1140/epjp/i2013-13015-3
- **[mulnix2026]** Mulnix, A. (2026). Structural theory of sonoluminescence via Klein–Gordon / Rayleigh–Plesset harmonic locking. Zenodo. DOI: 10.5281/zenodo.19079074

- [**pyyhtia2026**] Pyyhtiä, K., et al. (2026). Investigation of CR-39 damaging mechanisms in electrochemical environments. *Journal of Electroanalytical Chemistry*, 1013, 120104. DOI: 10.1016/j.jelechem.2026.120104. OA experimental support that HER nanobubble cavitation damages CR-39 and confounds pit=nuclear interpretations; does **not** prove LENR.

- [**nguyen2026**] Nguyen, T., et al. (2026). Quantum shell structure of proton–neutron pairing. *Nature*. DOI: 10.1038/s41586-026-10616-2

- [**floquetwkb2026**] Floquet–WKB analysis of lattice confinement fusion under a driven barrier (2026). DOI: 10.14293/PR2199.002776.v1

- [**parkhomov2018**] Parkhomov, A. G. (2018). Nuclear reactions in LENR: review of experimental results (LENR Academy curated subset).
