

Foundation of Analytics: Lecture 3

Dihui Lai

dlai@wustl.edu

August 2, 2020

Content

- Random Variables: Dependent, Independent, Correlation
- Linear Regression of One Variable
- Linear Regression of Multiple Variables
- Logistic Regression

Correlations between Random Variables

Let's look at a few pairs of data points?

- $\vec{x} = [0.5, 0.6, 0.1, -0.3, 2.3], \vec{y} = [0.5, 0.6, 0.1, -0.3, 2.3]$
- $\vec{x} = [0.5, 0.6, 0.1, -0.3, 2.3], \vec{y} = [0.6, 0.6, 0.12, -0.3, 2.3]$
- $\vec{x} = [0.5, 0.6, 0.1, -0.3, 2.3], \vec{y} = [0.02, -0.2, 0.2, 2.1, -0.5]$

What can you tell about the relationship between \vec{x} and \vec{y} ?

Correlations between Random Variables

Given two random variables X and Y , denote the mean and variance of the two variables as $E[X] = \mu_X$, $E[Y] = \mu_Y$, $Var[X] = \sigma_X^2$, $Var[Y] = \sigma_Y^2$.

The covariance of X and Y is the number defined by

$$\begin{aligned} Cov(X, Y) &= E[(X - \mu_X)(Y - \mu_Y)] \\ &= E[XY] - \mu_X \mu_Y \end{aligned}$$

Empirical Estimation of Covariance

$$\begin{aligned} Cov(X, Y) &= \frac{(x - \mu_x)(y^T - \mu_y)}{N} (\text{empirical}) \\ Cov(X, Y) &= \frac{(x - \mu_x)(y^T - \mu_y)}{N - 1} (\text{unbiased}) \end{aligned}$$

Correlations between Random Variables

The correlation of the two random variables is the number defined by

$$\rho_{XY} = \frac{Cov(X, Y)}{\sigma_X \sigma_Y}$$

Correlations between Random Variables

Calculate the covariance/correlation of

- Example 1:

$$\vec{x} = [2, -2, -2, 2], \vec{y} = [2, -2, -2, 2]$$

We have $\mu_x = 0$, $\mu_y = 0$, $\sigma_x^2 = 4$, $\sigma_y^2 = 4$, $E[XY] = 4$ Therefore
 $Cov(X, Y) = 4 - 0 = 4$ and $\rho_{xy} = 4/(2 * 2) = 1$

- Example 2:

$$\vec{x} = [2, -2, -2, 2], \vec{y} = [2, 0, -2, 0]$$

We have $\mu_x = 0$, $\mu_y = 0$, $\sigma_x^2 = 4$, $\sigma_y^2 = 2$, $E[XY] = 2$ Therefore
 $Cov(X, Y) = 2 - 0 = 2$ and $\rho_{xy} = 2/(2 * \sqrt{2}) = 1/\sqrt{2}$

Linear Regression with One Variable

Data set:

$$y = \begin{bmatrix} y^1 \\ y^2 \\ \vdots \\ y^n \end{bmatrix}, X = \begin{bmatrix} x^1 \\ x^2 \\ \vdots \\ x^n \end{bmatrix}$$

Linear Regression with One Variable

Assume y is linearly depending on x i.e.

$$\hat{y} = \beta_0 + \beta_1 x$$

Find $\hat{\beta}$ that minimize the estimation error

$$\epsilon = \sum_{i=1}^n (y^i - \hat{y}^i)^2 = \sum_{i=1}^n (y^i - \beta_0 - \beta_1 x^i)^2$$

i.e.

$$\frac{\partial \epsilon}{\partial \beta_1} = 0 \rightarrow \sum_{i=1}^n (y^i - \beta_0 - \beta_1 x^i) x^i = 0$$

$$\frac{\partial \epsilon}{\partial \beta_0} = 0 \rightarrow \sum_{i=1}^n (y^i - \beta_0 - \beta_1 x^i) = 0$$

$$\beta_0 \sum_{i=1}^n x^i = \sum_{i=1}^n y^i x^i - \beta_1 \sum_{i=1}^n x^i x^i$$

$$\beta_0 = \frac{1}{n} \sum_{i=1}^n (y^i - \beta_1 x^i) = \bar{y} - \beta_1 \bar{x}$$

Insert the second equation to the first, we have

$$n\bar{x}\bar{y} - \beta_1 n\bar{x}\bar{x} = \sum_{i=1}^n y^i x^i - \beta_1 \sum_{i=1}^n x^i x^i$$

Therefore,

$$\beta_1 = \frac{\frac{1}{n} \sum_{i=1}^n x^i y^i - \bar{x}\bar{y}}{\frac{1}{n} \sum_{i=1}^n x^i x^i - \bar{x}^2} = \frac{Cov(X, Y)}{Var(X)} = \rho_{XY} \frac{\sigma_Y}{\sigma_X}$$

Multivariate Linear Regression

Data set:

$$\begin{bmatrix} y^1 & x_0^1 & x_1^1 & x_2^1 & \dots & x_m^1 \\ y^2 & x_0^2 & x_1^2 & x_2^2 & \dots & x_m^2 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ y^n & x_0^n & x_1^n & x_2^n & \dots & x_m^n \end{bmatrix}$$

Multivariate Linear Regression

Assume y is a linear superposition of multiple x 's

$$\hat{y} = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \dots + \beta_m x_m$$

or simply

$$\hat{y} = \sum_{j=1}^m \beta_j x_j$$

Multivariate Linear Regression

Estimate β 's that best fits the data, we need to minimize the error

$$\begin{aligned}\epsilon &= \sum_{i=1}^n (y^i - \hat{y}^i)^2 \\ &= (y - \hat{y})^T (y - \hat{y})\end{aligned}$$

Use basic calculus we know, we want to have the β s satisfy the following equation set:

$$\frac{\partial \epsilon}{\partial \beta_j} = 0, j = 1, 2, 3, 4 \dots m$$

i.e.

$$\sum_{i=1}^n \frac{\partial (y^i - \hat{y}^i)^2}{\partial \beta_j} = 0$$

$$\sum_{i=1}^n (y^i - \hat{y}^i) \frac{\partial \hat{y}^i}{\partial \beta_j} = 0$$

$$\sum_{i=1}^n (y^i - \hat{y}^i) x_j^i = 0$$

Multivariate Linear Regression

Written in matrix formula we require

$$(y - X\beta)^T X = 0$$

or after transposing

$$X^T y - X^T X \beta = 0$$

Therefore

$$\beta = (X^T X)^{-1} X^T y$$