Supporting Information

for

Photo-induced coupling of CO and vinylidene ligands – formation of cyclobutane-1,3-diones

Mokhles M. Abd-Elzaher[#], Bernhard Weibert, and Helmut Fischer*

Fachbereich Chemie, Universität Konstanz, Fach M727, 78457 Konstanz, Germany

3. Experimental

3.1. General

All operations were carried out under nitrogen by using conventional Schlenk techniques. Solvents were dried by refluxing over sodium-benzophenone ketyl (THF, diethyl ether) or CaH₂ (pentane, dichloromethane) and were freshly distilled prior to use. The silica gel used for chromatography (J. T. Baker, silica gel for flash chromatography) was nitrogen saturated. The yields refer to analytically pure compounds and were not optimized.

The vinylidene complexes $\mathbf{1a}^1$ and $\mathbf{2a}^2$ and the allenylidene complex $\mathbf{5}^3$ were synthesized as previously described. Complexes $\mathbf{1b}$ and $\mathbf{2b}$ wee synthesized analogously. [Me₃O]BF₄ were prepared according to the literature procedures.⁴ Metal hexacarbonyls (Aldrich), 3,3-dimethylbut1-yne (Aldrich), pent-1-yne (Fluka), diphenyl diazene (Aldrich), and trimethylene sulfide were commercial products and were used without further purification.

IR: FT-IR spectrophotometer (FTS70), Bio-Rad. All IR measurements were carried out at 298 K. ¹H-NMR, ¹³C-NMR: Bruker AC 250. Unless specifically mentioned, NMR spectra were recorded in CDCl₃ at room temperature. Chemical shifts are reported in ppm relative to internal TMS. MS: Finnigan MAT 312 (EI). Elemental analysis: CHN-Analyser (CHN-O-RAPID) Heraeus. Photolysis reactions were carried out in a duran glass apparatus using a mercury high pressure lamp (TQ 150, Fa. Heraeus).

General procedure for photolysis of the vinylidene complexes 1a,b and 2a,b

About 5 mmol of the vinylidene complex was dissolved in 300 ml of pentane/dichloromethane (1:1) and photolyzed at -30°C for about 1 h in a Duran-glass apparatur (cut-off wavelength: 300 nm). The reaction was followed by IR spectroscopy. The photolysis was stopped when the vinylidene complex could not be detected any more. The solution was allowed to warmed to room temp. and the volume of the solution was reduced to about 10 ml. The solution was

chromatography at -30° C on silica. First, with pentane [M(CO)₆] was eluted. Subsequent elution with pentane/dichloromethane (ratio vide supra) gave the a yellow band containing the cyclobutane-1,3-dione.

2,4-Bis(1,2,2-trimethylpropylidene)-cyclobutane-1,3-dione (3a)

Photolysis of [(CO)₅Cr=C=C(t-Bu)Me] (**1a**) and elution of the yellow band with pentane/dichloromethane (9:1) gave 1.73 mmol of **3a** (62 % based on **1a**). Photolysis of [(CO)₅W=C=C(t-Bu)Me] (**2a**) and chromatography gave 0.75 mmol of **3a** (27 % based on **2a**). The microcrystalline powder obtained by recrystalization from pentane/dichloromethane (9:1) contained small amounts of dichloromethane. Mp: 175°C. IR (pentane): ν (CO) 1693 m, ν (C=C) 1590 m cm⁻¹. ¹H-NMR: δ 2.41 (s, 6H, Me), 1.30 (s, 18H, t-Bu). ¹³C-NMR: δ 188.9 (C=O), 175.3 (C=CCH₃), 150.4 (C=CCH₃), 38.8 (C(CH₃)₃), 28.5 (C(CH₃)₃)), 17.7 (=CCH₃). MS (m/z, %): 248 (21.7) [M⁺], 233 (19.4) [M⁺- Me], 205 (15.7) [M⁺ - Me - CO], 190 (16.4) [M⁺ - Me - CO - Me]. Anal. Calcd. for C₁₆H₂₄O₂ (248.37) + 0.1 CH₂Cl₂: C, 75.21; H, 9.42. Found: C, 75.22; H. 9.58. Elemental analysis: C₁₆H₂₄O₂ (248.37) C, 77.36; H, 9.75;

2,4-Bis(1-methylbutylidene)cyclobutane-1,3-dione (3b)

Photolysis of [(CO)₅Cr=C=C(Pr)Me] (**1b**) and elution of the yellow band with pentane/dichloromethane (8:2) gave 1.73 mmol of **3b** (58 % based on **1b**). Photolysis of [(CO)₅W=C=C(Pr)Me] (**2b**) chromatography gave 0.48 mmol of **3b** (19 % based on **2b**). Mp: 182° C. IR (pentane): v(CO) 1687 m, v(C=C) 1585 m cm⁻¹. ¹H-NMR: δ 2.63 (t, J = 7.5 Hz, 4H, CH₂), 2.25 (s, 6H, =CCH₃), 1.59 (sext, J = 7.5 Hz, 4H, CH₂), 0.96 (t, J = 7.4 Hz, 6H, CH₃). ¹³C-NMR: δ 190.4 (C=O), 163.5 (=CCH₃), 150.9 (C=CCH₃), 36.9 (=CCH₃), 21.4 (=CCH₂), 19.5 (CH₂CH₃), 13.8 (CH₂CH₃). MS (m/z, %): 220 (81.0) [M⁺], 177 (50.4) [M⁺ - CH₂CH₂CH₃], 134

(80.6) [M $^+$ - 2Pr]. Anal. Calcd. for $C_{14}H_{20}O_2$ (220.3): C, 76.31; H, 9.16. Found: C, 75.63; H, 9.24.

Photolysis of 1a in the presence of PPh₃

The photolysis of **1a** in the presence of 2.5 equivalents of PPh₃ was carried out as decribed above. Chromatography at -30°C on silica first with pentane gave [Cr(CO)₆] and then with pentane/dichloromethane (9:1) the cyclobutadione. Finally, with dichloromethane a second yellow band containing [CO)₅Cr(PPh₃)] was eluted. The complex was identified by comparison of its spectra with published data.⁵

Photolysis of 2a in the presence of PPh₃

The photolysis of **2a** in the presence of 2.5 equivalents of PPh₃ was carried out analogously. The reaction was followed by IR spectroscopy. Photolysis was stopped when the vinylidene complexes could not be detected any more. The solution was removed in vacuo. The cyclobutandione and [W(CO)₆] were extracted twice with 10 ml of pentane each and then twice with 15 ml of pentane/dichloromethane (1:1). The remaining [CO)₅W(PPh₃)] was still contaminated with small amounts of [W(CO)₆]. The complex was identified by comparison of its spectra with those published.⁵

Photolysis of 1a and 2a in the presence of trimethylene sulfide

The photolysis of 1a in the presence of 2.5 eq. of trimethylene sulfide was carried out as described above. Chromatography at -30° C on silica first with pentane gave [Cr(CO)₆] and then with pentane/dichloromethane (9:1) the cyclobutadione 3a. Finally, with pentane/dichloromethane (7:3) a second yellow band containing the complex 5 was eluted. Yield: 1.35 mmol (27 % based on 1a). The complex was fully characterized and additionally identified by comparison of its spectroscopic data with those published earlier. Mp: 29 - 30°C. IR (pentane): v(CO) 1938 s, 1946 vs, 2069 m cm⁻¹. H-NMR: δ 3.40 (t, J = 8.0 Hz, 4H, CH₂),

2.94 (quint, J = 8.0 Hz, 2H, CH₂). ¹³C-NMR: δ 221.5 (trans-CO), 214.8 (cis-CO), 35.9 (2CH₂), 25.9 (CH₂). MS (m/z, %): 266 (4.2) [M⁺], 238 (3.9) [M⁺ - CO], 210 (4.5) [M⁺ - 2CO], 182 (4.6) [M⁺ - 3CO], 154 (7.7) [M⁺ - 4CO], 126 (19.9) [M⁺ - 5CO], 98 (15.3) [M⁺ - 5CO - 2CH₂], 84 (29.8) [M⁺ - 5CO - 3CH₂]. Anal. Calc. for C₈H₆O₅CrS (266.19): C, 36.10; H, 2.27. Found C, 35.84; H 2.31.

Photolysis of 2a, chromatography and separation of $[(CO)_5W\{S(CH_2)_3\}]$ from the reaction products were carried analogously. The complex was identified by comparison of its spectra with published data.⁶

Photolysis of pentacarbonyl(diphenylallenylidene)chromium

A solution of 2.72 g (7.1 mmol) of pentacarbonyl(diphenylallenylidene)chromium (5) in 300 ml of pentane/dichloromethane (1:1) was photolyzed at -30°C for 4 h. The color of the solution gruadually changed from blue to green. The reaction was followed by IR spectroscopy. After 4 h the irradiation was stopped and the volume of the solution reduced to ca. 10 ml. Chromatography at -30°C on silica gave 5 fractions: the first one (slightly yellow, elution with pentane) contained [Cr(CO)₆], the second one (blue, elution with pentane/dichloromethane (20:1)) contained starting allenylidene complex, the third and the forth ones (different degree of green, elution with pentane/dichloromethane (10:1) and (8:2)) yielded mixtures of the allenylidene complex and (red-brown. tetraphenylhexapentaene **(6)**. Finally, the fifth band elution with pentane/dichloromethane (3:2)) contained pure 6. The compound 6 was identified by comparison of its spectra with published data.⁷

Yield: 1.13g (83 % based on **5**). Mp: 204°C (ref. f: 202-207°C). IR: ν(cumulene) 2154 m, 2125 s, 2054 m, ν(C=C) 1599 m cm⁻¹. ¹H-NMR: δ 7.82-6.63 (m, 20H, Ph). MS (m/z, %): 380 (0.91)

[M⁺], 303 (0.8) [M⁺ - Ph], 226 (1.1) [M⁺ - 2Ph], 214 (0.9) [M⁺ - 2Ph - C], 202 (1.3) [M⁺ - 2Ph - 2C].

X-Ray structural analyses of 3a

Single crystals suitable for X-ray structural analyses were obtained from pentane/dichloromethane (9:1). The measurements were performed at 188(2) K with a crystal mounted on a glass fiber on a Siemens P4 diffractometer (graphite monochromator, Mo-K α radiation, $\lambda = 0.71073$ Å). For crystal data and refinement details see the Table 1. The structure was solved by direct methods using the SHELXTL-97 program package. The positions of the hydrogen atoms were found and refined isotropically. All other atoms were refined anisotropically.

References

- 1 Karl, C. C.; Joneleit, S.; Abd-Elzaher, M. M.; Weibert, B.; Fischer, H. . *J. Organomet. Chem.* **2003**, 677, 46.
- Fischer, H.; Podschadly, O.; Früh, A.; Troll, C.; Stumpf, R.; Schlageter, A. Chem. Ber.1992, 125, 2667.
- 3 Fischer, H.; Reindl, D.; Roth, G. Z. Naturforsch. **1994**, 49b, 1207.
- 4 Meerwein, H.; Hinz, G.; Hofmann, P.; Kroning, E.; Pfeil, E. *J. prakt. Chem.* **1937**, *147*, 257.
- 5 Magee, T. A.; Matthews, C. N.; Wang, T. S.; Wotiz, J. H. *J. Am. Chem. Soc.* **1961**, *83*, 3200.
- Adams, R.D.; Fallon, S. B.; Perrin, J. R.; Queisser, J. A.; Yamamoto, J. H. *Chem. Ber.* 1996, 129, 313.
- 7 Hartzler, H. D. J. Am. Chem. Soc. **1961**, 83, 4990:

8 Sheldrick, G. M. SHELX-97, Programs for Crystal Structure Analysis, University of Göttingen, Göttingen (Germany), **1997**.

Structure of compound $(CO)_2[C=C(t-Bu)Me]_2(3a)$




Table 1. Crystal data and structure refinement for compound 3a.

Empirical formula $C_8H_{12}O$

Formula weight 124.18

Temperature 188(2) K

Wavelength 0.71073 Å

Crystal system, space group Monoclinic, P2(1)/n

Unit cell dimensions a = 6.829(2) Å $\alpha = 90 \text{ deg}$.

b = 10.524(3) Å $\beta = 96.11(2) \text{ deg.}$

c = 10.179(3) Å $\gamma = 90 \text{ deg.}$

Volume 727.4(4) Å³

Z, Calculated density 4, 1.134 Mg/m³

Absorption coefficient 0.073 mm⁻¹

F(000) 272

Crystal size, colour, habit 0.4 x 0.35 x 0.3 mm, yellow, block

Theta range for data collection 2.79 to 26.99 deg.

Limiting indices -8 <= h <= 8, -13 <= k <= 13, -13 <= l <= 13

Reflections collected / unique 3618 / 1586 [R(int) = 0.0869]

Completeness to theta = 26.99 99.9 %

Absorption correction None

Refinement method Full-matrix least-squares on F²

Data / restraints / parameters 1586 / 0 / 130

Goodness-of-fit on F² 1.021

Final R indices [I>2 \square (I)] R1 = 0.0451, wR2 = 0.1044

R indices (all data) R1 = 0.0632, wR2 = 0.1129

Largest diff. peak and hole 0.308 and -0.161 e· Å⁻³

Table 2. Bond lengths [Å] and angles [deg] for compound **3a**.

C(5)-C(6)-H(6C)	108.2(12)	
H(6A)-C(6)-H(6C)	111.7(17)	
H(6B)-C(6)-H(6C)	106.4(16)	
C(5)-C(7)-H(7C)	109.3(10)	
C(5)-C(7)-H(7B)	113.5(10)	
H(7C)-C(7)-H(7B)	109.8(17)	
C(5)-C(7)-H(7A)	110.2(12)	
H(7C)-C(7)-H(7A)	104.0(16)	
H(7B)-C(7)-H(7A)	109.6(15)	
C(5)-C(8)-H(8A)	110.7(11)	
C(5)-C(8)-H(8B)	108.3(12)	
H(8A)-C(8)-H(8B)	113.8(19)	
C(5)-C(8)-H(8C)	110.1(15)	
H(8A)-C(8)-H(8C)	106.8(16)	
H(8B)-C(8)-H(8C)	107.0(18)	

Table 3. Torsion angles [deg] for compound **3a**.

C(4)-C(3)-C(2)-C(1)	0.4(2)
C(5)-C(3)-C(2)-C(1)	178.57(13)
C(4)-C(3)-C(2)-C(1)#1	-177.35(16)
C(5)-C(3)-C(2)-C(1)#1	0.8(3)
C(3)-C(2)-C(1)-O(1)	2.0(3)
C(1)#1-C(2)-C(1)-O(1)	-179.5(2)
C(3)-C(2)-C(1)-C(2)#1	-178.52(18)
C(1)#1-C(2)-C(1)-C(2)#1	0.0
C(2)-C(3)-C(5)-C(7)	55.00(17)
C(4)-C(3)-C(5)-C(7)	-126.82(15)
C(2)-C(3)-C(5)-C(8)	175.07(13)
C(4)-C(3)-C(5)-C(8)	-6.74(19)
C(2)-C(3)-C(5)-C(6)	-65.42(17)
C(4)-C(3)-C(5)-C(6)	112.76(15)

Symmetry transformations used to generate equivalent atoms: #1 -x,-y+2,-z+1

```
data_comp3a
audit creation method
                                 SHELXL-97
_chemical_name_systematic
 ?
_chemical_name_common
_chemical_melting_point
__chemical_formula moiety
__chemical_formula_sum
'C8 H12 O'
chemical formula weight
                                124.18
loop
 _atom_type symbol
 _atom_type_description
 _atom_type_scat_dispersion_real
 _atom_type_scat_dispersion_imag
 atom_type_scat_source
 _atom_type_scat_scares
'C' 'C' 0.0033 0.0016
 'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
 'H' 'H' 0.0000 0.0000
 'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
 '0' '0' 0.0106 0.0060
 'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
_symmetry_cell_setting
                                 Monoclinic
                                 P2(1)/n
symmetry space group name H-M
loop_
 symmetry equiv pos as xyz
 'x, y, z'
 -x+1/2, y+1/2, -z+1/2
 '-x, -y, -z'
 x-1/2, -y-1/2, z-1/2
cell length a
                                 6.829(2)
_cell_length b
                                 10.524(3)
_cell_length c
                                10.179(3)
cell angle alpha
                                90.00
cell angle beta
                                96.11(2)
cell angle gamma
                                 90.00
cell volume
                                 727.4(4)
cell formula units Z
cell measurement temperature
                                 188(2)
cell measurement reflns used
                                 26
cell measurement theta min
                                 4.45
cell measurement theta max
                                 7.75
_exptl_crystal_description
                                 block
_exptl_crystal_colour
                                 yellow
exptl crystal size max
                                 0.4
exptl crystal size mid
                                 0.35
_exptl_crystal_size min
                                 0.3
exptl crystal density meas
_exptl_crystal_density_diffrn
                                1.134
exptl crystal density method
                                'not measured'
exptl crystal F 000
                                 272
exptl absorpt coefficient mu
                                 0.073
exptl absorpt correction type
```

none

```
exptl absorpt correction T min
exptl_absorpt_correction_T_max
exptl absorpt process details
exptl special details
 ?
;
diffrn ambient temperature
                                  188(2)
diffrn radiation wavelength
                                   0.71073
diffrn radiation type
                                  MoK\a
_diffrn_radiation source
                                   'fine-focus sealed tube'
\_diffrn\_radiation\_monochromator
                                  graphite
_diffrn_measurement device type
                                   'Bruker P4'
_diffrn_measurement method
                                   '\w scans'
_diffrn_detector_area resol mean
_diffrn_standards number
_diffrn_standards_interval_count
                                  97
_diffrn_standards_interval_time
_diffrn_standards_decay %
_diffrn_reflns number
                                   3618
_diffrn_reflns_av_R equivalents
                                   0.0869
_diffrn_reflns_av_sigmaI/netI
                                   0.0801
_diffrn_reflns_limit_h_min
                                   - 8
_diffrn_reflns_limit_h_max
                                   8
_diffrn_reflns_limit_k_min
                                   -13
_diffrn_reflns_limit_k_max
                                  13
_diffrn_reflns_limit_l_min
                                  -13
_diffrn_reflns_limit_l_max
                                  13
_diffrn_reflns_theta min
                                  2.79
diffrn reflns theta max
                                  26.99
reflns number total
                                  1586
_reflns_number_gt
                                  1210
reflns threshold expression
                                  >2sigma(I)
_computing data collection
                                   'Bruker XSCANS'
                                   'Bruker XSCANS'
_computing_cell_refinement
computing data reduction
                                   'Bruker SHELXTL'
computing structure solution
                                   'SHELXS-97 (Sheldrick, 1990)'
                                   'SHELXL-97 (Sheldrick, 1997)'
computing structure refinement
computing molecular graphics
                                   'Bruker SHELXTL'
computing publication material
                                   'Bruker SHELXTL'
_refine_special details
 Refinement of F^2^ against ALL reflections. The weighted R-factor wR and
 goodness of fit S are based on F^2^, conventional R-factors R are based
 on F, with F set to zero for negative F^2^. The threshold expression of
 F^2 > 2sigma(F^2) is used only for calculating R-factors(gt) etc. and is
 not relevant to the choice of reflections for refinement. R-factors based
 on F^2^ are statistically about twice as large as those based on F, and R-
 factors based on ALL data will be even larger.
refine ls structure factor coef Fsqd
refine ls matrix type
                                   full
refine ls weighting scheme
                                  calc
_refine_ls_weighting_details
-\ calc \overline{w}=1/[\s^2(\overline{fo^2}^2)+(0.0408P)^2+0.0758P] where P=(Fo^2^+2Fc^2^)/3'
atom sites solution primary
                                  direct
atom sites solution secondary
                                  difmap
```

```
atom sites solution hydrogens
                                   qeom
_refine_ls_hydrogen_treatment
                                   mixed
refine ls extinction method
                                   none
refine ls extinction coef
refine ls number reflns
                                   1586
refine ls number parameters
                                   130
refine ls number restraints
                                   Λ
refine ls R factor all
                                   0.0632
refine ls_R_factor_gt
                                   0.0451
refine ls wR factor ref
                                   0.1129
refine ls wR factor gt
                                   0.1044
refine ls goodness of fit ref
                                   1.021
refine ls restrained S all
                                   1.021
_refine_ls_shift/su max
                                   0.000
_refine_ls shift/su mean
                                   0.000
loop
 _atom_site label
 _atom_site_type_symbol
 \_atom\_site\_fract x
 _atom_site_fract_y
 _atom_site_fract z
 _atom_site_U_iso_or_equiv
 _atom_site_adp_type
 _atom_site_occupancy
 _atom_site_symmetry_multiplicity
 _atom_site_calc_flag
 _atom_site_refinement_flags
 _atom_site_disorder_assembly
  atom site disorder group
O1 O 0.09723(16) 0.90135(12) 0.31910(9) 0.0365(3) Uani 1 1 d . . .
C3 C 0.27454(19) 1.14689(13) 0.48778(12) 0.0243(3) Uani 1 1 d . . .
C2 C 0.12080(19) 1.06594(14) 0.49831(12) 0.0246(3) Uani 1 1 d . . .
C1 C 0.04408(19) 0.95418(14) 0.41605(12) 0.0251(3) Uani 1 1 d . . .
C4 C 0.3964(2) 1.12747(18) 0.37524(16) 0.0337(4) Uani 1 1 d . . .
C5 C 0.3242(2) 1.25825(14) 0.58276(13) 0.0283(3) Uani 1 1 d . . .
C6 C 0.1531(2) 1.35491(17) 0.56481(18) 0.0372(4) Uani 1 1 d . . .
C7 C 0.3515(3) 1.20938(18) 0.72628(15) 0.0386(4) Uani 1 1 d . . .
H7C H 0.469(3) 1.154(2) 0.7385(17) 0.048(5) Uiso 1 1 d . . .
C8 C 0.5151(3) 1.3271(2) 0.55563(19) 0.0422(4) Uani 1 1 d . .
H7B H 0.234(3) 1.162(2) 0.7527(16) 0.045(5) Uiso 1 1 d . . .
H4C H 0.535(3) 1.115(2) 0.4072(16) 0.050(5) Uiso 1 1 d . . .
H4B H 0.352(3) 1.055(2) 0.3160(17) 0.056(6) Uiso 1 1 d . . .
H4A H 0.385(3) 1.206(2) 0.3174(17) 0.052(5) Uiso 1 1 d . . .
H7A H 0.385(3) 1.282(2) 0.7889(17) 0.057(6) Uiso 1 1 d . . .
H8A H 0.506(3) 1.358(2) 0.4638(19) 0.047(5) Uiso 1 1 d . . .
H8B H 0.543(3) 1.396(3) 0.6239(19) 0.064(6) Uiso 1 1 d . . .
H8C H 0.630(3) 1.266(3) 0.567(2) 0.078(7) Uiso 1 1 d . . .
H6A H 0.141(2) 1.387(2) 0.4717(17) 0.049(5) Uiso 1 1 d . . .
H6B H 0.028(3) 1.315(2) 0.5864(16) 0.051(5) Uiso 1 1 d . . .
H6C H 0.182(3) 1.425(2) 0.6296(18) 0.057(6) Uiso 1 1 d . . .
loop
 atom site aniso label
 atom site aniso U 11
 atom site aniso U 22
 atom site aniso U 33
 atom site aniso U 23
 atom site aniso U 13
  atom_site_aniso U 12
01 \ 0.0425(6) \ 0.0315(6) \ 0.0374(6) \ -0.0116(5) \ 0.0135(4) \ -0.0058(5)
 \texttt{C3} \ \ 0.0263(6) \ \ 0.0179(7) \ \ 0.0278(6) \ \ 0.0040(5) \ \ -0.0014(5) \ \ 0.0019(6)
```

```
C2 0.0285(6) 0.0192(7) 0.0258(6) -0.0021(5) 0.0006(5) 0.0017(6)
\mathtt{C1} \ \ 0.0268(6) \ \ 0.0203(7) \ \ 0.0278(6) \ \ -0.0004(6) \ \ 0.0010(5) \ \ 0.0009(6)
C4 0.0349(8) 0.0285(9) 0.0388(8) 0.0014(7) 0.0086(6) -0.0029(7)
C5 0.0305(7) 0.0212(7) 0.0324(7) -0.0011(6) -0.0015(5) -0.0032(6)
C6 0.0416(9) 0.0226(8) 0.0457(9) -0.0049(7) -0.0033(7) 0.0042(7)
 \texttt{C7} \ \ 0.0460(9) \ \ 0.0347(10) \ \ 0.0327(8) \ \ -0.0014(7) \ \ -0.0060(7) \ \ -0.0042(8) 
 \texttt{C8} \ \ 0.0401(9) \ \ 0.0349(10) \ \ 0.0509(10) \ \ -0.0045(8) \ \ 0.0019(7) \ \ -0.0126(8) 
_geom_special_details
All esds (except the esd in the dihedral angle between two l.s. planes)
 are estimated using the full covariance matrix. The cell esds are taken
 into account individually in the estimation of esds in distances, angles
 and torsion angles; correlations between esds in cell parameters are only
 used when they are defined by crystal symmetry. An approximate (isotropic)
 treatment of cell esds is used for estimating esds involving l.s. planes.
loop
 _geom_bond_atom_site label 1
 _geom_bond_atom_site_label_2
 _geom_bond_distance
_geom_bond_site_symmetry 2
 geom bond publ flag
\overline{01} C1 \overline{1.2211}(16) .
C3 C2 1.365(2) . ?
C3 C4 1.500(2) . ?
C3 C5 1.534(2) . ?
C2 C1 1.505(2) . ?
C2 C1 1.5105(18) 3 576 ?
C1 C2 1.5105(18) 3 576 ?
C4 H4C 0.979(19) . ?
C4 H4B 1.00(2) . ?
C4 H4A 1.01(2) . ?
C5 C7 1.541(2) . ?
C5 C8 1.542(2) . ?
C5 C6 1.545(2) . ?
C6 H6A 1.001(18) .
C6 H6B 0.99(2) . ?
C6 H6C 0.99(2) . ?
C7 H7C 0.99(2) . ?
C7 H7B 1.00(2) . ?
C7 H7A 1.00(2) . ?
C8 H8A 0.99(2) . ?
C8 H8B 1.01(2) . ?
C8 H8C 1.01(3) . ?
loop
 geom angle atom site label 1
 geom angle atom site label 2
 _geom_angle_atom_site_label 3
 _geom angle
 _geom_angle site symmetry 1
 geom angle site symmetry 3
  geom angle publ flag
C2 C3 C4 117.96(13) . . ?
C2 C3 C5 123.20(12) . . ?
C4 C3 C5 118.82(13) . . ?
C3 C2 C1 132.00(12) . . ?
C3 C2 C1 139.42(13) . 3 576 ?
C1 C2 C1 88.56(10) . 3 576 ?
```

O1 C1 C2 133.60(13) . . ?

16

```
O1 C1 C2 134.96(14) . 3 576 ?
C2 C1 C2 91.44(10) . 3_576 ?
C3 C4 H4C 111.2(10) . . ?
C3 C4 H4B 113.9(12) . . ?
H4C C4 H4B 108.7(18) . . ?
C3 C4 H4A 108.7(11) . . ?
H4C C4 H4A 108.2(17) . . ?
H4B C4 H4A 105.8(15) . . ?
C3 C5 C7 109.81(13) . . ?
C3 C5 C8 112.36(12) . . ?
C7 C5 C8 107.88(13) . . ?
C3 C5 C6 108.34(11) . .
C7 C5 C6 110.21(13) . .
C8 C5 C6 108.21(14) . . ?
C5 C6 H6A 108.7(11) . . ?
C5 C6 H6B 110.5(13) . . ?
H6A C6 H6B 111.2(14) . . ?
C5 C6 H6C 108.2(12) . . ?
H6A C6 H6C 111.7(17) . . ?
H6B C6 H6C 106.4(16) . . ?
C5 C7 H7C 109.3(10) . . ?
C5 C7 H7B 113.5(10) . . ?
H7C C7 H7B 109.8(17) . . ?
C5 C7 H7A 110.2(12) . . ?
H7C C7 H7A 104.0(16) . . ?
H7B C7 H7A 109.6(15) . . ?
C5 C8 H8A 110.7(11) . . ?
C5 C8 H8B 108.3(12) . . ?
H8A C8 H8B 113.8(19) . . ?
C5 C8 H8C 110.1(15) . . ?
H8A C8 H8C 106.8(16) . . ?
H8B C8 H8C 107.0(18) . . ?
loop
 _geom_torsion_atom_site_label_1
 _geom_torsion_atom_site_label_2
 _geom_torsion_atom_site_label_3
 _geom_torsion_atom_site_label_4
 geom torsion
 geom torsion site symmetry 1
 geom torsion site symmetry 2
 geom torsion site symmetry 3
 _geom_torsion_site symmetry 4
  geom torsion publ flag
C\overline{4} C3 \overline{C}2 C1 0.4(2) . . . ?
C5 C3 C2 C1 178.57(13) . . . . ?
C4 C3 C2 C1 -177.35(16) . . . 3 576 ?
C5 C3 C2 C1 0.8(3) . . . 3_576 ?
C3 C2 C1 O1 2.0(3) . . . . ?
C1 C2 C1 O1 -179.5(2) 3_576 . . . ?
C3 C2 C1 C2 -178.52(18) . . . 3 576 ?
C1 C2 C1 C2 0.0 3 576 . . 3 576 ?
C2 C3 C5 C7 55.00(17) . . . . ?
C4 C3 C5 C7 -126.82(15) . . . . ?
C2 C3 C5 C8 175.07(13) . . . . ?
C4 C3 C5 C8 -6.74(19) . . . . ?
C2 C3 C5 C6 -65.42(17) . . . . ?
C4 C3 C5 C6 112.76(15) . . . . ?
diffrn measured fraction theta max
                                        0.999
diffrn reflns theta full
                                        26.99
diffrn measured fraction theta full
                                        0.999
```

```
_refine_diff_density_max 0.308
_refine_diff_density_min -0.161
_refine_diff_density_rms 0.039
```