

1 Fitness tracking reveals task-specific associations  
2 between memory, mental health, and exercise

3 Jeremy R. Manning<sup>1, \*</sup>, Gina M. Notaro<sup>1,2</sup>, Esme Chen<sup>1</sup>, and Paxton C. Fitzpatrick<sup>1</sup>

4 <sup>1</sup>Dartmouth College, Hanover, NH

5 <sup>2</sup>Lockheed Martin, Bethesda, MD

6 \*Address correspondence to [jeremy.r.manning@dartmouth.edu](mailto:jeremy.r.manning@dartmouth.edu)

7 September 17, 2021

8 **Abstract**

9 Physical exercise can benefit both physical and mental well-being. Different forms of exercise  
10 (i.e., aerobic versus anaerobic; running versus walking versus swimming versus yoga; high-  
11 intensity interval training versus endurance workouts; etc.) impact physical fitness in different  
12 ways. For example, running may substantially impact leg and heart strength but only moderately  
13 impact arm strength. We hypothesized that the mental benefits of exercise might be similarly  
14 differentiated. We focused specifically on how different forms of exercise might related to different  
15 aspects of memory and mental health. To test our hypothesis, we collected nearly a century's  
16 worth of fitness data (in aggregate). We then asked participants to fill out surveys asking them  
17 to self-report on different aspects of their mental health. We also asked participants to engage in  
18 a battery of memory tasks that tested their short and long term episodic, semantic, and spatial  
19 memory. We found that participants with similar exercise habits and fitness profiles tended to  
20 also exhibit similar mental health and task performance profiles.

## 21 Introduction

22 Engaging in physical activity (exercise) can improve our physical fitness by increasing muscle  
23 strength (Crane et al., 2013; Knuttgen, 2007; Lindh, 1979; Rogers and Evans, 1993), increasing bone  
24 density (Bassey and Ramsdale, 1994; Chilibeck et al., 2012; Layne and Nelson, 1999), increasing  
25 cardiovascular performance (Maiorana et al., 2000; Pollock et al., 2000), increasing lung capac-  
26 ity (Lazovic-Popovic et al., 2016) (although see Roman et al., 2016), increasing endurance (Wilmore  
27 and Knuttgen, 2003), and more. Exercise can also improve mental health (Basso and Suzuki, 2017;  
28 Callaghan, 2004; Deslandes et al., 2009; Mikkelsen et al., 2017; Paluska and Schwenk, 2000; Raglin,  
29 1990; Taylor et al., 1985) and cognitive performance (Basso and Suzuki, 2017; Brisswalter et al.,  
30 2002; Chang et al., 2012; Etnier et al., 2006).

31 The physical benefits of exercise can be explained by stress-responses of the affected body tis-  
32 sues. For example, skeletal muscles that are taxed during exercise exhibit stress responses (Morton  
33 et al., 2009) that can in turn affect their growth or atrophy (Schiaffino et al., 2013). By comparison,  
34 the benefits of exercise on mental health are less direct. For example, one hypothesis is that ex-  
35 ercise leads to specific physiological changes, such as increased aminergic synaptic transmission  
36 and endorphin release, which in turn act on neurotransmitters in the brain (Paluska and Schwenk,  
37 2000).

38 Speculatively, if different exercise regimens lead to different neurophysiological responses, one  
39 might be able to map out a spectrum of signalling and transduction pathways that are impacted  
40 by a given type, duration, and intensity of exercise in each brain region. For example, prior work  
41 has shown that exercise increases acetylcholine levels, starting in the vicinity of the exercised  
42 muscles (Shoemaker et al., 1997). Acetylcholine is thought to play an important role in memory  
43 formation (Palacios-Filardo et al., 2021, e.g., by modulating specific synaptic inputs from entorhinal  
44 cortex to the hippocampus, albeit in rodents;). Given the central role of these medial temporal  
45 lobe structures play in memory, changes in acetylcholine might lead to specific changes in memory  
46 formation and retrieval.

47 In the present study, we hypothesize that (a) different exercise regimens will have different,

48 quantifiable impacts on cognitive performance and mental health, and that (b) these impacts will  
49 be consistent across individuals. To this end, we collected a year of fitness tracking data from  
50 each of 113 participants. We then asked each participant to fill out a brief survey in which they  
51 self-evaluated several aspects of their mental health. Finally, we ran each participant through a  
52 battery of memory tasks, which we used to evaluate their memory performance along several  
53 dimensions. We examined the data for potential associations between memory, mental health, and  
54 exercise.

## 55 Results

- 56 • exploratory analysis (correlations)
  - 57 – Memory-memory
  - 58 – fitness-fitness
  - 59 – survey-survey
  - 60 – (fitness + survey)-memory
- 61 • predictive analysis (regressions)
  - 62 – Predict memory performance on held-out task from other tasks
  - 63 – Predict memory performance on each task using fitness data
  - 64 – Predict memory performance on each task using survey data
- 65 • Reverse correlations: look at recent changes versus baseline trends
  - 66 – Fitness profile that predicts performance on each task (barplots + timelines)
  - 67 – Fitness profile for each survey demographic (barplots + timelines)
  - 68 \* Select out mental health demographics (based on meds, stress levels)

## 69 Discussion

- 70 • summarize key findings
- 71 • correlation versus causation
- 72 • what can vs. can't we know? we can identify correlations, but not causal direction– e.g. we  
73 cannot know whether exercise *causes* mental changes versus whether people with particular  
74 neural profiles might tend to engage in particular exercise behaviors. that being said, we *can*  
75 separate out baseline tendencies (e.g., how people tend to exercise in general) versus recent  
76 changes (e.g., how they happened to have exercised prior to the experiment).
- 77 • related work (exercise/memory, exercise/mental health), what this study adds
- 78 • future direction: towards customized physical exercise recommendation engine for optimiz-  
79 ing mental health and mental fitness

## 80 Methods

### 81 Experiment

#### 82 Participants

#### 83 Tasks

#### 84 Intake survey.

#### 85 Free recall.

#### 86 Naturalistic recall.

#### 87 Foreign language flashcards.

#### 88 Spatial learning.

89 **Fitness tracking using Fitbit devices**

90 **Processing Fitbit data**

91 **Raw metrics.**

92 **Comparing recent versus baseline measurements.**

93 **Exploratory correlation analyses**

94 **Imputation and interpolation of missing data.**

95 **Regression-based prediction analyses**

96 **Reverse correlation analyses**

## 97 **Acknowledgements**

98 We acknowledge useful discussions with David Bucci, Emily Glasser, Andrew Heusser, Avigail  
99 Bartolome, Lorie Loeb, Lucy Owen, and Kirsten Ziman. Our work was supported in part by  
100 the Dartmouth Young Minds and Brains initiative. The content is solely the responsibility of the  
101 authors and does not necessarily represent the official views of our supporting organizations. This  
102 paper is dedicated to the memory of David Bucci, who helped to inspire the theoretical foundations  
103 of this work. Dave served as a mentor and colleague on the project prior to his passing.

## 104 **Data and code availability**

105 All analysis code and data used in the present manuscript may be found [here](#).

## 106 **Author contributions**

107 Concept: J.R.M. Implementation: G.M.N. Analyses: G.M.N., E.C., and P.C.F. Writing: J.R.M.

## 108 **Competing interests**

109 The authors declare no competing interests.

## 110 **References**

- 111 Bassey, E. J. and Ramsdale, S. J. (1994). Increase in femoral bone density in young women following  
112 high-impact exercise. *Osteoporosis International*, 4:72–75.
- 113 Basso, J. C. and Suzuki, W. A. (2017). The effects of acute exercise on mood, cognition, neurophys-  
114 iology, and neurochemical pathways: a review. *Brain Plasticity*, 2(2):127–152.
- 115 Brisswalter, J., Collardeau, M., and René, A. (2002). Effects of acute physical exercise characteristics  
116 on cognitive performance. *Sports Medicine*, 32:555–566.
- 117 Callaghan, P. (2004). Exercise: a neglected intervention in mental health care? *Psychiatric and*  
118 *Mental Health Nursing*, 11(4):476–483.
- 119 Chang, Y. K., Labban, J. D., Gapin, J. I., and Etnier, J. L. (2012). The effects of acute exercise on  
120 cognitive performance: a meta-analysis. *Brain Research*, 1453:87–101.
- 121 Chilibeck, P. D., Sale, D. G., and Webber, C. E. (2012). Exercise and bone mineral density. *Sports*  
122 *Medicine*, 19:103–122.
- 123 Crane, J. D., MacNeil, L. G., and Tarnopolsky, M. A. (2013). Long-term aerobic exercise is associated  
124 with greater muscle strength throughout the life span. *The Journals of Gerontology: Series A*,  
125 68(6):631–638.
- 126 Deslandes, A., Moraes, H., Ferreira, C., Veiga, H., Silveira, H., Mouta, R., Pompeu, F. A. M. S.,  
127 Coutinho, E. S. F., and Laks, J. (2009). Exercise and mental health: many reasons to move.  
128 *Neuropsychobiology*, 59:191–198.

- 129 Etnier, J. L., Nowell, P. M., Landers, D. M., and Sibley, B. A. (2006). A meta-regression to examine the  
130 relationship between aerobic fitness and cognitive performance. *Brain Research: Brain Research*  
131 *Reviews*, 52(1):119–130.
- 132 Knuttgen, H. G. (2007). Strength training and aerobic exercise: comparison and contrast. *Journal of*  
133 *Strength and Conditioning Research*, 21(3):973–978.
- 134 Layne, J. E. and Nelson, M. E. (1999). The effects of progressive resistance training on bone density:  
135 a review. *Medicine and Science in Sports and Exercise*, 31(1):25–30.
- 136 Lazovic-Popovic, B., Zlatkovic-Svenda, M., Durmic, T., Djelic, M., Saranovic, D., and Zugic, V.  
137 (2016). Superior lung capacity in swimmers: some questions, more answers! *Revista Portuguesa*  
138 *de Pneumologia*, 22(3):151–156.
- 139 Lindh, M. (1979). Increase of muscle strength from isometric quadriceps exercises at different knee  
140 angles. *Scandinavian Journal of Rehabilitation Medicine*, 11(1):33–36.
- 141 Maiorana, A., O'Driscoll, G., Cheetham, C., Collis, J., Goodman, C., Rankin, S., Taylor, R., and  
142 Green, D. (2000). Combined aerobic and resistance exercise training improves functional capacity  
143 and strength in CHF. *Journal of Applied Physiology*, 88(1565–1570).
- 144 Mikkelsen, K., Stojanovska, L., Polenakovic, M., Bosevski, M., and Apostolopoulos, V. (2017).  
145 Exercise and mental health. *Maturitas*, 106:48–56.
- 146 Morton, J. P., Kayani, A. C., McArdle, A., and Drust, B. (2009). The exercise-induced stress response  
147 of skeletal muscle, with specific emphasis on humans. *Sports Medicine*, 39:643–662.
- 148 Palacios-Filardo, J., Udakis, M., Brown, G. A., Tehan, B. G., Congreve, M. S., Nathan, P. J., Brown, A.  
149 J. H., and Mellor, J. R. (2021). Acetylcholine prioritises direct synaptic inputs from entorhinal cor-  
150 tex to CA1 by differential modulation of feedforward inhibitory circuits. *Nature Communications*,  
151 12(5475):doi.org/10.1038/s41467-021-25280-5.
- 152 Paluska, S. A. and Schwenk, T. L. (2000). Physical activity and mental health. *Sports Medicine*,  
153 29(3):167–180.

154 Pollock, M. L., Franklin, B. A., Balady, G. J., Chaltman, B. L., Fleg, J. L., Fletcher, B., Limacher, M.,  
 155 na, I. L. P., Stein, R. A., Williams, M., and Bazzarre, T. (2000). Resistance exercise in individuals  
 156 with and without cardiovascular disease. *Circulation*, 101:828–833.

157 Raglin, J. S. (1990). Exercise and mental health. *Sports Medicine*, 9:323–329.

158 Rogers, M. A. and Evans, W. J. (1993). Changes in skeletal muscle with aging: effects of exercise  
 159 training. *Exercise and Sport Sciences Reviews*, 21:65–102.

160 Roman, M. A., Rossiter, H. B., and Casaburi, R. (2016). Exercise, ageing and the lung. *European*  
 161 *Respiratory Journal*, 48:1471–1486.

162 Schiaffino, S., Dyar, K. A., Ciciliot, S., Blaauw, B., and Sandri, M. (2013). Mechanisms regulating  
 163 skeletal muscle growth and atrophy. *The febs Journal*, 280(17):4294–4314.

164 Shoemaker, J. K., Halliwill, J. R., Hughson, R. L., and Joyner, M. J. (1997). Contributions of  
 165 acetylcholine and nitric oxide to forearm blood flow at exercise onset and recovery. *Vascular*  
 166 *Physiology*, 273(5):2388–2395.

167 Taylor, C. B., Sallis, J. F., and Needle, R. (1985). The relation of physical activity and exercise to  
 168 mental health. *Public Health Reports*, 100(2):195–202.

169 Wilmore, J. H. and Knuttgen, H. G. (2003). Aerobic exercise and endurance. *The Physician and*  
 170 *Sportsmedicine*, 31(5):45–51.