



Question: Can you identify the abnormality on this plain film of the pelvis?

<image 1>

Options:

- (A) Bone cyst
- (B) Acute hip fracture
- (C) Osteoarthritis
- (D) Surgical hardware
- (E) Resection of the pubic symphysis
- (F) Bone tumor
- (G) Hip dislocation
- (H) Osteoporosis
- (I) Paget's disease
- (J) Bone infection

Explanation: The key finding is a smooth cortical bone defect at the pubic symphysis, suggesting a previous pubic symphysis resection rather than acute separation or trauma. The sacroiliac and hip joints are well-matched, with no fracture lines or metal implants, and no signs of osteoarthritis, Paget's disease, osteoporosis, infection, or tumor.

Qwen3-VL-8B-Thinking: <think> The video shows a diffuse radiolucent area in the pelvis, accompanied by widespread cortical thinning and loss of normal trabecular detail. The hip joint is symmetrical, without displacement, and no focal osteolytic or sclerotic lesions, fracture lines, periosteal reaction, or metallic density shadows are observed. The joint space is intact, with no osteophyte formation, suggesting the absence of osteoarthritis, and lacks the typical features of Paget's disease (bone expansion, cortical thickening) or infection (invasive destruction, soft tissue swelling). The lesion is **diffuse rather than focal**, therefore osteoporosis is the most plausible explanation. </think><answer>H</answer>

Error Reason: The model incorrectly overinterpreted diffuse radiolucent areas as osteoporosis, neglecting a crucial finding—smooth cortical spaces at the pubic symphysis, consistent with surgical resection. The apparent "demineralization" may be related to bone protrusion/exposure, and there is no corroborating evidence of osteopenia prevalent in other sites.

OneThinker: <think> Based on the pelvis plain film and the listed options, the abnormality most plausibly involves **the hip joint**. Typical signs of a bone cyst, tumor, osteoporosis, Paget's disease, osteoarthritis, pubic symphysis resection, hip dislocation, or bone infection are not clearly demonstrated, and an acute hip fracture is not convincingly visible. With the absence of definitive features for these alternatives, the most specific and common explanation in this context is a hip-related abnormality consistent with surgical hardware. Therefore, the best answer is D. Surgical hardware. </think><answer>D</answer>

Error Reason: Although no metallic density was observed on the imaging, the model incorrectly inferred the presence of a "surgical implant" and overlooked the smooth pubic symphysis and cortical absence—characteristic features of previous resections. Furthermore, the diagnosis incorrectly described the abnormality as related to the hip joint, rather than located at the pubic symphysis.

Qwen3-VL-8B-Instruct (SFT+RL on VideoKR): <think> The video shows **significant destruction at the pubic symphysis**, with a marked gap or separation, which is uncommon in a normal pelvis. Therefore, there is a significant abnormality in the pubic symphysis region. The **interpubic cortical space is smooth**, and the surrounding bone structure appears intact, **but this is consistent with previous surgical resection rather than acute trauma**. The hip joint is well-aligned, without displacement or dislocation; no cortical fracture or trabecular bone destruction is observed, nor is there an acute hip fracture; no focal osteolytic/sclerotic lesions, bone cysts, tumors, Paget's disease, or infection are seen. Diffuse demineralization suggests osteoporosis, but is insufficient for a definitive diagnosis; typical features of osteoarthritis such as joint space narrowing or osteophyte formation are also absent. No metallic feel or surgical implants are observed. Therefore, the pathological findings are most consistent with a diagnosis following pubic symphysis resection. </think><answer>E</answer>

Note: VideoKR focuses on the pubic symphysis and utilizes radiological prior information (depicting the pelvic ring and checking joint consistency) to avoid the biases of other models that are hip-centric and diffuse osteoporosis.