

Question: By how much does the electronic energy gap of NbO2 widen as it transitions from a bond alternation level of 17.1% to its completely dimerized insulating form?

GT Answer: 0.26 eV

Crucial Evidence:

The occupied split-off flat bands can also exist in nonmagnetic *d*-electron transition metal oxides such as the tetragonal compound NbO₂. This compound shows a metal-insulator transition with an embedded Peierls instability [82], where the Nb-Nb dimerization of structural symmetry breaking determines the electronic structures. Without Nb-Nb dimerization, as shown in Fig. 4(a), there is an internal energy gap, and the Fermi level resides in the principal CB, which makes the system an electron metal. The space group is *P4₂/mmn* (rutile structure) for this metallic phase. With Nb-Nb dimerization, as shown in Fig. 4(b), the Fermi level is shifted down, entering the internal energy gap, creating occupied split-off flat bands, and opening a band gap of 0.40 eV. The space group is *I4₁* for this body-centered insulating phase. The total energy will be lowered by 64 meV/f.u. after symmetry breaking. The band edges change from (*p*, *d*^{*}) to (*d*, *d*^{*}) during this C2 process, reflecting that *d*-like resonance states within the principal CB shift into the internal energy gap, as also shown in the PDOS of Fig. 4(b). The split-off flat bands will trap one Nb-*d* electron per formula unit in the Nb-Nb dimers, as shown in Fig. 5(a). (See the Supplemental Material [54] for the occupied split-off flat bands trapping electrons in LaTiO₃.)

where $d_{\text{Nb-Nb}}^{\text{long}}$ and $d_{\text{Nb-Nb}}^{\text{short}}$ are the long and short Nb-Nb bonding lengths, respectively. The snapshots of crystallographic structures at different DBAs are generated by linear interpolations between the initial symmetry-unbroken phase (DBA = 0%) and the final symmetry-broken phase (DBA = 21.3%), which are relaxed by DFT calculations with the electronic structures shown in Fig. 4. We pick up four representative snapshots of density of states (DOS) for NbO₂ in Figs. 8(a)–8(d) to show the evolution of occupied split-off flat bands: The Nb-*d* resonance states reside within the broad principal CB at DBA = 4.3% [Fig. 8(a)]; gradually shift down, still connecting the principal CB at DBA = 15.0% [Fig. 8(b)]; enter the internal energy gap, becoming occupied split-off flat bands, producing a band gap of 0.14 eV at DBA = 17.1% [Fig. 8(c)]; and further move away from the principal CB with a larger band gap of 0.34 eV at DBA = 20.3% [Fig. 8(d)]. We can see in Fig. 8(e) the whole C2 process with the band gap as a function of DBA: The band gap remains closed with increasing perturbations of DBA, starts to appear until exceeding minimal threshold perturbations of DBA = 15.1%, and then grows as a linear function of DBA. Note that we do not use the effective *U* for calculations of nonmagnetic NbO₂. This continuous C2 process is also associated with

Gemini-2.5-Pro Response:

According to the document, the electronic energy gap of NbO2 at a bond alternation (DBA) level of 17.1% is 0.14 eV.

The completely dimerized insulating form of NbO2 has a band gap of 0.40 eV.

Therefore, the widening of the electronic energy gap is the difference between these two values:
0.40 eV - 0.14 eV = 0.26 eV.

metry-broken phase (DBA = 21.3%), which
FT calculations with the electronic structures
4. We pick up four representative snapshots
ates (DOS) for NbO₂ in Figs. 8(a)–8(d) to
ion of occupied split-off flat bands: The Nb-
tes reside within the broad principal CB at
Fig. 8(a)]; gradually shift down, still connect-
l CB at DBA = 15.0% [Fig. 8(b)]; enter the
gap, becoming occupied split-off flat bands,
d gap of 0.14 eV at DBA = 17.1% [Fig. 8(c)];
e away from the principal CB with a larger


compound NbO₂. This compound shows a transition with an embedded Peierls instability Nb-Nb dimerization of structural symmetry l mines the electronic structures. Without Nb-N as shown in Fig. 4(a), there is an internal ei the Fermi level resides in the principal CB, the system an electron metal. The space grov (rutile structure) for this metallic phase. With ization, as shown in Fig. 4(b), the Fermi level i entering the internal energy gap, 5 ating occ flat bands, and opening a band gap of 0.40 group is *I4₁* for this body-centered insulati total energy will be lowered by 64 meV/f.u. ; breaking. The band edges change from (*p*, during this C2 process, reflecting that *d*-like r within the principal CB shift into the internal also shown in the PDOS of Fig. 4(b). The spli will trap one Nb-*d* electron per formula unit dimers, as shown in Fig. 5(a). (See the Suppl [54] for the occupied split-off flat bands trap in LaTiO₃.)

The cation-cation dimerization can also prc split-off flat bands in the metal insulator tra

Evaluation: Rec.=0.5, Rel.=4, Ans.=5, SAA=1 

Qwen3-VL-8B Response:

The electronic energy gap of NbO₂ widens by 0.2 eV as it transitions from a bond alternation level of 17.1% to its completely dimerized insulating form. This is determined by analyzing Figure 8 on page 14, which shows the evolution of the electronic structure of NbO₂ with increasing bond alternation (DBA). The figure includes a plot of the band gap as a function of DBA. At DBA = 17.1% (Figure 8c), the band gap is 0.34 eV. At the completely dimerized state, which corresponds to DBA = 20.3% (Figure 8d), the band gap is 0.54 eV. The difference is 0.54 eV - 0.34 eV = 0.20 eV.



Evaluation: Rec.=0, Rel.=1, Ans.=1, SAA=0 