

Extended data for: Towards human Super EEG?

Lucy L. W. Owen, Andrew C. Heusser, and Jeremy R. Manning
Department of Psychological and Brain Sciences
Dartmouth College, Hanover, NH 03755, USA
Corresponding author: jeremy.r.manning@dartmouth.edu

October 9, 2018

Overview

This document provides additional details about the methods we used in the main text. We also include some additional analyses referenced in the main text.




Figure S1: Reconstruction quality for datasets 1 and 2. A. Distributions of correlation coefficients. Although patients in dataset 1 all participated in the same experiment, we did not limit our analyses to task specific data. Instead we incorporated data from their entire recording session (sometimes up to 14 hours of data) and classify this as training our model on across experiment information. The split violin plot shows the distributions of reconstruction accuracy using models trained on data from all other patients (Across Subject, in black) and all other electrodes from the same patient (Within Subject, in gray). **B. Distributions of correlation coefficients.** Since all patients participated in two experiments in dataset 2, we limit our analyses to task specific data. The violin plots compare distributions of reconstruction accuracy using data from the same task (Within Experiment), the other task (Across Experiment), and from both tasks (All Experiment), using models trained on data from all other patients (Across Subject, in black) and all other electrodes from the same patient (Within Subject, in gray).




Figure S2: Reconstruction quality for experiments 1 and 2. A. Distributions of correlation coefficients. For experiment 1, the split violin plots compare distributions of reconstruction accuracy using data from just experiment 1 (Within Experiment), just experiment 2 (Across Experiment), and from both experiments (All Experiment), using models trained on data from all other patients (Across Subject, in black) and all other electrodes from the same patient (Within Subject, in gray). **B. Distributions of correlation coefficients.** For experiment 2, the split violin plots compare distributions of reconstruction accuracy using data from just experiment 2 (Within Experiment), just experiment 1 (Across Experiment), and from both experiments (All Experiment), using models trained on data from all other patients (Across Subject, in black) and all other electrodes from the same patient (Within Subject, in gray).




Figure S3: Reconstruction quality for each experiment. A. & C. Distributions of correlation coefficients. Across all electrodes from all patients in the labeled experiment from dataset 2, the panel displays the distribution of correlations between the observed and reconstructed LFP data using models trained on data from all other patients (Across, in black) and all other electrodes from the same patient (Within, in gray). **B. & D. Correlation maps.** The glass brain maps display the average correlation between the observed LFP data and the across-subjects model reconstructed data by location, for each labeled experiment.