tenet DIAGNOSTICS
It's Good to Know

MC-2977

PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE

Name	: **MRS.NAGA LAXMI J**	TID/SID	: UMR2162620/ 30644477
Age / Gender	: 70 Years / Female	Registered on	: 07-Nov-2025 / 07:58 AM
Ref.By	: DR RAJASHEKAR CHAKRAVATHY M	Collected on	: 07-Nov-2025 / 08:01 AM
Req.No	: 25BNJH0012386	Reported on	: 07-Nov-2025 / 11:52 AM
Sample Type	: Serum		

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Liver Function Test (LFT)

Investigation	Observed Value	Biological Reference Interval
Total Bilirubin. Method:Diazo Method	0.38	<1.2 mg/dL
Direct Bilirubin. Method:Diazo Method	0.17	<0.30 mg/dL
Indirect Bilirubin. Method:Calculated	0.21	<0.9 mg/dL
Alanine Aminotransferase ,(ALT/SGPT) Method:UV wtihout P5P	20	<34 U/L
Aspartate Aminotransferase,(AST/SGOT) Method:UV wtihout P5P	**37**	<31 U/L
ALP (Alkaline Phosphatase). Method:PNPP-AMP Buffer	**155**	35-104 U/L
Gamma GT. Method:GCNA	**64**	6-42 U/L
Total Protein. Method:Biuret	7.4	6.6-8.7 g/dL
Albumin. Method:Bromocresol Green (BCG)	3.8	3.5-5.2 g/dL
Globulin. Method:Calculated	3.60	1.8-3.8 g/dL
A/GRatio. Method:Calculated	1.06	0.8-2.0
AST/ALT Ratio Method:Calculated	**1.85**	<1.00
Note	Kindly correlate clinically	

Interpretation: Liver functions tests help to identify liver disease, its severity, and its type. Generally these tests are performed in combination, are abnormal in liver disease, and the pattern of abnormality is indicative of the nature of liver disease. An isolated abnormality of a single liver function test usually means a non-hepatic cause. If several liver function tests are simultaneously abnormal, then hepatic etiology is likely.

* Sample processed at National Reference Laboratory,Tenet Diagnostics,54, Kineta Towers, Journalist Colony, Banjara Hills

--- End Of Report ---

Dr.Abdur Rehman Asif
Consultant Biochemist
Reg.No - APMC/FMR/78102