tenet DIAGNOSTICS
It's Good to Know

PLEASE SCAN QR CODE
TO VERIFY THE REPORT ONLINE

Name	: **MRS.NAGA LAXMI J**	TID/SID	: UMR2162620/ 30691307
Age / Gender	: 71 Years / Female	Registered on	: 14-Nov-2025 / 10:30 AM
Ref.By	: DR RAJASEKARA CHAKRAVARTHI M	Collected on	: 14-Nov-2025 / 10:32 AM
Req.No	: 25BNJH0013256	Reported on	: 14-Nov-2025 / 17:17 PM
Sample Type	: Urine		

TEST REPORT

DEPARTMENT OF CLINICAL CHEMISTRY I

Protein / Creatinine Ratio, Spot Urine

Investigation	Observed Value	Biological Reference Interval
Protein Spot Urine Method:Benzethonium	**22.3**	<15 mg/dL
Creatinine Spot Urine Method:Alkaline Picrate Kinetic	108	25-400 mg/dL
Protein/Creatinine Ratio (Spot Urine) Method:Calculated	**0.21**	Normal: <0.20 Low grade proteinuria: 0.20-1.00 Moderate proteinuria: 1.00-5.00 Nephrosis: >5.00
Note	Kindly correlate clinically	

Comments:
Under most circumstances untimed (Spot) sample be used to detect and monitor Proteinuria in children and adults. Patients with 2 or more positive results within a period of 1-2 weeks should be labeled as having persistent proteinuria and undergo further evaluation an management for chronic kidney disease. Orthostatic proteinuria must be excluded by repeat measurement on first morning specimen if the initial findings of proteinuria was obtained on a random specimen.

* Sample processed at National Reference Laboratory,Tenet Diagnostics,54, Kineta Towers, Journalist Colony, Banjara Hills

--- End Of Report ---

Dr.Abdur Rehman Asif
Consultant Biochemist
Reg.No - APMC/FMR/78102