

Natural language statement

Domain

GroundTruth

Theorem 2.3.1. Let Ω be a regular bounded domain. Assume that for any x in Ω there exists a function $h(x, \cdot)$ in $C^2(\Omega)$ such that $\Delta_y h(x, y) = 0$ in Ω , $h(x, y) = K(x, y)$ for y in $\partial\Omega$. Define the Green's function $G(x, y) = K(x, y) - h(x, y)$ so that, in particular, $\Delta_y G(x, y) = 0$ in Ω , $G(x, y) = 0$ for y in $\partial\Omega$. Then for any $C^2(\Omega)$ -solution u of the Dirichlet problem (2.3.1) and x in Ω we have $u(x) = \int_{\Omega} G(x, y) \Delta_y u(y) dy + \int_{\partial\Omega} G(x, y) \frac{\partial u}{\partial n_y}(y) dS_y$, where $H(x, y) := -\Delta_y G(x, y)$ in Ω , y in $\partial\Omega$, is the so-called Poisson kernel.

PDE

```

noncomputable def directionalDerivativeList {d : N} :
  List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R
| [], u => u
| i :: indices, u => fun x =>
  fderiv R (directionalDerivativeList indices u) x (Pi.single i 1)

noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> N)) :
  List (Fin d) :=
  Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i

noncomputable def multiDerivative {d : N} (alpha : (Fin d -> N))
  (u : (Fin d -> R) -> R) : (Fin d -> R) -> R :=
  directionalDerivativeList (multiIndexDirections alpha) u

noncomputable def laplacian {d : N} (u : (Fin d -> R) -> R) (x : (Fin d ->
R)) : R :=
  sum i, directionalDerivativeList [i, i] u x

def RegularBoundedDomain {d : N} (Omega : Set (Fin d -> R)) : Prop :=
  IsOpen Omega /\ Bornology.IsBounded Omega /\ Omega.Nonempty /\
  forall z in frontier Omega, exists barrier : (Fin d -> R) -> R,
    ContinuousOn barrier (closure Omega) /\
    barrier z = 0 /\ (forall x in closure Omega, x != z -> 0 < barrier x)
    /\
    forall x in Omega, laplacian barrier x <= 0

def LaplaceDirichletSolution {d : N} (Omega : Set (Fin d -> R))
  (f g u : (Fin d -> R) -> R) : Prop :=
  ContDiffOn R 2 u Omega /\ ContinuousOn u (closure Omega) /\
  (forall x in Omega, laplacian u x = f x) /\ forall x in frontier Omega,
  u x = g x

theorem krylov_2_3_1_green_poisson_representation
  {d : N} {Omega : Set (Fin d -> R)}
  {K h G H : (Fin d -> R) -> (Fin d -> R) -> R} {f g u : (Fin d -> R) ->
R}
  (hOmega : RegularBoundedDomain Omega) (boundaryMeasure : Measure (Fin d
-> R))
  (hmeasure : boundaryMeasure (frontier Omega)^c = 0)
  (normal : (Fin d -> R) -> (Fin d -> R))
  (hnormal : forall y in frontier Omega, ||normal y|| = 1)
  (hharmonic : forall x in Omega, HarmonicIn Omega (h x))
  (hboundary : forall x in Omega, forall y in frontier Omega, h x y = K x
y)
  (hgreen : forall x y, G x y = K x y - h x y)
  (hgreenHarmonic : forall x in Omega, HarmonicIn (Omega \ {x}) (G x))
  (hgreenBoundary : forall x in Omega, forall y in frontier Omega, G x y
= 0)
  (hpoisson : forall x in Omega, forall y in frontier Omega,
  H x y = -fderiv R (G x) y (normal y))
  (hu : LaplaceDirichletSolution Omega f g u) :
  forall x in Omega, u x = integral y in Omega, G x y * f y +
  integral y, H x y * g y dboundaryMeasure := by
  sorry

```

Theorem 2.5.2. Let Ω be a domain and u in $C^2_{loc}(\Omega)$. Inter $C(\Omega)$ be a harmonic function in Ω . Then u is infinitely differentiable in Ω and for any multi-index α and any x in Ω we have $|D^\alpha u(x)| \leq N R^{|\alpha|} \sup_{B_R(x)} |u|$ whenever $B_R(x) \subset \Omega$.

PDE

```

noncomputable def directionalDerivativeList {d : N} :
  List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R
| [], u => u
| i :: indices, u => fun x =>
  fderiv R (directionalDerivativeList indices u) x (Pi.single i 1)

noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> N)) :
  List (Fin d) :=
  Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i

noncomputable def multiDerivative {d : N} (alpha : (Fin d -> N))
  (u : (Fin d -> R) -> R) : (Fin d -> R) -> R :=
  directionalDerivativeList (multiIndexDirections alpha) u

noncomputable def laplacian {d : N} (u : (Fin d -> R) -> R) (x : (Fin d ->
R)) : R :=

```

Natural language statement

Domain

GroundTruth

		<pre> sum i, directionalDerivativeList [i, i] u x def HarmonicIn {d : N} (Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : Prop := ContDiffOn R 2 u Omega /\ forall x in Omega, laplacian u x = 0 theorem krylov_2_5_2_harmonic_smooth_interior_estimates {d : N} {Omega : Set (Fin d -> R)} {u : (Fin d -> R) -> R} (hOmega : IsOpen Omega /\ IsConnected Omega) (hu : HarmonicIn Omega u) : ContDiffOn R top u Omega /\ forall alpha : (Fin d -> N), forall x in Omega, exists C : R, 0 <= C /\ forall R : R, 0 < R -> Metric.closedBall x R subset Omega -> multiDerivative alpha u x <= C * R ^ (-(sum i, alpha i : Z)) * sSup { u y y in Metric.closedBall x R} := by sorry </pre>
Theorem 2.9.2. Let Ω be a domain in $\mathbb{R}^d$ and u be a bounded and continuous function on Ω and $u = 0$ on $\partial\Omega$ if $\partial\Omega$ is empty (that is, if $\Omega \neq \mathbb{R}^d$). Moreover, assume that u is in $C^2_{loc}(\Omega)$. Finally, let $a(x)$, $b(x)$ be bounded and $c(x) \leq -\lambda$ where the constant $\lambda > 0$. Then in Ω, $u \leq \lambda^{-1} \sup_{\Omega} (Lu)^{-}$ and $u \leq \lambda^{-1} \sup_{\Omega} Lu$.	PDE	<pre> noncomputable def directionalDerivativeList {d : N} : List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R [], u => u i :: indices, u => fun x => fderiv R (directionalDerivativeList indices u) x (Pi.single i 1) noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> N)) : List (Fin d) := Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i noncomputable def multiDerivative {d : N} (alpha : (Fin d -> N)) (u : (Fin d -> R) -> R) : (Fin d -> R) -> R := directionalDerivativeList (multiIndexDirections alpha) u structure EllipticOperatorData {d : N} (m : N) (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) where terms : Finset ((Fin d -> N)) order_le : forall alpha in terms, sum i, alpha i <= m coefficient : (Fin d -> N) -> (Fin d -> R) -> R formula : forall u x, L u x = sum alpha in terms, coefficient alpha x * multiDerivative alpha u x ellipticityConstant : R ellipticityConstant_pos : 0 < ellipticityConstant principalSymbol : forall (x xi : (Fin d -> R)), ellipticityConstant * xi ^ m <= sum alpha in terms with sum i, alpha i = m, coefficient alpha x * prod i, (xi i) ^ (alpha i) def SecondOrderEllipticOperator {d : N} (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) (lam : R) : Prop := exists data : EllipticOperatorData 2 L, exists zeroIndex in data.terms, (forall i, zeroIndex i = 0) /\ (forall x, data.coefficient zeroIndex x <= -lam) /\ exists C : R, forall alpha in data.terms, forall x, data.coefficient alpha x <= C noncomputable def functionSupNorm {X : Type*} (Omega : Set X) (u : X -> R) : R:=0inf := x : Omega, ENNReal.ofReal u x theorem krylov_2_9_2_bounded_maximum_principle_resolvent {d : N} {Omega : Set (Fin d -> R)} {L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R} {lam : R} {u : (Fin d -> R) -> R} (hOmega : IsOpen Omega) (hDiff : ContDiffOn R 2 u Omega) (huContinuous : ContinuousOn u (closure Omega)) (huBounded : Bornology.IsBounded (u '' Omega)) (huBoundary : forall x in frontier Omega, u x = 0) (hlam : 0 < lam) (hL : SecondOrderEllipticOperator L lam) : functionSupNorm Omega (fun x => max (u x) 0) <= (ENNReal.ofReal lam)^-1 * functionSupNorm Omega (fun x => max (-(L u x)) 0) /\ functionSupNorm Omega u <= (ENNReal.ofReal lam)^-1 * functionSupNorm Omega (L u) := by </pre>

Natural language statement

Domain

GroundTruth

		sorry
Theorem 3.7.2. Let $\lambda \neq 0$, $k \geq 0$ be an integer and $0 < \delta < 1$. Then for any $f \in C^{k+\delta}(\mathbb{R}^d)$ there exists a unique solution $u \in C^{k+m+\delta}(\mathbb{R}^d)$ of the equation $L_\lambda u(x) = f(x)$, $x \in \mathbb{R}^d$.	PDE	<pre> noncomputable def directionalDerivativeList {d : N} : List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R [], u => u i :: indices, u => fun x => fderiv R (directionalDerivativeList indices u) x (Pi.single i 1) noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> N)) : List (Fin d) := Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i noncomputable def multiDerivative {d : N} (alpha : (Fin d -> N)) (u : (Fin d -> R) -> R) : (Fin d -> R) -> R := directionalDerivativeList (multiIndexDirections alpha) u noncomputable def holderGauge {d : N} (k : N) (delta : R) (Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : R >= 0 := (alpha : {alpha : (Fin d -> N) // sum i, alpha i <= k}, x : Omega, ENNReal.ofReal multiDerivative alpha u x) alpha : {alpha : (Fin d -> N) // sum i, alpha i = k}, x : Omega, y : Omega, (_ : (x : (Fin d -> R)) != y), ENNReal.ofReal (multiDerivative alpha u x - multiDerivative alpha u y / (x : (Fin d -> R)) - y ^ delta) def HolderOn {d : N} (r : R) (Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : Prop := exists k : N, exists delta : R, r = k + delta /\ 0 <= delta /\ delta < 1 /\ ContDiffOn R k u Omega /\ holderGauge k delta Omega u < inf structure EllipticOperatorData {d : N} (m : N) (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) where terms : Finset ((Fin d -> N)) order_le : forall alpha in terms, sum i, alpha i <= m coefficient : (Fin d -> N) -> (Fin d -> R) -> R formula : forall u x, L u x = sum alpha in terms, coefficient alpha x * multiDerivative alpha u x ellipticityConstant : R ellipticityConstant_pos : 0 < ellipticityConstant principalSymbol : forall (x xi : (Fin d -> R)), ellipticityConstant * xi ^ m <= sum alpha in terms with sum i, alpha i = m, coefficient alpha x * prod i, (xi i) ^ (alpha i) def ConstantCoefficientEllipticOperator {d : N} (m : N) (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) : Prop := exists data : EllipticOperatorData m L, forall alpha in data.terms, forall x y, data.coefficient alpha x = data.coefficient alpha y def ShiftedEllipticEquation {d : N} (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) (lam : R) (u f : (Fin d -> R) -> R) : Prop := forall x, L u x - lam * u x = f x theorem krylov_3_7_2_constant_coefficient_holder_solvability {d m k : N} {delta lam : R} {L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R} (hm : 0 < m) (hdelta : 0 < delta /\ delta < 1) (hlam : lam != 0) (hL : ConstantCoefficientEllipticOperator m L) : forall f, HolderOn (k + delta) univ f -> exists! u, HolderOn (k + m + delta) univ u /\ ShiftedEllipticEquation L lam u f := by sorry </pre>
Theorem 4.2.1. Let the assumptions of Theorem 4.1.2 be satisfied, and let $k \geq 0$ be an integer, $K_1 \geq 1$ be a constant. Assume that for any α we have $\alpha _\alpha \leq K_1$. Then for any λ, the inclusions	PDE	<pre> noncomputable def directionalDerivativeList {d : N} : List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R [], u => u i :: indices, u => fun x => </pre>

Natural language statement

Domain

GroundTruth

u in $C^{m+\delta}(R^d)$ and $L_\lambda u$ in $C^{k+\delta}(R^d)$ imply that u in $C^{k+m+\delta}(R^d)$. Moreover, if we take $\lambda \geq \lambda_0$ from Theorem 4.1.2 and take a real λ so that $|\lambda| \geq \lambda_0$, then there exists a constant $N > 0$ depending only on k, m, δ, K, d , such that for any u in $C^{k+m+\delta}(R^d)$ we have $|u|_{k+m+\delta} + |\lambda|^{k+m+\delta} |u|_0 \leq N(|L_\lambda u|_{k+\delta} + |\lambda|^{k+\delta} |u|_0)$.

```

fderiv R (directionalDerivativeList indices u) x (Pi.single i 1)

noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> R)) :
List (Fin d) :=
  Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i

noncomputable def multiDerivative {d : N} (alpha : (Fin d -> R))
(u : (Fin d -> R) -> R) : (Fin d -> R) -> R :=
  directionalDerivativeList (multiIndexDirections alpha) u

noncomputable def holderGauge {d : N} (k : N) (delta : R)
(Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : R>=0inf :=
  ( alpha : {alpha : (Fin d -> R) // sum i, alpha i <= k},
    x : Omega, ENNReal.ofReal [multiDerivative alpha u x])
  alpha : {alpha : (Fin d -> R) // sum i, alpha i = k},
  x : Omega, y : Omega, ( _ : (x : (Fin d -> R)) != y),
  ENNReal.ofReal
  (|multiDerivative alpha u x - multiDerivative alpha u y| /
    ||(x : (Fin d -> R)) - y|| ^ delta)

def HolderOn {d : N} (r : R) (Omega : Set (Fin d -> R))
(u : (Fin d -> R) -> R) : Prop :=
  exists k : N, exists delta : R, r = k + delta /\ 0 <= delta /\ delta < 1
  /\
    ContDiffOn R k u Omega /\ holderGauge k delta Omega u < inf

structure EllipticOperatorData {d : N} (m : N)
  (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) where
  terms : Finset ((Fin d -> R))
  order_le : forall alpha in terms, sum i, alpha i <= m
  coefficient : (Fin d -> R) -> (Fin d -> R) -> R
  formula : forall u x, L u x = sum alpha in terms, coefficient alpha x *
    multiDerivative alpha u x
  ellipticityConstant : R
  ellipticityConstant_pos : 0 < ellipticityConstant
  principalSymbol : forall (x xi : (Fin d -> R)),
    ellipticityConstant * ||xi|| ^ m <=
      |sum alpha in terms with sum i, alpha i = m,
        coefficient alpha x * prod i, (xi i) ^ (alpha i)|

def VariableCoefficientEllipticOperator {d : N} (m : N)
(L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) : Prop :=
  Nonempty (EllipticOperatorData m L)

def OperatorCoefficientsHolder {d : N} (m : N) (r : R)
(L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) : Prop :=
  exists data : EllipticOperatorData m L,
    forall alpha in data.terms, HolderOn r univ (data.coefficient alpha)

def ShiftedEllipticEquation {d : N}
(L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) (lam : R)
(u f : (Fin d -> R) -> R) : Prop :=
  forall x, L u x - lam * u x = f x

theorem krylov_4.2.1_better_regular_data_better_regular_solution
{d m k : N} {delta lam lam0 : R} {L : ((Fin d -> R) -> R) -> (Fin d ->
R) -> R}
{u f : (Fin d -> R) -> R} (hk : 0 < k) (hdelta : 0 < delta /\ delta < 1)
(hL : VariableCoefficientEllipticOperator m L)
(hcoeff : OperatorCoefficientsHolder m (k + delta) L)
(hu : HolderOn (m + delta) univ u) (hLu : ShiftedEllipticEquation L lam
u f)
(hf : HolderOn (k + delta) univ f) :
  HolderOn (k + m + delta) univ u /\
    (lam0 <= |lam| -> exists C : R>=0inf, C < inf /\
      holderGauge (k + m) delta univ u <=
        C * holderGauge k delta univ (fun x => L u x - lam * u x)) := by
  sorry

```

Theorem 4.5.1. Let $L = L(x) = \sum_{|\alpha| \leq m} a^\alpha(x) D^\alpha$ be a uniformly elliptic operator, and $k \geq 0$

PDE

```

noncomputable def directionalDerivativeList {d : N} :
List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R

```

Natural language statement

Domain

GroundTruth

be an integer. Assume that a^α in $C^{k+\delta}(R^d)$ for any α . Define the constant $\lambda_{\text{bda}0}$ depending only on the ellipticity constant κ and m, δ, d and maximum of $|a^\alpha|_{\delta}$ as in Theorem 4.1.2, and take any real λ_{bda} such that $|\lambda_{\text{bda}}| \geq \lambda_{\text{bda}0}$. Then for any f in $C^{k+\delta}(R^d)$ there exists a unique solution u in $C^{k+m+\delta}(R^d)$ of the equation $L_{\lambda_{\text{bda}}} u(x) = f(x)$, x in R^d .

```

| [], u => u
| i :: indices, u => fun x =>
  fderiv R (directionalDerivativeList indices u) x (Pi.single i 1)

noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> R)) :
List (Fin d) :=
  Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i

noncomputable def multiDerivative {d : N} (alpha : (Fin d -> R))
(u : (Fin d -> R) -> R) : (Fin d -> R) -> R :=
  directionalDerivativeList (multiIndexDirections alpha) u

noncomputable def holderGauge {d : N} (k : N) (delta : R)
(Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : R>=0inf :=
  ( alpha : {alpha : (Fin d -> R) // sum i, alpha i <= k},
    x : Omega, ENNReal.ofReal |multiDerivative alpha u x|)
  alpha : {alpha : (Fin d -> R) // sum i, alpha i = k},
  x : Omega, y : Omega, ( _ : (x : (Fin d -> R)) != y),
  ENNReal.ofReal
  (|multiDerivative alpha u x - multiDerivative alpha u y| /
    ||(x : (Fin d -> R)) - y|| ^ delta)

def HolderOn {d : N} (r : R) (Omega : Set (Fin d -> R))
(u : (Fin d -> R) -> R) : Prop :=
  exists k : N, exists delta : R, r = k + delta /\ 0 <= delta /\ delta < 1
  /\
    ContDiffOn R k u Omega /\ holderGauge k delta Omega u < inf

structure EllipticOperatorData {d : N} (m : N)
(L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) where
terms : Finset ((Fin d -> R))
order_le : forall alpha in terms, sum i, alpha i <= m
coefficient : (Fin d -> R) -> (Fin d -> R) -> R
formula : forall u x, L u x = sum alpha in terms, coefficient alpha x *
  multiDerivative alpha u x
ellipticityConstant : R
ellipticityConstant_pos : 0 < ellipticityConstant
principalSymbol : forall (x xi : (Fin d -> R)),
  ellipticityConstant * ||xi|| ^ m <=
    |sum alpha in terms with sum i, alpha i = m,
      coefficient alpha x * prod i, (xi i) ^ (alpha i)|

def VariableCoefficientEllipticOperator {d : N} (m : N)
(L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) : Prop :=
  Nonempty (EllipticOperatorData m L)

def OperatorCoefficientsHolder {d : N} (m : N) (r : R)
(L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) : Prop :=
  exists data : EllipticOperatorData m L,
    forall alpha in data.terms, HolderOn r univ (data.coefficient alpha)

def ShiftedEllipticEquation {d : N}
(L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) (lam : R)
(u f : (Fin d -> R) -> R) : Prop :=
  forall x, L u x - lam * u x = f x

theorem krylov_4_5_1_variable_coefficient_global_solvability
{d m k : N} {delta lam lam0 : R} {L : ((Fin d -> R) -> R) -> (Fin d ->
R) -> R}
(hdelta : 0 < delta /\ delta < 1) (hlarge : lam0 <= |lam|)
(hL : VariableCoefficientEllipticOperator m L)
(hcoeff : OperatorCoefficientsHolder m (k + delta) L) :
  forall f, HolderOn (k + delta) univ f ->
    exists! u, HolderOn (k + m + delta) univ u /\ ShiftedEllipticEquation
      L lam u f := by
  sorry

```

Theorem 6.5.3. For any f in $C^{k+\delta}(\Omega)$ and g in $C^{k+2+\delta}(\Omega)$ there exists a unique function u in $C^{k+2+\delta}(\Omega)$ satisfying the equation $Lu = f$ in Ω and equal to g on $\partial\Omega$.

PDE

```

noncomputable def directionalDerivativeList {d : N} :
  List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R
| [], u => u
| i :: indices, u => fun x =>

```

Natural language statement

Domain

GroundTruth

		<pre> fderiv R (directionalDerivativeList indices u) x (Pi.single i 1) noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> N)) : List (Fin d) := Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i noncomputable def multiDerivative {d : N} (alpha : (Fin d -> N)) (u : (Fin d -> R) -> R) : (Fin d -> R) -> R := directionalDerivativeList (multiIndexDirections alpha) u noncomputable def holderGauge {d : N} (k : N) (delta : R) (Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : R>=0inf := (alpha : {alpha : (Fin d -> N) // sum i, alpha i <= k}, x : Omega, ENNReal.ofReal multiDerivative alpha u x) alpha : {alpha : (Fin d -> N) // sum i, alpha i = k}, x : Omega, y : Omega, (_ : (x : (Fin d -> R)) != y), ENNReal.ofReal (multiDerivative alpha u x - multiDerivative alpha u y / (x : (Fin d -> R)) - y ^ delta) def HolderOn {d : N} (r : R) (Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : Prop := exists k : N, exists delta : R, r = k + delta /\ 0 <= delta /\ delta < 1 /\ ContDiffOn R k u Omega /\ holderGauge k delta Omega u < inf def SmoothBoundedDomain {d : N} (Omega : Set (Fin d -> R)) : Prop := RegularBoundedDomain Omega /\ exists rho : (Fin d -> R) -> R, ContDiff R top rho /\ Omega = {x rho x < 0} /\ forall x in frontier Omega, fderiv R rho x != 0 structure EllipticOperatorData {d : N} (m : N) (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) where terms : Finset ((Fin d -> N)) order_le : forall alpha in terms, sum i, alpha i <= m coefficient : (Fin d -> N) -> (Fin d -> R) -> R formula : forall u x, L u x = sum alpha in terms, coefficient alpha x * multiDerivative alpha u x ellipticityConstant : R ellipticityConstant_pos : 0 < ellipticityConstant principalSymbol : forall (x xi : (Fin d -> R)), ellipticityConstant * xi ^ m <= sum alpha in terms with sum i, alpha i = m, coefficient alpha x * prod i, (xi i) ^ (alpha i) def VariableCoefficientEllipticOperator {d : N} (m : N) (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) : Prop := Nonempty (EllipticOperatorData m L) def EllipticDirichletSolution {d : N} (Omega : Set (Fin d -> R)) (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) (f g u : (Fin d -> R) -> R) : Prop := (forall x in Omega, L u x = f x) /\ forall x in frontier Omega, u x = g x theorem krylov_6_5_3_smooth_domain_dirichlet_solvability {d k : N} {delta : R} {Omega : Set (Fin d -> R)} {L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R} (hdelta : 0 < delta /\ delta < 1) (hOmega : SmoothBoundedDomain Omega) (hL : VariableCoefficientEllipticOperator 2 L) (hcoeff : OperatorCoefficientsHolder 2 (k + delta) L) : forall f g, HolderOn (k + delta) Omega f -> HolderOn (k + 2 + delta) (closure Omega) g -> exists u, HolderOn (k + 2 + delta) Omega u /\ EllipticDirichletSolution Omega L f g u /\ forall v, HolderOn (k + 2 + delta) Omega v -> EllipticDirichletSolution Omega L f g v -> Set.EqOn v u (closure Omega) := by sorry </pre>
Theorem 7.1.2. Let Ω be a domain in $\mathbb{R}^d$ and u in $C^{m+\delta}(\Omega)$. Assume that L_λ is	PDE	<pre> noncomputable def directionalDerivativeList {d : N} : List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R </pre>

Natural language statement

Domain

GroundTruth

$C^{k+\delta}(\Omega)$ for some λ . Then u in $C^{k+m+\delta}(\Omega)$.

```

| [], u => u
| i :: indices, u => fun x =>
  fderiv R (directionalDerivativeList indices u) x (Pi.single i 1)

noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> R)) :
List (Fin d) :=
  Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i

noncomputable def multiDerivative {d : N} (alpha : (Fin d -> R))
  (u : (Fin d -> R) -> R) : (Fin d -> R) -> R :=
  directionalDerivativeList (multiIndexDirections alpha) u

noncomputable def holderGauge {d : N} (k : N) (delta : R)
  (Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : R>=0inf :=
  ( alpha : {alpha : (Fin d -> R) // sum i, alpha i <= k},
    x : Omega, ENNReal.ofReal |multiDerivative alpha u x|)
  alpha : {alpha : (Fin d -> R) // sum i, alpha i = k},
  x : Omega, y : Omega, ( _ : (x : (Fin d -> R)) != y),
  ENNReal.ofReal
  (|multiDerivative alpha u x - multiDerivative alpha u y| /
    ||(x : (Fin d -> R)) - y|| ^ delta)

def HolderOn {d : N} (r : R) (Omega : Set (Fin d -> R))
  (u : (Fin d -> R) -> R) : Prop :=
  exists k : N, exists delta : R, r = k + delta /\ 0 <= delta /\ delta < 1
  /\
    ContDiffOn R k u Omega /\ holderGauge k delta Omega u < inf

structure EllipticOperatorData {d : N} (m : N)
  (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) where
  terms : Finset ((Fin d -> R))
  order_le : forall alpha in terms, sum i, alpha i <= m
  coefficient : (Fin d -> R) -> (Fin d -> R) -> R
  formula : forall u x, L u x = sum alpha in terms, coefficient alpha x *
    multiDerivative alpha u x
  ellipticityConstant : R
  ellipticityConstant_pos : 0 < ellipticityConstant
  principalSymbol : forall (x xi : (Fin d -> R)),
    ellipticityConstant * ||xi|| ^ m <=
    |sum alpha in terms with sum i, alpha i = m,
      coefficient alpha x * prod i, (xi i) ^ (alpha i)|

def ShiftedEllipticEquation {d : N}
  (L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R) (lam : R)
  (u f : (Fin d -> R) -> R) : Prop :=
  forall x, L u x - lam * u x = f x

theorem krylov_7_1_2_interior_holder_regularization
  {d m k : N} {delta lam : R} {Omega : Set (Fin d -> R)}
  {L : ((Fin d -> R) -> R) -> (Fin d -> R) -> R} {u f : (Fin d -> R) -> R}
  (hdelta : 0 < delta /\ delta < 1) (hOmega : IsOpen Omega)
  (hL : VariableCoefficientEllipticOperator m L)
  (hcoeff : OperatorCoefficientsHolder m (k + delta) L)
  (hu : HolderOn (m + delta) Omega u)
  (hLu : ShiftedEllipticEquation L lam u f) (hf : HolderOn (k + delta)
    Omega f) :
  HolderOn (k + m + delta) Omega u := by
  sorry

```

Theorem 8.7.3. For any f in $C^{\delta/2, \delta/2}(R^{d+1})$ there exists a unique function u in $C^{1+\delta/2, 2+\delta/2}(R^{d+1})$ satisfying the equation $\Delta u - u_t - u = f$ in R^{d+1} .

PDE

```

noncomputable def directionalDerivativeList {d : N} :
  List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R
| [], u => u
| i :: indices, u => fun x =>
  fderiv R (directionalDerivativeList indices u) x (Pi.single i 1)

noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> R)) :
List (Fin d) :=
  Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i

noncomputable def multiDerivative {d : N} (alpha : (Fin d -> R))
  (u : (Fin d -> R) -> R) : (Fin d -> R) -> R :=

```

Natural language statement

Domain

GroundTruth

		<pre> directionalDerivativeList (multiIndexDirections alpha) u noncomputable def laplacian {d : N} (u : (Fin d -> R) -> R) (x : (Fin d -> R)) : R := sum i, directionalDerivativeList [i, i] u x def ParabolicHolderOn {d : N} (r : R) (Q : Set (R x (Fin d -> R))) (u : (R x (Fin d -> R)) -> R) : Prop := (forall t, HolderOn r {x (t, x) in Q} fun x => u (t, x)) /\ exists C : R, 0 <= C /\ forall x s t, (s, x) in Q -> (t, x) in Q -> u (t, x) - u (s, x) <= C * t - s ^ (r / 2) def ShiftedHeatEquation {d : N} (u f : (R x (Fin d -> R)) -> R) : Prop := forall t x, laplacian (fun y => u (t, y)) x - deriv (fun s => u (s, x)) t - u (t, x) = f (t, x) theorem krylov_8_7_3_shifted_heat_holder_solvability {d : N} {delta : R} (hdelta : 0 < delta /\ delta < 1) : forall f : (R x (Fin d -> R)) -> R, ParabolicHolderOn delta univ f -> exists! u : (R x (Fin d -> R)) -> R, ParabolicHolderOn (2 + delta) univ u /\ ShiftedHeatEquation u f := by sorry </pre>
Theorem 10.3.3. For any f in $C^{\{\delta/2, \delta\}}(Q)$ and g in $C^{\{1+\delta/2, 2+\delta\}}(Q)$ there exists a unique function u in $C^{\{1+\delta/2, 2+\delta\}}(Q)$ satisfying the equation $Lu - u_t = f$ in Q and equal g on $d'Q$.	PDE	<pre> noncomputable def directionalDerivativeList {d : N} : List (Fin d) -> ((Fin d -> R) -> R) -> (Fin d -> R) -> R [], u => u i :: indices, u => fun x => fderiv R (directionalDerivativeList indices u) x (Pi.single i 1) noncomputable def multiIndexDirections {d : N} (alpha : (Fin d -> N)) : List (Fin d) := Finset.univ.toList.flatMap fun i => List.replicate (alpha i) i noncomputable def multiDerivative {d : N} (alpha : (Fin d -> N)) (u : (Fin d -> R) -> R) : (Fin d -> R) -> R := directionalDerivativeList (multiIndexDirections alpha) u noncomputable def holderGauge {d : N} (k : N) (delta : R) (Omega : Set (Fin d -> R)) (u : (Fin d -> R) -> R) : R>=0inf := (alpha : {alpha : (Fin d -> N) // sum i, alpha i <= k}, x : Omega, ENNReal.ofReal multiDerivative alpha u x) alpha : {alpha : (Fin d -> N) // sum i, alpha i = k}, x : Omega, y : Omega, (_ : (x : (Fin d -> R)) != y), ENNReal.ofReal (multiDerivative alpha u x - multiDerivative alpha u y / (x : (Fin d -> R)) - y ^ delta) def ParabolicHolderOn {d : N} (r : R) (Q : Set (R x (Fin d -> R))) (u : (R x (Fin d -> R)) -> R) : Prop := (forall t, HolderOn r {x (t, x) in Q} fun x => u (t, x)) /\ exists C : R, 0 <= C /\ forall x s t, (s, x) in Q -> (t, x) in Q -> u (t, x) - u (s, x) <= C * t - s ^ (r / 2) def ParabolicOperator {d : N} (L : ((R x (Fin d -> R)) -> R) -> (R x (Fin d -> R)) -> R) : Prop := exists (a : (R x (Fin d -> R)) -> Matrix (Fin d) (Fin d) R) (b : (R x (Fin d -> R)) -> (Fin d -> R)) (c : (R x (Fin d -> R)) -> R) (kappa : R), 0 < kappa /\ (forall (p : (R x (Fin d -> R))) (xi : (Fin d -> R)), kappa * xi ^ 2 <= sum i, sum j, a p i j * xi i * xi j) /\ forall u p, L u p = sum i, sum j, a p i j * directionalDerivativeList [i, j] (fun x => u (p.1, x)) p.2 + sum i, b p i * fderiv R (fun x => u (p.1, x)) p.2 (Pi.single i 1) + c p * u p def parabolicBoundary {d : N} (Q : Set (R x (Fin d -> R))) : Set (R x (Fin d -> R)) := </pre>

Natural language statement

Domain

GroundTruth

		<pre>{p in frontier Q forall eps : R, 0 < eps -> exists q in Q, q.1 <= p.1 /\ dist q p < eps} def ParabolicDirichletSolution {d : N} (Q : Set (R x (Fin d -> R))) (L : ((R x (Fin d -> R)) -> R) -> (R x (Fin d -> R)) -> R) (f g u : (R x (Fin d -> R)) -> R) : Prop := (forall p in Q, L u p - deriv (fun t => u (t, p.2)) p.1 = f p) /\ forall p in parabolicBoundary Q, u p = g p theorem krylov_10_3_3_parabolic_dirichlet_domain_solvability {d : N} {delta : R} {Q : Set (R x (Fin d -> R))} {L : ((R x (Fin d -> R)) -> R) -> (R x (Fin d -> R)) -> R} (hdelta : 0 < delta /\ delta < 1) (hL : ParabolicOperator L) : forall f g, ParabolicHolderOn delta Q f -> ParabolicHolderOn (2 + delta) Q g -> exists u, ParabolicHolderOn (2 + delta) Q u /\ ParabolicDirichletSolution Q L f g u /\ forall v, ParabolicHolderOn (2 + delta) Q v -> ParabolicDirichletSolution Q L f g v -> Set.EqOn v u (Q union parabolicBoundary Q) := by sorry</pre>
--	--	---