

Theorem 1.1: formal statement and verified foundations

Falconer distance-gauge formalization project
Yongxi Lin — October 8, 2026

Status. This is an incomplete formalization notebook. The main theorem is an explicit proof obligation; it has not passed Comparator and is not registered on Palomar. The results below record the proved foundation.

The exact target

The uploaded manuscript defines

$$h_\theta(r) = r \exp\left(-(\log(1/r))^\theta\right) \quad (0 < r < 1), \quad h_\theta(0) = 0.$$

Its Theorem 1.1 asserts that, for a compact set $E \subseteq \mathbb{R}^2$,

$$\frac{2}{3} < \theta < 1, \quad \mathcal{H}^{h_\theta}(E) > 0 \implies \mathcal{L}^1(\Delta(E)) > 0.$$

$$\Delta(E) = \{\|x - y\| : x, y \in E\}.$$

The formalization uses the Euclidean norm and Mathlib's arbitrary-gauge Hausdorff measure constructor. For radii at least one, the gauge is extended by the identity; this does not affect its small-radius Hausdorff measure. The covering characterization has been proved:

$$\mathcal{H}^{h_\theta}(E) = \sup_{\rho > 0} \inf_{\substack{E \subseteq \bigcup_{n \geq 0} U_n \\ \text{diam}(U_n) \leq \rho}} \sum_{\substack{n \geq 0 \\ U_n \neq \emptyset}} h_\theta(\text{diam}(U_n)).$$

The independent `Challenge.lean` imports only Mathlib and repeats all custom definitions. The solution does not import Challenge. The theorem statement has no auxiliary Frostman, projection, or energy hypothesis.

The geometric foundation

Proposition 1. *For $0 < r < 1$, the gauge is positive and satisfies $h_\theta(r) \leq r$. Consequently $\mathcal{H}^{h_\theta}(E) \leq \mathcal{H}^1(E)$, and countable sets have zero gauge measure. Positive gauge measure supplies two distinct points of E .*

Proof. The exponential is strictly positive. The logarithm and its real power are nonnegative, so the exponential is at most one. Mathlib's gauge comparison theorem gives the measure inequality. Singletons are null for $\mathcal{H}^1$, hence also for the gauge measure; countable subadditivity gives the countable case. A set containing at most one point is countable. $\square$

The all-pairs distance map is continuous. It therefore maps the compact product $E \times E$ to a compact, Lebesgue-measurable distance set. Two distinct points give a positive distance; that fact alone does not give positive length.

The scalar summability foundation

Proposition 2. *For every $s, c > 0$, one has $\log n \leq cn^s$ for all sufficiently large natural numbers n . For every $\theta, c > 0$,*

$$\sum_{n=0}^{\infty} \exp(-cn^{\theta}) < \infty.$$

Proof. Mathlib proves $\log x = o(x^s)$ as $x \rightarrow \infty$ for $s > 0$. Restricting its eventual norm bound to natural numbers proves the first assertion. Applying that assertion with coefficient $c/2$ gives

$$\log n \leq \frac{c}{2}n^{\theta} \implies \exp(-cn^{\theta}) \leq \exp(-2 \log n) = n^{-2}.$$

The inverse-square series converges. The omitted finite initial segment has no effect on summability, so comparison proves the claim. $\square$

Proposition 3. *The branch-counting exponent and filter-error power obey*

$$3\theta - 2 > 0 \iff \theta > \frac{2}{3}, \quad \theta > \frac{2}{3} \implies -8(2\theta - 1) < -1.$$

Consequently, in the theorem's parameter range, every positive coefficient eventually absorbs the logarithm into $n^{3\theta-2}$, and

$$\sum_{n=0}^{\infty} n^{-8(2\theta-1)} < \infty.$$

Proof. The exponent inequalities are linear arithmetic. Apply the logarithmic domination result with exponent $3\theta - 2$, and apply Mathlib's real-power series criterion to the filter-error exponent. $\square$

These results justify scalar convergence once suitable shell energies and filter errors are supplied. They do not establish the geometric estimates that produce those bounds.

The exact rounded parameters

The gain has the proved exact form $\beta_N = c_{\theta}N^{\theta-1}$, with $c_{\theta} > 0$. The ceiling definitions are retained in the block parameter κ_N and tolerance ε_N . The verified lower bounds imply

$$\varepsilon_N N \geq \frac{(c_{\theta}/1000)^2}{10^6} N^{2\theta-1}, \quad \kappa_N^3 N \geq (c_{\theta}/1000)^3 N^{3\theta-2}.$$

Hence the actual rounded filter errors $(\varepsilon_N N)^{-8}$ are summable, and $\kappa_N^3 N$ eventually absorbs every fixed logarithmic cost when $\theta > 2/3$. The full Lemma 2.2 is now proved: all four budget groups hold above one common threshold. The coarse bound $T + 1 \leq 242N$ makes its derivative constants polynomial; logarithmic domination absorbs them into $2^{\varepsilon_N N/8}$. The remaining count budgets follow from the inverse tolerance bound and the growth of $\kappa_N^3 N$. All three terms in the asserted filter bound are now summable: $(\varepsilon_N N)^{-8}$, $N2^{-\varepsilon_N N}$, and $2^{-\kappa_N N/4}$. The geometric removed-mass estimate remains to be proved.

The verified gauge Frostman construction

Proposition 4. *For $0 < \theta \leq 1$ and a compact set E of positive gauge measure, there are a probability measure μ and a constant $C > 0$ such that*

$$\mu(E) = 1, \quad \text{supp } \mu \subseteq E, \quad \mu(\overline{B}(x, r)) \leq Ch_\theta(r) \quad (0 < r < 1).$$

Proof. Positive Hausdorff gauge measure gives positive gauge content. Finite dyadic trees supply nonnegative weights with uniformly positive total mass and saturated covering bounds. Choosing atoms on the compact set and normalizing gives probability measures carried by E . These measures form a weakly compact family, so a subsequence converges to a probability measure carried by E . Portmanteau transfers open-ball bounds to the limit. Comparing an arbitrary radius with consecutive dyadic scales, and applying the proved gauge doubling estimate, yields the closed-ball bound. The probability bound handles radii bounded away from zero. $\square$

Proposition 5. *The resulting measure has no atoms and has finite critical logarithmic energy for every $\gamma \geq 1$:*

$$J_\gamma(\mu) = \iint \frac{(1 + \log^+(1/\|x - y\|))^\gamma}{\|x - y\|} d\mu(x) d\mu(y) < \infty.$$

Proof. The ball bound tends to zero with the radius, proving atomlessness. The formal kernel is infinite on the diagonal. Decompose each pointwise potential into dyadic annuli; the gauge ball bound controls its n -th annular contribution by a constant times

$$(n + 1)^\gamma e^{-cn^\theta}.$$

Logarithmic domination absorbs the polynomial into half the stretched exponential. The resulting summable majorant uniformly bounds the potential. Integrating against the finite measure proves the energy bound. $\square$

The actual scalar Gaussian kernel is also controlled:

$$K_\gamma(r) = \sum_{n \geq 0} (n + 1)^\gamma 2^n e^{-4^n r^2} \leq C_\gamma \frac{(1 + \log(1/r))^\gamma}{r} \quad (0 < r < 1, \gamma \geq 0).$$

A split at the dyadic index nearest the inverse radius bounds the finite prefix and compares the shifted tail with $K_\gamma(1)$. This kernel now also controls the actual Fourier energy described on the next page.

The verified logarithmic Fourier energy

Proposition 6. *For every planar probability measure μ and $\gamma \geq 0$, a finite positive constant depending only on γ satisfies*

$$\int_{\mathbb{R}^2} |\widehat{\mu}(\xi)|^2 \frac{\log^\gamma(e + \|\xi\|)}{\|\xi\|} d\xi \leq C_\gamma(1 + J_\gamma(\mu)).$$

Proof. Mathlib's evaluated Gaussian Fourier transform and Fubini give the precise duality between Gaussian-weighted squared Fourier modulus and the spatial Gaussian kernel. Tonelli sums the non-negative dyadic Gaussian contributions. The proved scalar kernel bound controls their sum by the critical spatial energy. A dyadic scale nearest each high frequency supplies the desired logarithmic weight. At low frequencies, the probability bound controls the Fourier modulus; the extended inverse norm has a proved finite integral on the planar unit ball. Combining the regions proves Lemma 5.2 with a finite explicit constant and the manuscript's literal Fourier phase. $\square$

Separated probabilities and the prepared geometry

Two compact pieces of positive mass are extracted from the atomless Frostman probability. Their normalized restrictions retain gauge ball bounds and all finite logarithmic energies. The positive similarity used in Proposition 5.6 places the pieces in discs of radius $1/400$ inside the unit square, with centers exactly $1/4$ apart. Every cross distance lies in $[0.24, 0.26]$. These geometric and energy preparation statements are proved; the radial Orlicz density preparation remains to be established.

The actual product-distance pushforward has mass one on the all-pairs distance set. Its absolute continuity therefore gives positive length. Preparation multiplies this length by the positive similarity scale, so positivity transfers back to the original set.

The complete summable reconstruction criterion

Lemma 4.2 is now proved for a finite positive measure α and dominated approximants $\tau_N \leq \alpha$ for $N \geq N_0$. Its exact hypotheses are

$$\sum_{N \geq N_0} (\alpha(\mathbb{R}) - \tau_N(\mathbb{R})) < \infty, \quad \sum_{N \geq N_0} \int_{2^{N-1} \leq |r| \leq 2^{N+1}} |\widehat{\tau}_N(r)|^2 dr < \infty.$$

They imply $\alpha \ll \mathcal{L}^1$. The concrete convolution errors and Fourier bands form convergent L^1 and L^2 series. Telescoping the low-pass convolutions identifies their pairing with the original measure. The final absolute-continuity argument is given on the next page.

The verified reconstruction bridges

Dyadic frequency shells overlap at most three times. The quadratic L^2 bound proves convergence of square-summable shell pieces. Actual smooth low-pass kernels preserve mass; their differences have uniformly bounded L^1 norm and the precise dyadic Fourier support. Convolving removed measures with these differences produces summable L^1 errors whenever the removed masses are summable. Low-pass convolutions converge to the original measure on Schwartz tests.

Proposition 7. *Suppose a finite positive measure α satisfies*

$$\int \varphi d\alpha = \int \varphi f dx + \int (\mathcal{F}^{-1}\varphi)g d\xi \quad \text{for all Schwartz tests } \varphi, \quad f \in L^1, \quad g \in L^2.$$

Then α is absolutely continuous with respect to Lebesgue measure.

Proof. Let $d\sigma = |f| dx$. Cauchy–Schwarz and Plancherel give

$$\left| \int \varphi d\alpha \right| \leq \int |\varphi| d\sigma + \|g\|_2 \|\varphi\|_2.$$

Use smooth compact cutoffs equal to one on a compact subset of an open set U , zero off U , and bounded between zero and one. Their L^2 norm is at most $|U|^{1/2}$. Inner regularity yields

$$\alpha(U) \leq \sigma(U) + \|g\|_2 |U|^{1/2}.$$

For a Lebesgue-null set, outer regularity provides open covers whose Lebesgue and σ masses both tend to zero. The bound forces its α mass to be zero. $\square$

The actual L^2 shell pieces and the telescoped series pairing have been constructed, completing this part of the reconstruction proof.

Remaining analytic work and verification

The proof still needs orthogonal and radial Orlicz projection estimates, quantitative regular decomposition, four uniform energy estimates, and the multiscale budget induction. Fixed-parameter estimates need proved uniform dependence before use at shrinking parameters.

The main theorem currently has one explicit proof hole. Comparator permits the three standard axioms and must reject the target’s proof hole. A successful ordinary build of this skeleton is not a successful theorem verification.

The public project includes a separate sandboxed Comparator job and a manual full Palomar preflight. Registration requires a completed proof, verification of the exact public commit, truthful metadata, and the registry review.

[Public project: CoolRmal/falconer-theta-gauge](#)