


LAB MEETING 1

Katie Levandosky

July 10, 2023

SINGLE CELL

- ▶ Rac and Rho particles inside the cell and on the cell membrane
- ▶ Bind to the membrane with rate k_{on}
- ▶ Unbind from the membrane with rate k_{off}
- ▶ Recruit particles with rate k_{fb}
- ▶ Particles can also diffuse along the membrane




SINGLE CELL

- ▶ Branched and bundled actin networks
- ▶ Rac forms branched networks with actin
- ▶ Rho forms bundled networks with actin
- ▶ Branched actin is protrusive and bundled actin is contractile



Equations for branched actin (a)
and bundled actin (b):

$$\frac{\partial a(s, t)}{\partial t} = K_a(a(1 + \alpha n_{rac}(s, t)) - a^2) - m_0 ab + D_a \Delta a$$

$$\frac{\partial b(s, t)}{\partial t} = K_b(b(1 + \alpha n_{rho}(s, t)) - b^2) - m_0 ab + D_b \Delta b$$




SINGLE CELL




TWO CELLS

- ▶ Define a contact region between the two cells
- ▶ Uncoupled: nothing different happens at the contact region
- ▶ Determine if the cells are going in the same direction; within 45 degrees = yes




RAC/RHO BINDING AND UNBINDING

Change binding rate (k_{on}) or unbinding rate (k_{off}) for rac (x) or rho (y) in the contact region.




INCREASED ANTAGONISM BETWEEN RAC AND RHO

When a rac particle binds in the overlap region in cell 1, remove 1 or more nearby ($\varepsilon = 0.1, 0.5$) rho particles in cell 1. Do the same for cell 2.




CIL AND COA

- CIL = Contact Inhibition of Locomotion; when cells are in contact, down-regulate rac and up-regulate rho so cells move away from each other
- COA = Co-attraction; when cells are nearby, up-regulate rac so cells move towards each other




RAC/RHO PROPORTIONAL TO ITSELF

At each point in the contact region of cell 1, change the rac/rho binding/unbinding rate depending on the amount of rac/rho nearby in cell 1. Do the same for cell 2.



RAC/RHO PROPORTIONAL TO OTHER CELL

At each point in the contact region of cell 1, change the rac/rho binding/unbinding rate depending on the amount of rac/rho nearby in cell 2. Do the same for cell 2.




$$\frac{da}{dt} = K_a \left((1 - n_{rac}) * a - a^2 \right) - m_0 * a * b + \text{diffusion}$$

$$\frac{db}{dt} = K_b \left((1 - n_{rho}) * b - b^2 \right) - m_0 * a * b + \text{diffusion}$$


BRANCHED/BUNDLED ACTIN FORMATION

Change the rate of branched or bundled actin formation by changing the coefficient K_a or K_b in the contact region.




BRANCHED/BUNDLED PROPORTIONAL TO OTHER CELL

At each point in the contact region of cell 1, change the branched/bundled coefficient depending on the amount of branched/bundled actin nearby in cell 2. Do the same for cell 2.



RAC/RHO BINDING AND UNBINDING



NEXT: ADDING CADHERINS

- ▶ Add cadherins to incorporate more interaction between the two cells
- ▶ Cadherins will also bind and unbind from the cell membrane like rac and rho
- ▶ They will stabilize when both cells place cadherins near each other in the contact region
- ▶ Cells will place more rac in the areas with cadherins, so implement CIL in those spots

