

---


# Mass Spectrometry based Protein Quantification

---



Marc Vaudel

*Center for Medical Genetics and Molecular Medicine,  
Haukeland University Hospital, Bergen, Norway*



*KG Jebsen Center for Diabetes Research, Department of Clinical Science,  
University of Bergen, Norway*









# Tandem Mass Spectrometry



# Tandem Mass Spectrometry




# Targeted Quantification



Heavy peptides are spiked as internal control

# Data interpretation using Skyline




# Sample Labelling and Multiplexing

## A - Metabolic labelling

Control


Patient



## B - Chemical labelling

Control


Patient




## C - Label free

Control


Patient



# Tandem Mass Spectrometry





# Spectrum Counting




## Protein R

### Peptide R1




## Protein B


### Peptide B1





### Peptide B2



### Peptide B3




### Peptide B4




## Protein G



### Peptide G1



### Peptide G2



### Peptide G3



## Protein R

$$NSAF = \frac{3}{252} = 1.19 \times 10^{-2}$$

$$emPAI = 10^{\frac{1}{20}} - 1 = 0.12$$

## Protein B

$$NSAF = \frac{7}{422} = 1.66 \times 10^{-2}$$

$$emPAI = 10^{\frac{4}{37}} - 1 = 0.28$$


## Protein G

$$NSAF = \frac{4}{123} = 3.52 \times 10^{-2}$$


$$emPAI = 10^{\frac{3}{8}} - 1 = 1.37$$



# Reporter ion based quantification



# Label-free quantification



# Label-free quantification

