

# Strong self-regulation and widespread facilitative interactions between genera of phytoplankton

Coralie Picoche & Frédéric Barraquand

March 9, 2019

## Abstract

The persistence of phytoplanktonic diversity in spite of competition for basic resources has long been a source of wonder and inspiration to ecologists. To sort out, among the many coexistence mechanisms suggested by theory and experiments, which ones actually maintain diversity in natural ecosystems, long-term field studies are paramount. Here, we analyse a large dataset of phytoplankton abundance time series, counted every two weeks over 20 years, at 10 sites along the French coastline. We estimate biotic interactions using dynamic, multispecies autoregressive models. We show that a strong self-regulation, with competition strength within a genus an order of magnitude higher than between genera, was present in all phytoplanktonic interaction networks. Furthermore, positive net effects between phytoplanktonic taxa constituted at least 40% of non-null interactions in all sites. Both strong self-regulation and widespread net facilitation should therefore be key features of coexistence mechanisms intending to explain phytoplankton diversity maintenance. (148 words)

## Introduction

How species or close genera can coexist together in spite of competition is one of the main puzzles of community ecology, especially for primary producers that seemingly share the same basic resources<sup>1</sup>. Many theoretical studies of competition models have shown that competitive exclusion is likely in those circumstances<sup>2,3</sup>, unless mechanisms involving spatial or temporal variation are at play<sup>4,5,6,7</sup>. Neutral theory, that assumes a non-equilibrium coexistence maintained by dispersal and equal competitive abilities for all species<sup>8</sup> (though there are exceptions<sup>9,10</sup>) has been proposed as a solution to explain highly diverse communities<sup>8,11</sup>.

However, the evidence gathered from terrestrial plant communities starts to suggest that, in fact, niche rather than neutral processes may be paramount to explain coexistence, with intraspecific competition dwarfing interspecific competition in most cases<sup>12,13</sup>. Whether these conclusions drawn from studies of annual plants and forest trees apply to other ecosystems and taxa is currently little known (but see Mutshinda et al.<sup>14</sup>).

Moreover, even within a single trophic level, competition may not be the rule: the meta-analysis by Adler et al.<sup>13</sup> reported a large number of facilitative interactions (30%) and several reviews<sup>15,16</sup> have highlighted that facilitation

may be much more widespread than ecologists usually tend to think. Although some theoretical studies suggest that facilitative interactions can be destabilizing (*sensu* resilience) and therefore undermine coexistence in Lotka-Volterra models<sup>17</sup>, multiple other modelling<sup>18</sup> and empirical<sup>15,19</sup> studies have suggested that facilitative interactions can to a large degree benefit coexistence, especially when multiple interaction types are considered simultaneously<sup>20,21</sup>.

Here, we study a large multi-species dataset consisting of several multivariate long-term time series of phytoplankton dynamics along the French coastline, which we then analyse using multivariate autoregressive (MAR) time series models, allowing for interactions between genera. Although many ecological studies focus on interactions between species, competition has been shown experimentally to occur between different genera of phytoplankton<sup>22,23</sup>. Because competition does - according to both theory and experiments - occur between phytoplankton genera, and the genus level is a rather fine taxonomic scale for phytoplankton interaction studies (most studies are restricted to interaction between different classes or even phyla<sup>24,25,26</sup>), we study here interactions between different genera of phytoplankton, which belong mostly to diatoms and dinoflagellates. To put our results into a more general context, we then compare our interaction strength estimates to previously published interaction networks produced under the same statistical framework, both in plankton and other empirical systems.

## Results

**MAR(1) estimates** Using MAR(1) autoregressive models, we have produced interaction matrices<sup>24,27</sup> - i.e., Jacobian community matrices on the logarithmic scale<sup>24</sup>. Best-fitting models corresponded to a phylogenetically-structured interaction scenario, where interaction only occurred between closely related genera (Supplementary Fig. 3). This led to sparse, modular matrices that have two main features. First, we observe a strong self-regulation for all sites (Fig. 1, diagonal elements of all matrices), a feature that we have previously highlighted in a more detailed analysis on one of the considered study regions<sup>28</sup>. The ratio of mean intragenus to intergenus interaction coefficients varies between 6 and 10, not counting coefficients set to 0 in the estimation process. If we include the zeroes in the interaction matrix in the computation of the intra/inter mean interaction strength, the ratio rises to 21-43. Therefore, intragenus interactions are on average much stronger than intergenus interactions, approximately 10 to 20 times stronger.

Second, although the percentage of facilitative interactions seemed to vary among sites (between 40% and 71% of interactions in the selected models), facilitation remained predominant in 9 sites out of 10 (only Lazaret, in the Mediterranean Sea, has 60% negative interactions). Our observational setup being nested, with sites within regions, we can examine whether locally positive interactions remain positive in a regional context: the percentage of consistently positive interactions at the regional level varies between 30% and 53%, higher than the percentage of similarly defined negative interactions (between 15% and 40%), except for sites in the Mediterranean Sea.

We have found that the percentage of true mutualism (+/+) is substantial: averaged over all sites, 32% of all interactions are (+/+) while only 12% of them are (-/-). The sign correspondence is not always maintained between

62 French regions: the only interaction that is non-zero in the 10 sites (CHA/SKE) is mutualistic in Men er Roue only  
 63 (Brittany) and mixed (+/-) at all other sites. Within the same region, however, interactions measured at different  
 64 sites tend to keep the same sign. In the 3 sites of Marennes-Oléron, for instance, there were 4 interactions which  
 65 remained positive on both sides (CHA/GUI, DIT/GUI, LEP/THP, SKE/THP), 3 of them being also mutualistic  
 66 in some of the Brittany sites. This contradicts previous observations that mutualistic interactions tend to be more  
 67 context-dependent than competitive interactions<sup>29</sup>.




Figure 1: **Interaction matrices estimated in 10 sites along the French coastline.** Four regions are distinguished: Brittany (a), Marennes-Oléron (b), Arcachon (d) and the Mediterranean Sea (e). Only interactions between clades (pennate and centric diatoms, dinoflagellates, other planktonic taxa) are allowed, as this is the best fitting interaction scenario (Supplementary Fig. 3). Taxon  $j$  (in columns) has an effect illustrated by the bar height on taxon  $i$ 's growth rate (in rows). We present the interaction matrix minus the identity matrix ( $\mathbf{B} - \mathbf{I}$ ) because this compares unambiguously intra- and intergenera interactions. The scale for the coefficient values is given at the bottom left of panel a). 95% significance of coefficients was determined by bootstrapping and is marked by asterisks (\*). The community composition is given in Supplementary Table 2. The fraction of positive interactions in each matrix is given by points in c) while the dashed (resp., dotted) line represents the ratio of interactions remaining positive (resp., negative) for all sites of a given region.

**Interaction network analysis** The stability (*sensu* resilience<sup>30</sup>) of all interaction matrices was not strongly affected by the percentage of positive interactions or their connectivity properties (Fig. 2). The maximum modulus of the interaction matrix eigenvalues remained between 0.65 and 0.80. There was also a slight increase in stability with weighted connectance, with a drop in eigenvalue modulus for weighted connectances between 0.09 and 0.1.




Figure 2: **Relation between stability and complexity of the interaction networks.** The maximum modulus of the interaction matrix  $\mathbf{B}$  eigenvalues indicates stability *sensu* resilience. Each point color corresponds to a given region. Metrics formula for weighted connectance and linkage density are given in Supplementary Information.

Given that a direct complexity-stability link was not obvious, we investigated whether the matrix coefficients had some particular structure that could help theoretical ecology to make better null models of joint community dynamics and interactions<sup>31</sup>. We defined two scores, vulnerability (summed effect of others on the focal species growth rate, Supplementary Eq. 5) and impact (summed effect of the focal species onto other species growth rates, Supplementary Eq. 6). Relations between inter- and intragroup interactions emerged (Fig. 3): genera that were more self-regulating also had also a higher vulnerability score and a lower impact score. Those two influences are likely to trade-off: a high degree of self-regulation somehow buffers outside influences. Taxa that were less self-regulating were also more likely to have a broad range of effects onto other taxa. As these genera tended to be more abundant (Supplementary Fig. 6), they were more likely to affect other genera. It is important to note, however, that these trends are weak and there is therefore a considerable amount of randomness dominating the interaction matrix: many scenarios of self-regulation vs limitation by others are therefore possible.




Figure 3: **Relation between vulnerability/impact and self-regulation.** Average vulnerability (effects of others onto the focal taxon growth rate, a-b) and impact (effects of the focal taxon unto other's growth rates, c-d), as well as self-regulation, are computed for untransformed (a-c) or absolute (b-d) values of the coefficients of the interaction matrix ( $\mathbf{B} - \mathbf{I}$ ) for the 10 study sites. Each color corresponds to a given region (Supplementary Fig. 1). The p-value of the Pearson correlation between vulnerability (respectively impact) and self-regulation is given in the top left of each panel and the slope of the linear regression is given by the black line across scatter plots.

83 Aside from these trade-offs, some of which promote some stability (*sensu* invariability), we found no remarkable  
 84 patterns of covariation between matrix elements (other than a mean-variance scaling of interaction coefficients,  
 85 Supplementary Fig. 5).

**Literature comparison** Finally, we sought to put these results in a broader context by compiling the intra vs inter group estimates of previous MAR(1) studies of long-term observational count data (listed in Supplementary Table 3). We found that the order of magnitude of intra/inter interaction strengths considered here is not particularly above those found for most planktonic systems to which MAR(1) models have been fitted, considering that our systems are relatively high-dimensional and that the higher the dimension, the larger the intraspecific regulation<sup>32</sup>. We included in Fig. 4 not only plankton studies but also a couple of vertebrate or insect studies on less diverse communities, where interactions are stronger. The conclusion from this comparison seems to be that any diverse system of competitors and facilitators has evolved large niche differences making intragroup competition much larger in magnitude than intergroup interactions.




Figure 4: **Ratio of intra- to inter-group interaction strength in Multivariate AutoRegressive (MAR) models in the ecological literature.** The name of each studies, corresponding to each code, is given in Supplementary Table 3. The symbol color corresponds to the sparsity of the interaction matrix (e.g., the proportion of null interactions in the matrix) and the symbol shape corresponds to the number of species taken into account. Intergroup interactions were set to 0 when they were not specified in the articles (in most cases, articles removed non-significant interactions at the 95% threshold). Last column corresponds to the present study.

## Discussion

**Strong self-regulation and facilitation** We found very large niche differences between genera, translating into much higher intragenus than intergenus effects on growth rates (i.e., strong self-regulation), together with a high

98 degree of facilitative net interactions.

99 The rather high intra/intertaxon interaction strength ratio<sup>33</sup> that we found, from 5 to 20, depending on how  
 100 one counts the interactions set to zero in the estimation process, could appear extremely high in light of previous  
 101 intra/interspecific competition strength estimates of 4 to 5 by Adler et al.<sup>13</sup>. Even though their model is a different  
 102 one, i.e., Lotka-Volterra competition, we prove in the SI that the intra/inter ratio should remain commensurate.  
 103 The difference in the intra/inter ratio that we found should therefore lie elsewhere, which requires some explanation.  
 104 First, one could argue that such high intra/inter ratio arises because we consider the genus as our baseline taxonomic  
 105 unit, rather than the species. Although it is logical that niche differentiation increases as one gets up the phylogenetic  
 106 tree, and that getting down to the species level could decrease slightly that ratio, there are two good arguments  
 107 suggesting that the niche differences found here extend to the species level. First, species belonging to the various  
 108 genera considered here are often found to compete in experiments<sup>22,34,23</sup>. There is therefore a massive difference  
 109 between niches in the lab and in the field<sup>28</sup>. Second, the only other study that managed to provide MAR(1)  
 110 estimates down to the species level for phytoplankton, that of Huber and Gaedke<sup>35</sup>, provides an intra/interspecific  
 111 strength ratio similar to ours (point 7a in Fig. 4). Strong self-regulation seems therefore a genuine feature of field  
 112 phytoplanktonic communities.

113 Another main finding of our study is the large frequency of facilitative interactions, between 40 and 70%, with  
 114 30% truly mutualistic (+/+) interactions. This again can be compared to the meta-analysis by Adler et al.<sup>13</sup> who  
 115 found facilitative interactions, but with a lower fraction ( $\approx 30\%$ ). However, Adler et al.<sup>13</sup>'s review contain many  
 116 experiments while the plant literature is replete with field examples of facilitation<sup>15,16</sup>, so that facilitation could be  
 117 much higher in the field. We note that several authors using MAR(1) models previously forbade positive interactions  
 118 within the same trophic level, so that the fraction of facilitative interactions in phytoplankton cannot be computed  
 119 from literature-derived MAR(1) estimates. At the moment, it is therefore unknown how the predominance of  
 120 facilitative interactions that we found in phytoplankton compares to facilitation in terrestrial plants.

121 The large niche differences and facilitative interactions that arise when considering a single trophic level are  
 122 an emergent property, arising from hidden effects of resources or predator partitioning/sharing<sup>3</sup>. In our previous  
 123 publication investigating in detail the Arcachon study sites<sup>28</sup>, we have argued that for phytoplankton, the strong  
 124 intragroup density-dependence could arise from effects of natural enemies<sup>36,28</sup>. Natural enemies could also very  
 125 well create apparent mutualism between prey species<sup>37,38,39</sup>. We believe this to be likely true for the present  
 126 study as well, given that the new study regions (Oléron, Brittany, Mediterranean) have similar predators to the  
 127 Arcachon site (zooplankton<sup>40,41,42</sup>) and parasites (viruses<sup>43</sup>, fungi). Though natural enemies are good candidates  
 128 to explain the observed niche differences and emerging facilitation, one must bear in mind that other known drivers  
 129 of phytoplankton dynamics such as allelopathy<sup>44</sup>, auxotrophy<sup>45</sup> or hydrodynamics<sup>46</sup> can all, in theory, help create  
 130 different niches and an emerging facilitation (see last subsection of the Discussion). Finally, resources that are  
 131 usually considered limiting for all species might in fact not always be: the changes in phytoplankton absorption  
 132 spectrum documented by Burson et al.<sup>47</sup> are examples of fine-scale resource partitioning of one resource, light, that



is usually believed to be limiting for all species and genera.

### **No complexity-stability relationship but connections between self-regulation and intergroup inter-**

**actions** There was no relation between the complexity of the communities (measured as either the weighted connectance or linkage density of the interaction matrices) and their stability, as measured by the dominant eigenvalue of the interaction matrix, which measures the return time to a point equilibrium. This result is conditional upon our model being a good approximate description of the system (i.e., no multiyear limit cycles or chaotic attractors as the mapping between eigenvalues and actual stability is distorted in that case<sup>48</sup>), but we showed on a subset of this data that a fixed point in a MAR(1) model, perturbed by seasonality and abiotic variables, is an accurate description of the system<sup>28</sup>. Therefore, we are confident that the absence of complexity-resilience found here is genuine.

This absence of direct link between complexity and stability could be an actual feature of empirical systems, as shown previously by Jacquet et al.<sup>49</sup> using a different technique, even though it does contradict previous results on random matrices, especially for competitive and/or mutualistic networks<sup>50</sup>. We also found that the percentage of mutualistic interactions, that is thought to affect the stability of the network<sup>20,17,51</sup>, does not have a major impact on the network's resilience.

In addition to weighted connectance, indices at the network-level (e.g., linkage density) and at the species or genus level (vulnerability and impact) approximate the average effects exerted and sustained by any given taxa in the different study sites. While, at the network level, network structure (either complexity measures or the percentage of mutualistic interactions) did not affect stability, a relation emerged between self-regulation, necessary for coexistence, and genus-level indices. We found that the more a genus is self-regulated, the more it tends to be vulnerable to other genera impacts and the less it impacts other genera. High self-regulation usually indicates large niche differences with the rest of the community, and it makes therefore sense that a species/genus whose needs strongly differ from the others only marginally impacts the resources of the other coexisting species. Furthermore, a low self-regulation was correlated with strong average abundance, which echoes findings by Yenni et al.<sup>52</sup> who found that rare species usually show stronger self-regulation. This correlation between rarity and self-regulation could explain the lesser impact effect of high self-regulated species/genus: a species which dominate the community composition can have a major effect on the others, especially as they usually cover more space, while it is harder for rare, localized species have large impacts. However, it was more difficult to explain the reverse link: a genus that is more self-regulated and rarer was found here to be on average more vulnerable to other genera's increases in densities. Such relation implies greater stability for the network as a whole, because the taxa that are the most vulnerable to other taxa's impacts are also those that are the most self-regulated and whose dynamics are therefore less perturbed. By which mechanisms this could happen is so far unclear and open to speculation. We caution, however, that the relationships between vulnerability, impact and self-regulation that we evidenced are all relatively weak: considerable stochasticity dominates the distribution of interaction matrix coefficients.

## **Ghosts of competition past and present**

Overall, the dominance of niche differentiation in observational plankton studies is similar to what has been recently found in plant community studies<sup>10,13</sup>, if not more frequent, or empirical food webs including horizontal diversity<sup>32</sup>. Large niche differences might be due to the ghost of the competition past, i.e., competition has occurred in the past, leading to strong selection and subsequent evolution leading to progressive niche separation. In this scenario, species have evolved niches that allow them not to compete or to interact only weakly (very strong facilitative effects might be likewise destabilizing<sup>17</sup>). The likely predator effects that we highlighted above could be comprised within such niche differentiation *sensu largo*: specialized predators can make strong conspecific density-dependence emerge<sup>53,54</sup>, while switching generalists can also promote diversity<sup>55</sup>. Both predators and resources have often symmetrical effects and can therefore contribute almost equally to such past niche differentiation<sup>3</sup>.

An intriguing new possibility, dubbed the “ghost of competition present”<sup>56</sup>, suggests by contrast that spatial distributions in relation to abiotic factors might have a large impact on the interaction strengths inferred from temporal interaction models such as ours. Recent combinations of model fitting and removal experiments have shown that the model fitting usually underestimates the effect of competitors that are uncovered by removal experiments<sup>56,57</sup>. This could occur for instance if species are spatially segregated (at a small scale) because each species only exists within a domain where it is relatively competitive, while a focal species could spread out if competitors were removed. This means that a species can be limited by competitors, but act so as to minimize competition (a little like avoidance behaviour in animals) so that competition is in effect hard to detect when all species are present. This would require some fine-scale segregation between phytoplankton species at the scale of interactions, i.e., at the microscale. At the moment, it is known that turbulence generates inhomogeneities at the microscale<sup>58,59</sup> but it is quite unclear how this affects multivariate spatial patterns of species distributions (*sensu* Bolker and Pacala<sup>60</sup> or Murrell and Law<sup>61</sup>). Also, even if turbulence generates spatial structure with segregation between species, it is not quite clear that the “ghost of competition present” mechanism could work for plankton, because turbulence rather than organism movement may somehow dictates where the phytoplankton patches can or cannot appear.

## **Methods**

### **Sampling methods**

All phytoplankton counts were collected as part of the National Phytoplankton and Phycotoxin Monitoring Network (REPHY<sup>62</sup>). Since 1987, this monitoring program has required 26 sites along the French coastline to be sampled every 2 weeks within 2 hours of high tide to determine both biotic (phytoplankton counts) and abiotic (water temperature, salinity) parameters. We focused on sites which had the longest and most recent time series. We also excluded time series which had missing data for over 6 months or an average delay between sampling dates above

20 days. This reduced the number of study sites to 10 sites nested within 4 regions (Brittany, Marennes-Oléron, Arcachon and the Mediterranean Sea; Supplementary Fig. 1).

Abiotic variables (temperature, salinity) were measured directly from the boat during the sampling process while water samples for biotic analyses were fixed with a Lugol's solution and examined later. Phytoplankton cells above 20  $\mu\text{m}$  were identified at the lowest possible taxonomic level and counted with the Utermöhl method using an optical microscope<sup>63</sup>. Throughout the years and sites, more than 600 taxa were identified at different taxonomic levels. We aggregated them at the genus (or group of genera when not possible) level based on previous work<sup>64,28</sup>, except for cryptophytes and euglenophytes in Arcachon, which could not be identified below the family level. Although the taxonomic resolution used here may seem coarse in comparison to land plants, it is in fact more refined than 86% of the MAR(1) studies of phytoplankton listed in Supplementary Table 3.

For each region, the MAR(1) analysis focused on the most abundant and most frequently observed genera to avoid most of the gaps in the time series. When gaps did not exceed a month, missing values were linearly interpolated; remaining missing values were replaced by a random number between 0 and half of the lowest observed abundance<sup>65</sup>. We tested extensively this and other methods to deal with missing data in a previous publication on a subset of this dataset<sup>28</sup>. All time series were scaled and centered before MAR analyses.

## MAR(1) model

Multivariate autoregressive (MAR) models are used to determine the interspecific interactions and abiotic variables shaping a community's dynamics<sup>24</sup>. They are based on a stochastic, discrete-time Gompertz model which relates log-abundance of  $S$  species at time  $t + 1$  to interactions with the rest of the community at time  $t$ , and effects of  $V$  abiotic variables at time  $t + 1$ , following eq. 1.

$$\mathbf{n}_{t+1} = \mathbf{B}\mathbf{n}_t + \mathbf{C}\mathbf{u}_{t+1} + \mathbf{e}_t, \mathbf{e}_t \sim \mathcal{N}_S(0, \mathbf{Q}) \quad (1)$$

where  $\mathbf{n}_t$  is the  $1 \times S$  log abundance vector of abundance of phytoplankton groups,  $\mathbf{B}$  is the  $S \times S$  community (interaction) matrix,  $\mathbf{C}$  is the  $S \times V$  environment matrix describing the effects of  $V$  variables (stacked in vector  $\mathbf{u}_{t+1}$ ) on species growth, and  $\mathbf{e}_t$  is a  $1 \times S$  noise vector which covers both process and observation error, following a multivariate normal distribution with a variance-covariance matrix  $\mathbf{Q}$ .  $\mathbf{Q}$  is diagonal and we have previously showed that this parsimonious choice did not affect qualitatively the results<sup>28</sup>.

We used the MARSS package<sup>66</sup> v3.9, in R v3.3.2<sup>67</sup>, to estimate parameters with a maximum likelihood procedure.

We have previously published a detailed analysis of one of the dataset (Arcachon) for which more covariables were available<sup>28</sup>. Covariables included nutrients and hydrodynamics variables. We found that hydrodynamics variables were more influential than nutrients; nutrient dynamics contributed little to phytoplankton dynamics on the two-weeks timescale. Because temperature and salinity sum up seasonal changes in light as well as hydrology (salinity is inversely related to freshwater inflow), these represent the two key drivers needed to account for abiotic

influences<sup>68</sup>. The analysis of real data in Barraquand et al.<sup>28</sup> was complemented by that of phytoplankton-like simulated data, which confirmed the ability of MAR(1) models to infer biotic interactions and abiotic forcing (e.g., no need for extra non-linearities to model the storage effect, which was found to be nearly non-existent, as in previous analyses of plant data for which strong-self regulation was observed<sup>12,69</sup>). Furthermore, using two abiotic variables (temperature and salinity) in this study rather than the full set of used in Barraquand et al.<sup>28</sup> led to almost identical estimates to the ones obtained previously<sup>28</sup>. We are therefore confident that the MAR(1) models presented here are robust to small changes in model specification. In general, MAR(1) models tend to be fairly robust to small deviations of the underlying (non-linear) data-generating model, provided that one asks mainly order of magnitude of coefficients values (rather than precise estimates) and sign of interaction coefficients<sup>48</sup>, which is how these are models are used here. For ease of interpretation of those coefficients, we also prove the correspondance between the magnitude of intra/inter interaction strength in a MAR(1) model and a multispecies Beverton-Holt model, i.e., a discrete-time Lotka-Volterra model, in the SI.

In this study, the number of phytoplankton groups,  $S$ , varies between regions but we keep the same 2 covariates, i.e. water temperature and salinity, that could be measured for all study sites. Therefore, the dimension of the dynamical system only depends on the (square of the) number of phytoplankton groups we study, which ranges between 7 (Mediterranean Sea) and 14 (Brittany). The smallest system still requires 70 parameters to be estimated if we consider all possible interactions between species. To reduce this dimensionality and remove unnecessary parameters, we compared different ‘interaction scenarios’ based on BIC. The null interaction scenario assumed no interaction between groups of species (diagonal interaction matrix) and was compared to four other interaction scenarios. The first interaction scenario assumed that interactions could only occur between phylogenetically close organisms, i.e., within a class (namely, diatoms, dinoflagellates, and other phytoplanktonic organisms) while the second interaction scenario further differentiated pennate and centric diatoms. The third interaction scenario considered the inverse hypothesis, that only unrelated organisms could interact (i.e., a diatom could only interact with a dinoflagellate or a cryptophyte, but not with another diatom), and the last interaction scenario did not constrain the interactions at all (full interaction matrix). The second interaction scenario, hereafter called the pennate-centric scenario, had the lowest BIC for all sites and was therefore the most accurate while still parsimonious: we chose to further investigate the corresponding networks.

## Analysis of interaction strengths

The interaction matrix obtained from MAR(1) analyses can be used to determine the stability of a discrete-time dynamical system<sup>24</sup>. We compared the maximum modulus of the eigenvalues of the pennate/centric matrices in each site, as a proxy of stability, to network metrics which could be related to complexity, such as weighted connectance and linkage density<sup>59</sup>. Weighted connectance is a measure of the proportion of realized links, taking into account the shape of the flux distribution, while link density measures the average proportion and strength of interactions for a given species. These metrics are adapted to weighted interaction matrices but cannot accomodate for both

positive and negative coefficients: we therefore chose to focus on the absolute values of these coefficients, which can be linked to their strength, irrespective of interaction sign.

In addition to these network-level metrics, we also computed the average vulnerability (average effect of other taxa on a focal taxon, Supplementary Eq. 5) and impact (average effect of a focal taxon on other taxa, Supplementary Eq. 6) on both raw and absolute values of the coefficients, and compared these to the regulation a focal species exerted on itself. Raw values indicate the average effect (i.e., is the effect mostly positive or negative?) that can be expected on a species' growth rate from other planktonic species while absolute effects characterise the strength of all types of interactions on a species (i.e., is a species strongly affected by the others?).

Finally, we compared our results on self regulation/intraspecific interactions vs interspecific interactions to other published studies based on a MAR model. A list of references is given in Supplementary Table 3. Authors usually reported only coefficients that were significant at the 95% threshold, ignoring potentially many weak effects. We therefore computed both the mean value of all coefficients outside of the matrix diagonal, replacing missing values by 0 (Fig. 4, which decreases the mean intergroup interaction strength), and the mean value of statistically significant coefficients only (Supplementary Fig. 7, which increases the mean intergroup interaction strength). We should mention two potential biases associated with this comparison: low-dimensional matrices tended to be more complete (less sparse) than high-dimensional matrices, as these small interaction matrices were used to study known interaction phenomena (observed predation between organisms, for instance). There is therefore a positive correlation between sparsity and dimensionality (Supplementary Fig. 8). The number of parameters to estimate increase as the square of the number of interacting groups, leading authors to reduce this set before the estimation process. A second caveat is that while we informed our model selection by phylogeny (see above), several authors have reduced the number of estimated parameters by an automated procedure, usually based on the comparison of 100 randomly chosen interaction matrices, optimizing BIC<sup>24</sup>. The latter choice may bias high non-zero interactions in the previous literature, which is why we decided to present in the main text interaction matrices including coefficients set to zero, which should be more robust to the model selection method.

## References

- [1] G.E. Hutchinson. The paradox of the plankton. *The American Naturalist*, 95(882):137–145, 1961.
- [2] R.A. Armstrong and R. McGehee. Competitive exclusion. *The American Naturalist*, 115(2):151–170, 1980.
- [3] P. Chesson. Updates on mechanisms of maintenance of species diversity. *Journal of ecology*, 106(5):1773–1794, 2018.
- [4] R.A. Armstrong and R. McGehee. Coexistence of species competing for shared resources. *Theoretical population biology*, 9(3):317–328, 1976.

- 295 [5] P. Chesson and N. Huntly. The roles of harsh and fluctuating conditions in the dynamics of ecological com-  
296 munities. *The American Naturalist*, 150(5):519–553, 1997.
- 297 [6] J. Huisman and F.J. Weissing. Biological conditions for oscillations and chaos generated by multispecies  
298 competition. *Ecology*, 82(10):2682–2695, 2001.
- 299 [7] L. Li and P. Chesson. The effects of dynamical rates on species coexistence in a variable environment: the  
300 paradox of the plankton revisited. *The American Naturalist*, 188(2):E46–E58, 2016.
- 301 [8] S.P. Hubbell. *The Unified Neutral Theory of Biodiversity and Biogeography (MPB-32)*. Princeton University  
302 Press, 2001. ISBN: 978-0-691-02128-7.
- 303 [9] I. Volkov, J.R. Banavar, S.P. Hubbell, and A. Maritan. Neutral theory and relative species abundance in  
304 ecology. *Nature*, 424:1035–1037, 2003.
- 305 [10] I. Volkov, J.R. Banavar, S.P. Hubbell, and A. Maritan. Patterns of relative species abundance in rainforests  
306 and coral reefs. *Nature*, 450(7166):45–49, 2007.
- 307 [11] J Rosindell, SP Hubbell, and RS Etienne. The unified neutral theory of biodiversity and biogeography at age  
308 ten. *Trends in ecology & evolution*, 26(7):340, 2011.
- 309 [12] P.B. Adler, S.P. Ellner, and J.M. Levine. Coexistence of perennial plants: an embarrassment of niches. *Ecology*  
310 *letters*, 13(8):1019–1029, 2010.
- 311 [13] P.B. Adler, D. Smull, K.H. Beard, R.T. Choi, T. Furniss, A. Kulmatiski, J.M. Meiners, A.T. Tredennick, and  
312 K.E. Veblen. Competition and coexistence in plant communities: intraspecific competition is stronger than  
313 interspecific competition. *Ecology letters*, 21(9):1319–1329, 2018.
- 314 [14] C. M. Mutshinda, R. B. O’Hara, and I. P. Woiod. What drives community dynamics? *Proceedings of the*  
315 *royal society b: biological sciences*, 276(1669):2923–2929, 2009.
- 316 [15] R.W. Brooker, F.T. Maestre, R.M. Callaway, C.L. Lortie, L.A. Cavieres, G. Kunstler, P. Liancourt, K. Tiel-  
317 bürger, J.M.J. Travis, F. Anthelme, C. Armas, L. Coll, E. Corcket, S. Delzon, E. Forey, Z. Kikvidze, J. Olofsson,  
318 F. Pugnaire, C.L. Quiroz, P. Saccone, K. Schiffers, M. Seifan, B. Touzard, and R. Michalet. Facilitation in  
319 plant communities: the past, the present, and the future. *Journal of ecology*, 96(1):18–34, 2008.
- 320 [16] E.J.B. McIntire and A. Fajardo. Facilitation as a ubiquitous driver of biodiversity. *New phytologist*, 201(2):403–  
321 416, 2014.
- 322 [17] K.Z. Coyte, J. Schluter, and K.R. Foster. The ecology of the microbiome: Networks, competition, and stability.  
323 *Science*, 350(6261):663–666, 2015.
- 324 [18] K. Gross. Positive interactions among competitors can produce species-rich communities. *Ecology letters*,  
325 11(9):929–936, 2008.
- 326 [19] L.A. Cavieres and E.I. Badano. Do facilitative interactions increase species richness at the entire community  
327 level? *Journal of ecology*, 97(6):1181–1191, 2009.

- [20] A Mougi and M Kondoh. Diversity of interaction types and ecological community stability. *Science*, 337(6092):349–351, 2012.
- [21] D. García-Callejas, R. Molowny-Horas, and M.B. Araújo. The effect of multiple biotic interaction types on species persistence. *Ecology*, 99(10):2327–2337, 2018.
- [22] D. Titman. Ecological competition between algae: experimental confirmation of resource-based competition theory. *Science*, 192(4238):463–465, 1976.
- [23] B. Descamps-Julien and A. Gonzalez. Stable coexistence in a fluctuating environment: an experimental demonstration. *Ecology*, 86(10):2815–2824, 2005.
- [24] A. R. Ives, B. Dennis, K. L. Cottingham, and S. R. Carpenter. Estimating community stability and ecological interactions from time-series data. *Ecological monographs*, 73(2):301–330, 2003.
- [25] S.E. Hampton, L.R. Izmet’Eva, M.V. Moore, S.L. Katz, B. Dennis, and E.A. Silow. Sixty years of environmental change in the world’s largest freshwater lake - Lake Baikal, Siberia. *Global change biology*, 14(8):1947–1958, 2008.
- [26] J.R. Griffiths, S. Hajdu, A.S. Downing, O. Hjerne, U. Larsson, and M. Winder. Phytoplankton community interactions and environmental sensitivity in coastal and offshore habitats. *Oikos*, 125(8):1134–1143, 2015.
- [27] S.E. Hampton, E.E. Holmes, L.P. Scheef, M.D. Scheuerell, S.L. Katz, D.E. Pendleton, and E.J. Ward. Quantifying effects of abiotic and biotic drivers on community dynamics with multivariate autoregressive (MAR) models. *Ecology*, 94(12):2663–2669, 2013.
- [28] F. Barraquand, C. Picoche, D. Maurer, L. Carassou, and I. Auby. Coastal phytoplankton community dynamics and coexistence driven by intragroup density-dependence, light and hydrodynamics. *Oikos*, 127(12):1834–1852, 2018.
- [29] S.A. Chamberlain, J.L. Bronstein, and J.A. Rudgers. How context dependent are species interactions? *Ecology letters*, 17(7):881–890, 2014. Rampal Etienne, editor.
- [30] A. R. Ives and S. R. Carpenter. Stability and diversity of ecosystems. *Science*, 317(5834):58–62, 2007.
- [31] A. James, M.J. Plank, A.G. Rossberg, J. Beecham, M. Emmerson, and J.W. Pitchford. Constructing random matrices to represent real ecosystems. *The American Naturalist*, 185(5):680–692, 2015.
- [32] G. Barabás, M.J. Michalska-Smith, and S. Allesina. Self-regulation and the stability of large ecological networks. *Nature ecology & evolution*, 1(12):1870–1875, 2017.
- [33] J.M. Levine and J. HilleRisLambers. The importance of niches for the maintenance of species diversity. *Nature*, 461(7261):254–257, 2009.
- [34] D. Tilman, S.S. Kilham, and P. Kilham. Phytoplankton community ecology: the role of limiting nutrients. *Annual review of ecology and systematics*, 13:349–372, 1982.

- [35] V. Huber and U. Gaedke. The role of predation for seasonal variability patterns among phytoplankton and ciliates. *Oikos*, 114(2):265–276, 2006.
- [36] D. Haydon. Pivotal assumptions determining the relationship between stability and complexity: an analytical synthesis of the stability-complexity debate. *The American Naturalist*, 144(1):14–29, 1994.
- [37] P.A. Abrams, R.D. Holt, and J.D. Roth. Apparent competition or apparent mutualism? Shared predation when populations cycle. *Ecology*, 79(1):201–212, 1998.
- [38] F. Barraquand, L.F. New, S. Redpath, and J. Matthiopoulos. Indirect effects of primary prey population dynamics on alternative prey. *Theoretical population biology*, 103:44–59, 2015.
- [39] Peter C. de Ruiter and Ursula Gaedke. Emergent facilitation promotes biological diversity in pelagic food webs. *Food webs*, 10:15–21, 2017.
- [40] J.-L. Jamet, G. Boge, S. Richard, C. Geneys, and D. Jamet. The zooplankton community in bays of Toulon area (northwest Mediterranean Sea, France). *Hydrobiologia*, 557:155–165, 2001.
- [41] J. Modéran, P. Bouvais, V. David, S. Le Noc, B. Simon-Bouhet, N. Niquil, P. Miramand, and D. Fichet. Zooplankton community structure in a highly turbid environment (Charente estuary, France): Spatio-temporal patterns and environmental control. *Estuarine, coastal and shelf science*, 88(2):219–232, 2010.
- [42] S. Tortajada, N. Niquil, H. Blanchet, B. Grami, H. Montanié, V. David, C. Glé, B. Saint-Béat, G.A. Johnson, E. Marquis, Y. Del Amo, S. Dubois, D. Vincent, C. Dupuy, F. Jude, H.J. Hartmann, and B. Sautour. Network analysis of the planktonic food web during the spring bloom in a semi enclosed lagoon (Arcachon, SW France). *Acta oecologica*, 40:40–50, 2012.
- [43] P. Ory, H.J. Hartmann, F. Jude, C. Dupuy, Y. Del Amo, P. Catala, F. Mornet, V. Huet, B. Jan, D. Vincent, B. Sautour, and H. Montanié. Pelagic food web patterns: do they modulate virus and nanoflagellate effects on picoplankton during the phytoplankton spring bloom?: Effects of viruses and nanoflagellates on picoplankton. *Environmental microbiology*:2755–2772, 2010.
- [44] A.B. Felpeto, S. Roy, and V.M. Vasconcelos. Allelopathy prevents competitive exclusion and promotes phytoplankton biodiversity. *Oikos*, 127(1):85–98, 2018.
- [45] Y. Z. Tang, F. Koch, and C. J. Gobler. Most harmful algal bloom species are vitamin B1 and B12 auxotrophs. *Proceedings of the national academy of sciences*, 107(48):20756–20761, 2010.
- [46] M. Lévy, P.J.S. Franks, and K.S. Smith. The role of submesoscale currents in structuring marine ecosystems. *Nature communications*, 9(1):4758, 2018.
- [47] A. Burson, M. Stomp, E. Greenwell, J. Grosse, and J. Huisman. Competition for nutrients and light: testing advances in resource competition with a natural phytoplankton community. *Ecology*, 99(5):1108–1118, 2018.
- [48] G. Certain, F. Barraquand, and A. Gårdmark. How do MAR(1) models cope with hidden nonlinearities in ecological dynamics? *Methods in ecology and evolution*, 9(9):1975–1995, 2018.



- [49] C. Jacquet, C. Moritz, L. Morissette, P. Legagneux, F. Massol, P. Archambault, and D. Gravel. No complexity-stability relationship in empirical ecosystems. *Nature communications*, 7(1), 2016.
- [50] S. Allesina and S. Tang. Stability criteria for complex ecosystems. *Nature*, 483(7388):205–208, 2012.
- [51] D. García-Callejas, R. Molowny-Horas, and M.B. Araújo. Multiple interactions networks: towards more realistic descriptions of the web of life. *Oikos*, 127(1):5–22, 2018.
- [52] G. Yenni, P.B. Adler, and S. K. M. Ernest. Do persistent rare species experience stronger negative frequency dependence than common species?: Asymmetric NFD and persistent rare species. *Global ecology and biogeography*, 26(5):513–523, 2017.
- [53] R. Bagchi, R.E. Gallery, S. Gripenberg, S.J. Gurr, L. Narayan, C.E. Addis, R.P. Freckleton, and O.T. Lewis. Pathogens and insect herbivores drive rainforest plant diversity and composition. *Nature*, 506(7486):85–88, 2014.
- [54] L.S. Comita, S.A. Queenborough, S.J. Murphy, J.L. Eck, K. Xu, M. Krishnadas, N. Beckman, and Y. Zhu. Testing predictions of the Janzen-Connell hypothesis: a meta-analysis of experimental evidence for distance- and density-dependent seed and seedling survival. *Journal of ecology*, 102(4):845–856, 2014.
- [55] Sergio M Vallina, BA Ward, S Dutkiewicz, and MJ Follows. Maximal feeding with active prey-switching: a kill-the-winner functional response and its effect on global diversity and biogeography. *Progress in oceanography*, 120:93–109, 2014.
- [56] S.L. Tuck, J. Porter, M. Rees, and L.A. Turnbull. Strong responses from weakly interacting species. *Ecology letters*, 21(12):1845–1852, 2018.
- [57] P.B. Adler, A. Kleinhesselink, G. Hooker, J.B. Taylor, B. Teller, and S.P. Ellner. Weak interspecific interactions in a sagebrush steppe? Conflicting evidence from observations and experiments. *Ecology*, 99(7):1621–1632, 2018.
- [58] A.D. Barton, B.A. Ward, R.G. Williams, and M.J. Follows. The impact of fine-scale turbulence on phytoplankton community structure: Phytoplankton and turbulence. *Limnology and oceanography: fluids and environments*, 4(1):34–49, 2014.
- [59] R.E. Breier, C.C. Lalescu, D. Waas, M. Wilczek, and M.G. Mazza. Emergence of phytoplankton patchiness at small scales in mild turbulence. *Proceedings of the national academy of sciences*, 115(48):12112–12117, 2018.
- [60] B.M. Bolker and S.W. Pacala. Spatial Moment Equations for Plant Competition: Understanding Spatial Strategies and the Advantages of Short Dispersal. *The American Naturalist*, 153(6):575–602, 1999.
- [61] D.J. Murrell and R. Law. Heteromyopia and the spatial coexistence of similar competitors. *Ecology letters*, 6(1):48–59, 2003.

- [62] REPHY - French Observation, Monitoring program for Phytoplankton, and Hydrology in coastal waters. RePHY dataset - french observation and monitoring program for phytoplankton and hydrology in coastal waters. 1987-2016 metropolitan data. 2017. DOI: 10.17882/47248.
- [63] H Utermöhl. Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. *Mitt. int. ver. theor. angew. limnol.*, 9, 1958.
- [64] T. Hernández Fariñas, C. Bacher, D. Soudant, C. Belin, and L. Barillé. Assessing phytoplankton realized niches using a French national phytoplankton monitoring network. *Estuarine, coastal and shelf science*, 159:15–27, 2015.
- [65] S.E. Hampton, M.D. Scheuerell, and D.E. Schindler. Coalescence in the lake washington story: Interaction strengths in a planktonic food web. *Limnology and oceanography*, 51(5):2042–2051, 2006.
- [66] E. E. Holmes, E. J. Ward, and M. D. Scheuerell. Analysis of multivariate time-series using the MARSS package. *User guide: <http://cran.r-project.org/web/packages/marss/vignettes/userguide.pdf>*, 2014.
- [67] W. N. Venables and D. M. Smith. The R Core Team. *An introduction to r. notes on r: a programming environment for data analysis and graphics. url: [ran.r-project.org/doc/manuals/r-release/r-intro.pdf](http://cran.r-project.org/doc/manuals/r-release/r-intro.pdf)*, 2013.
- [68] L.P. Scheef, S.E. Hampton, and L.R. Izmet’eva. Inferring plankton community structure from marine and freshwater long-term data using multivariate autoregressive models. *Limnology and oceanography: methods*, 11:475–484, 2013.
- [69] S.P. Ellner, R.E. Snyder, and P.B. Adler. How to quantify the temporal storage effect using simulations instead of math. *Ecology letters*, 19(11):1333–1342, 2016.