

# Strong self-regulation and widespread facilitative interactions between genera of phytoplankton

Coralie Picoche<sup>1</sup>, Frédéric Barraquand<sup>1,2\*</sup>

**1** University of Bordeaux, Integrative and Theoretical Ecology, LabEx COTE, Bât. B2 - Allée Geoffroy St-Hilaire, 33615 Pessac, France; coralie.picoche@u-bordeaux.fr

**2** CNRS, Institute of Mathematics of Bordeaux, 351 Cours de la Libération, 33405 Talence, France; frederic.barraquand@u-bordeaux.fr

\* corresponding author

## Abstract

1. The persistence of phytoplanktonic diversity in spite of competition for basic resources has long been a source of wonder and inspiration to ecologists. To sort out, among the many coexistence mechanisms suggested by theory and experiments, which ones actually maintain diversity in natural ecosystems, long-term field studies are paramount.
2. We analysed a large dataset of phytoplankton abundance time series using dynamic, multi-variate autoregressive models. Phytoplankton was counted and identified down to the genus level, every two weeks over twenty years, at ten sites along the French coastline. Multivariate autoregressive models allowed to estimate biotic interaction networks, while also accounting for abiotic variables that may drive part of the phytoplankton fluctuations. We then analysed the ratio of intra- to inter-taxa interactions (measuring self-regulation, itself a measure of niche differentiation), the frequency of negative vs positive interactions, and how stability metrics (both at the network and genus level) relate to the network complexity and genus self-regulation or abundance.
3. We showed that a strong self-regulation, with competition strength within a taxon (genus) an order of magnitude higher than between taxa, was present in all phytoplanktonic interaction networks. This much stronger intragenus competition suggests that niche differentiation - rather than neutrality - is commonplace in phytoplankton. Furthermore, interaction networks were dominated by positive net effects between phytoplanktonic taxa (above 50% of non-zero interactions on average). While network stability (sensu resilience) was unrelated to complexity measures, we unveiled links between self-regulation, intergenus interaction strengths and abundance. The less common taxa tend to be more strongly self-regulated and can therefore maintain in spite of competition with abundant ones.
4. *Synthesis:* We prove that strong niche differentiation, widespread facilitation between phytoplanktonic taxa and stabilizing covariances between interactions strengths should be common features of coexisting phytoplankton communities in the field. These are structural properties

that we can expect to emerge from plausible mechanistic models of phytoplankton communities. We discuss mechanisms, such as predation or restricted microscale movement, that are consistent with these findings, which paves the way for further research.

**Keywords:** phytoplankton; coexistence; facilitation; mutualism; niche theory; time series; networks

# 1 Introduction

2 How species or close genera can coexist together in spite of competition is one of the main puzzles of  
3 community ecology, especially for primary producers that seemingly share the same basic resources  
4 (Hutchinson, 1961). Many theoretical studies of competition models have shown that competitive  
5 exclusion is likely in those circumstances (~~Armstrong & McGehee, 1980; Chesson, 2018~~), unless  
6 mechanisms involving spatial or temporal variation are at play (Armstrong & McGehee, 1976, 1980;  
7 Chesson & Huntly, 1997; Huisman & Weissing, 2001; Li & Chesson, 2016; Chesson, 2018). Neutral  
8 theory, that assumes a non-equilibrium coexistence maintained by dispersal and equal competitive  
9 abilities for all species (Hubbell 2001, though there are exceptions, see Volkov *et al.* 2003, 2007) has  
10 been proposed as a solution to explain highly diverse communities (Hubbell, 2001; Rosindell *et al.*,  
11 2011).

12 However, the evidence gathered from terrestrial plant communities starts to suggest that, in fact,  
13 niche rather than neutral processes may be paramount to explain coexistence, with intraspecific  
14 competition dwarfing interspecific competition in most cases (Adler *et al.*, 2010, 2018b). Whether  
15 these conclusions drawn mostly from studies of annual plants ~~and forest trees~~ apply to other  
16 ecosystems and taxa is currently little known (but see Mutshinda *et al.* 2009).

17 Moreover, competition may not be the rule: the meta-analysis by Adler *et al.* (2018b) reported a  
18 large number of facilitative interactions (30%) and several reviews (Brooker *et al.*, 2008; McIntire  
19 & Fajardo, 2014; Kinlock, 2019) have highlighted that facilitation may be much more widespread  
20 than ecologists usually tend to think. Although some theoretical studies suggest that facilitative  
21 interactions can be destabilizing (*sensu* resilience) and therefore undermine coexistence in Lotka-  
22 Volterra models (Coyte *et al.*, 2015), multiple other modelling (Gross, 2008) and empirical (Brooker  
23 *et al.*, 2008; Cavieres & Badano, 2009) studies have suggested that facilitative interactions can  
24 to a large degree benefit coexistence, especially when multiple interaction types are considered  
25 simultaneously (Mougi & Kondoh, 2012; García-Callejas *et al.*, 2018).

26 Here, we analyse a spatially replicated, long-term community-level dataset, consisting of ten  
27 multivariate time series of phytoplankton abundance along the French coastline. The time series are

modelled using multivariate autoregressive (MAR) models, allowing for interactions between genera. Although many ecological studies focus on interactions between species, competition has been shown experimentally to occur between different genera of phytoplankton (Titman, 1976; Descamps-Julien & Gonzalez, 2005). The genus level is also a rather fine taxonomic scale for phytoplankton interaction studies, as most studies are restricted to interactions between different classes or even phyla (Ives *et al.*, 2003; Hampton *et al.*, 2008; Griffiths *et al.*, 2015). Studying interactions between different genera of phytoplankton therefore both makes empirical sense in light of competition experiments and allows to estimate better-resolved networks. We focus here on genera that belong mostly to diatoms and dinoflagellates. To put our results into a more general context, we then compare our interaction strength estimates to previously published interaction networks produced under the same statistical framework, both in plankton and other empirical systems.

## Material and methods

### Sampling methods

All phytoplankton counts were collected by Ifremer coastal laboratories as part of the National Phytoplankton and Phycotoxin Monitoring Network (REPHY, 2017). Since 1987, this monitoring program has required the sampling of 26 sites along the French coastline every 2 weeks within 2 hours of high tide to document both biotic (phytoplankton counts) and abiotic (water temperature, salinity) variables. We focused on sites which had the longest time series. We also excluded time series which had missing data for over 6 months or an average delay between sampling dates above 20 days. This reduced the number of study sites to 10 sites nested within 4 regions (Brittany, Oléron, Arcachon and the Mediterranean Sea; Fig. S1 and Table S1).

Abiotic variables (temperature, salinity) were measured directly from the boat during the sampling process while water samples for biotic analyses were fixed with a Lugol's solution and examined later. Phytoplankton cells above 20  $\mu\text{m}$  were identified at the lowest possible taxonomic level and counted with the Utermöhl method using an optical microscope (Utermöhl, 1958). Throughout the years and sites, more than 600 taxa were identified at different taxonomic levels. We aggregated them at

the genus (or group of genera when not possible) level based on previous work (Table S2; [Hernández Fariñas \*et al.\* 2015](#); [Barraquand \*et al.\* 2018](#)), except for cryptophytes and euglenophytes in Arcachon, which could not be identified below the family level. Although the taxonomic resolution used here may seem coarse in comparison to land plants, it is in fact more refined than 86% of the MAR(1) studies of phytoplankton listed in Table S3.

For each region, the MAR(1) analysis focused on the most abundant and most frequently observed genera to avoid most of the gaps in the time series. Time series are plotted in Fig. S2. When gaps did not exceed a month, missing values were linearly interpolated; remaining missing values were replaced by a random number between 0 and half of the lowest observed abundance ([Hampton \*et al.\*, 2006](#)). We tested extensively this and other methods to deal with missing data in a previous publication on a subset of this dataset ([Barraquand \*et al.\*, 2018](#)). All time series were scaled and centered before MAR analyses.

## MAR(1) model

Multivariate autoregressive (MAR) models are used to determine the interspecific interactions and abiotic effects shaping a community's dynamics ([Ives \*et al.\*, 2003](#)). MAR(1) models are based on a stochastic, discrete-time Gompertz equation which relates the log-abundance of each of the  $S$  taxa at time  $t + 1$  to log-abundances of the whole community at time  $t$ , with possible interactions between taxa, and effects of  $V$  abiotic variables at time  $t + 1$ . These assumptions are encapsulated in eq. 1:

$$\mathbf{n}_{t+1} = \mathbf{B}\mathbf{n}_t + \mathbf{C}\mathbf{u}_{t+1} + \mathbf{e}_t, \mathbf{e}_t \sim \mathcal{N}_S(0, \mathbf{Q}) \quad (1)$$

where  $\mathbf{n}_t$  is the  $1 \times S$  vector of log-abundance of phytoplankton taxa,  $\mathbf{B}$  is the  $S \times S$  community (interaction) matrix,  $\mathbf{C}$  is the  $S \times V$  environment matrix describing the effects of  $V$  variables (stacked in vector  $\mathbf{u}_{t+1}$ ) on growth rates, with  $V=2$  in our case (temperature and salinity). and The noise  $\mathbf{e}_t$  is a  $1 \times S$  noise vector which covers both process and observation error, following a multivariate normal distribution with a variance-covariance matrix  $\mathbf{Q}$ .  $\mathbf{Q}$  is diagonal and we have previously showed that this parsimonious choice did not affect qualitatively the results ([Barraquand \*et al.\*,](#)

2018). We used the MARSS package (Holmes *et al.*, 2014) v3.9, in R v3.3.2 (Venables & Smith, 2013), to estimate parameters with a maximum likelihood procedure.

Our previous analysis of the Arcachon region, for which more covariables were available (Barraquand *et al.*, 2018), revealed that hydrodynamics and hydrology had more influence on phytoplankton dynamics than nutrients on the two-week timescale. Because temperature and salinity sum up seasonal changes in light as well as hydrology (salinity is inversely related to freshwater inflow), these represent the two key drivers needed to account for abiotic influences (Scheef *et al.*, 2013). They are therefore used to summarize the abiotic environment in the remainder of the article.

The analysis of real data in Barraquand *et al.* (2018) was complemented by that of simulated data mimicking the study design, which confirmed the ability of MAR(1) models to infer biotic interactions and abiotic forcings. ~~There was no need to account for extra non-linearities to model relative nonlinearities or a storage effect, as these were found to be non-existent (through threshold autoregressive models, Barraquand *et al.*, 2018).~~ A different type of model (threshold autoregressive model) showed no sign of extra non-linearities or a storage effect on a subset of the present dataset (Barraquand *et al.*, 2018). Other aspects of the MAR(1) modelling are likewise quite robust: using two abiotic variables (temperature and salinity) in this study rather than the full set used in Barraquand *et al.* (2018) led to almost identical covariate effects and interaction estimates for the Arcachon study sites. Even if some departures from the true data-generating model may not always be detectable through MAR(1) diagnostics (e.g., residuals), simulation ~~work~~analysis has showed that MAR(1) models are in general robust to nonlinearities ~~(Certain *et al.*, 2018)~~ if the inference focuses on interaction sign and order of magnitude of model coefficients (Certain *et al.*, 2018), which is how these models are used here. For ease of interpretation of MAR(1) interaction coefficients, we also prove the correspondence between the magnitude of intra/inter interaction strength in a MAR(1) model and a multispecies Beverton-Holt model, i.e., a discrete-time Lotka-Volterra model (Cushing *et al.*, 2004), in the Supporting Information.

In this study, the number of phytoplankton taxa ( $S$ ) and the community composition vary slightly between regions but sites share on average 67% of their taxa. In order to have comparable models, we also keep the same 2 covariates, i.e., water temperature and salinity, that were measured at all study

106 sites. Therefore, the dimension of the dynamical system depends on the (square of the) number of  
 107 phytoplankton taxa we study, which ranges between 7 (Mediterranean Sea) and 14 (Brittany). The  
 108 smallest system still requires 63 parameters to be estimated (49 for the 7x7 interaction matrices and  
 109 14 for the 2x7 environment matrices) if we consider all possible interactions between taxa. To reduce  
 110 this dimensionality and remove unnecessary parameters, we ~~compared~~built different ‘interaction  
 111 scenarios’ with phylogenetic grounds (Violle *et al.*, 2011; Narwani *et al.*, 2017). The ~~based-on~~  
 112 BIC-based comparison (Fig S3), ~~which~~ proved to be satisfactory in our previous analyses of both  
 113 real data and similar simulated datasets (Barraquand *et al.*, 2018, Appendix 2). The null interaction  
 114 scenario assumed no interaction between groups of species (diagonal interaction matrix) and was  
 115 compared to four other interaction scenarios. The first interaction scenario assumed that interactions  
 116 could only occur between phylogenetically close organisms, i.e., within a class (groups were then  
 117 diatoms, dinoflagellates, and other phytoplanktonic organisms) while the second interaction scenario  
 118 further differentiated pennate and centric diatoms. The third interaction scenario considered the  
 119 reverse hypothesis, that only unrelated organisms could interact (i.e., a diatom could only interact  
 120 with a dinoflagellate or a cryptophyte, but not with another diatom), and the last interaction  
 121 scenario did not constrain the interactions at all (full interaction matrix). The second interaction  
 122 scenario, hereafter called the pennate-centric scenario, had the lowest BIC for all sites (Fig. S3).  
 123 This parsimonious scenario was therefore chosen as the basis for further investigations of network  
 124 structure.

## 125 Analysis of interaction strengths

126 The interaction matrix obtained from MAR(1) analyses can be used to determine the stability of  
 127 a discrete-time dynamical system (Ives *et al.*, 2003). We compared the maximum modulus of the  
 128 eigenvalues of the pennate/centric matrices in each site, as a measure of resilience (similar to the real  
 129 part of the leading eigenvalue for continuous time model), to network metrics which could be related  
 130 to complexity, such as weighted connectance and linkage density (Bersier *et al.*, 2002). Weighted  
 131 connectance is a measure of the proportion of realized links, taking into account the shape of the  
 132 flux distribution, while link density measures the average proportion and strength of interactions for



a given taxon. These metrics are adapted to weighted interaction matrices but cannot accommodate for both positive and negative coefficients: we therefore chose to focus ~~on the absolute values of these coefficients, which can be linked to their strength,~~ on interaction strength (absolute values of the coefficients), irrespective of interaction sign.

In addition to these network-level metrics, we also computed the average vulnerability ~~(average effect of other taxa on a focal taxon, eq. S5)~~ and impact (respectively, average effect of other taxa on a focal taxon, eq. S5 and average effect of a focal taxon on other taxa, eq. S6, similar in-strength and out-strength in Kinlock, 2019) on both raw and absolute values of the coefficients. We then compared these to the regulation a focal species exerted on itself. Raw values indicate the average effect (i.e., is the effect of others mostly positive or negative?) that can be expected on a taxon's growth rate from other planktonic taxa while absolute effects characterise the strength of all types of interactions on a taxon (i.e., is a taxon strongly affected by the others?).

Finally, we compared ~~our results on the~~ the observed ratio between mean self-regulation (intra-taxon interaction strength) and mean intertaxa interaction strength to other published studies based on a MAR(1) model. A list of references is given in Table S3. Authors usually reported only coefficients that were significant at the 95% significance threshold, thus ignoring potentially many weak effects which were set to 0 by default. ~~This implies that~~ There are therefore two ways of computing the mean intertaxa interactions, i.e., taking the mean value of all coefficients outside of the matrix diagonal, including zeroes (which decreases the estimated mean intertaxa interaction strength, Fig. 4), and the mean value of statistically significant intertaxa coefficients only (which increases the estimated mean intertaxa interaction strength, Fig. S8). We considered both.

## Results

### Interaction estimates

Using MAR(1) autoregressive models, we have produced interaction matrices (Ives *et al.*, 2003; Hampton *et al.*, 2013) – i.e., Jacobian community matrices on the logarithmic abundance scale (Ives *et al.*, 2003). Best-fitting models corresponded to a phylogenetically-structured interaction

scenario, where interactions only occurred between closely related genera (Fig S3). This led to sparse, modular matrices that have two main features. First, we observe a strong self-regulation for all sites (Fig. 1, diagonal elements of all matrices), a feature that we have previously highlighted in a more detailed analysis on one of the considered study regions (Barraquand *et al.*, 2018). The ratio of mean intragenus to intergenus interaction coefficients varies between 6 and 10, not counting coefficients set to 0 in the estimation process. If we include the zeroes in the interaction matrix in the computation of the intra/inter mean interaction strength, the ratio rises to 21-43. Therefore, intragenus interactions are on average much stronger than intergenus interactions, ~~approximately~~ **10 to 20 times stronger**.

Second, although the percentage of facilitative interactions seemed to vary among sites (between 40% and 71% of interactions in the selected models), facilitation remained predominant in 9 sites out of 10 (only Lazaret, in the Mediterranean Sea, has 60% negative interactions). Our observational setup being nested, with sites within regions, we can examine whether locally positive interactions remain positive in a regional context: the percentage of consistently positive interactions at the regional level varies between 30% and 53%, higher than the percentage of similarly defined negative interactions (between 15% and 40%), except for sites in the Mediterranean Sea.

We have found that the percentage of true mutualism (+/+) is substantial: averaged over all sites, 32% of all interactions are (+/+) while only 12% of them are (-/-), see also Fig S5. The sign correspondence is not always maintained between French regions: the only interaction that is non-zero in the 10 sites (CHA/SKE) is mutualistic in Men er Roue only (Brittany) and mixed (+/-) at all other sites. Within the same region, however, interactions measured at different sites tend to keep the same sign. In the 3 sites of Oléron, for instance, there were 4 interactions which remained positive on both sides (CHA/GUI, DIT/GUI, LEP/THP, SKE/THP), 3 of them being also mutualistic in some of the Brittany sites. This contradicts previous observations that mutualistic interactions tend to be more context-dependent than competitive interactions (Chamberlain *et al.*, 2014).

## Interaction network analysis

The stability (*sensu* resilience, Ives & Carpenter 2007) of all interaction matrices was not strongly affected by the percentage of positive interactions or their connectivity properties (Fig. 2). The maximum modulus of the interaction matrix eigenvalues remained between 0.65 and 0.80. There was also a slight increase in stability with weighted connectance, with a drop in eigenvalue modulus for weighted connectances between 0.09 and 0.1.

Given that a direct complexity-stability link was not obvious, we investigated whether the matrix coefficients had some particular structure that could help theoretical ecology to make better null models of joint community dynamics and interactions (James *et al.*, 2015). We defined two scores, vulnerability (summed effect of others on the focal taxon growth rate, eq. S5) and impact (summed effect of the focal taxon onto other taxa's growth rates, eq. S6). Relations between inter- and intra-genus interactions emerged (Fig. 3): genera that were more self-regulating also had also a higher vulnerability score and a lower impact score. Those two influences are likely to trade-off: a high degree of self-regulation somehow buffers outside influences. Taxa that were less self-regulating were also more likely to have a stronger effect onto other taxa. As these genera tended to be more abundant (Fig S7), this could be mediated by the average density of a genus. It is important to note, however, that these trends are weak and there is therefore a considerable amount of randomness dominating the interaction matrix: many scenarios of self-regulation vs limitation by others are therefore possible.

Aside from the trade-offs of Fig. 3, we found no remarkable patterns of covariation between matrix elements other than a mean-variance scaling of interaction coefficients (Fig S6).

## Literature comparison

Finally, we sought to put these results in a broader context by compiling the intra vs inter group estimates of previous MAR(1) studies of long-term observational count data (listed in Table S3). We found that the order of magnitude of intra/inter interaction strengths considered here is not particularly above those found for most planktonic systems to which MAR(1) models have been

211 fitted, considering that our systems are relatively high-dimensional and that the higher the number  
 212 of taxa, the larger the intraspecific regulation (Barabás *et al.*, 2017). We included in Fig. 4 not only  
 213 plankton studies but also a couple of vertebrate or insect studies on less diverse communities, where  
 214 interactions are stronger, in order to provide lower bounds for the intra/inter ratio. The conclusion  
 215 from this comparison seems to be that, unlike small communities that can be tight-knit, any diverse  
 216 field system of competitors and facilitators has evolved large niche differences making intragroup  
 217 competition much larger in magnitude than intergroup interactions.

## 218 Discussion

### 219 Strong self-regulation and facilitation

220 We found very large niche differences between genera, translating into much higher intragenus  
 221 than intergenus effects on growth rates (i.e., strong self-regulation), together with a high degree of  
 222 facilitative net interactions.

223 The rather high intra/intertaxon interaction strength ratio (Levine & HilleRisLambers, 2009)  
 224 that we found, from 5 to 20, depending on how one counts the interactions set to zero in the  
 225 estimation process, could appear extremely high in light of previous intra/interspecific competition  
 226 strength estimates of 4 to 5 by Adler *et al.* (2018b). Even though their model is a different one,  
 227 i.e., Lotka-Volterra competition, we prove in the Supporting Information that the intra/inter ratio  
 228 should remain commensurate in a MAR(1) model. The difference in the intra/inter ratio that we  
 229 found should therefore lie elsewhere, which requires some explanation. ~~First,~~ One could argue that  
 230 such high intra/inter ratio arises because we consider the genus as our baseline taxonomic unit,  
 231 rather than the species. Although it is logical that niche differentiation increases as one gets up  
 232 the phylogenetic tree, and that getting down to the species level could slightly decrease that ratio,  
 233 there are two arguments suggesting that the niche differences found here extend to the species level.  
 234 First, phytoplankton species belonging to different genera are often found to compete in experiments  
 235 (Titman, 1976; Tilman *et al.*, 1982; Descamps-Julien & Gonzalez, 2005). In the field-based dataset  
 236 studied here, the same genera that are considered in experiments are found not to compete (or only

237 weakly), hence there must be some niche differentiation occurring in the field but not in the lab.  
238 Second, the only other study that managed to provide MAR(1) estimates down to the species level  
239 for phytoplankton, that of [Huber & Gaedke \(2006\)](#), provides an intra/interspecific strength ratio  
240 similar to ours (point 7a in Fig. 4). Strong self-regulation seems therefore a genuine feature of field  
241 phytoplanktonic communities.

242 Another main finding of our study is the large frequency of positive interactions, with 30% truly  
243 mutualistic (+/+) interactions and between 40 and 70% facilitative effects. Although a seasonal  
244 environment can generate some positive covariation between taxa, those effects have already been  
245 filtered out by the inclusion of our 2 abiotic covariates (Fig. S4). The facilitative effects shown  
246 here are therefore residual effects, once abiotic trends are accounted for. Between 40 and 70%  
247 facilitation can be compared to the meta-analysis by [Adler \*et al.\* \(2018b\)](#) who also found facilitative  
248 interactions, but less than here ( $\approx 30\%$ ). However, [Adler \*et al.\* \(2018b\)](#)'s review contains many  
249 experiments while the plant literature is replete with field examples of facilitation ([Brooker \*et al.\*,  
250 2008](#); [McIntire & Fajardo, 2014](#)), so that plant facilitation could be higher in the field. At the  
251 moment, it is therefore unknown how the predominance of facilitative interactions that we found  
252 in phytoplankton compares to facilitation in terrestrial plants. We note that several authors using  
253 MAR(1) models previously forbade positive interactions within the same trophic level, so that the  
254 fraction of facilitative interactions in plankton cannot be computed from literature-derived MAR(1)  
255 estimates.

256 The large niche differences and facilitative interactions that arise when considering a single trophic  
257 level are an emergent property, arising from hidden effects of resource or predator partitioning/sharing  
258 ([Chesson, 2018](#)). In our previous publication investigating in detail the Arcachon study sites  
259 ([Barraquand \*et al.\*, 2018](#)), we have argued that for phytoplankton, the strong intragroup density-  
260 dependence could arise from effects of natural enemies ([Haydon, 1994](#)). Natural enemies could also  
261 very well create apparent mutualism between prey species ([Abrams \*et al.\*, 1998](#); [Barraquand \*et al.\*,  
262 2015](#); [de Ruiter & Gaedke, 2017](#)). We believe this to be likely true for the present study as well,  
263 given that the new study regions (Oléron, Brittany, Mediterranean) have similar predators to the  
264 Arcachon site (zooplankton, e.g., [Jamet \*et al.\*, 2001](#); [Modéran \*et al.\*, 2010](#); [Tortajada \*et al.\*, 2012](#))

and parasites (viruses, e.g., [Ory \*et al.\*, 2010](#); fungi). Though natural enemies are good candidates to explain the observed niche differences and emerging facilitation, one must bear in mind that other known drivers of phytoplankton dynamics such as allelopathy ([Felpeto \*et al.\*, 2018](#)), auxotrophy ([Tang \*et al.\*, 2010](#)) or hydrodynamics ([Lévy \*et al.\*, 2018](#)) can all, in theory, help create different niches and an emerging facilitation (see last subsection of the Discussion). Finally, resources that are usually considered limiting for all species might in fact not always be: [Burson \*et al.\* \(2018\)](#) show that phytoplanktonic taxa specialize on different components of the light spectrum. This constitutes an example of fine-scale resource partitioning of one resource, light, that all species and genera are usually thought to compete for.

## **No complexity-stability relationship but connections between self-regulation and intergroup interactions**

There was no relation between the complexity of the communities (measured as either the weighted connectance or linkage density of the interaction matrices) and their stability, as measured by the dominant eigenvalue of the interaction matrix, which quantifies the return time to a point equilibrium (i.e., resilience). This result is conditional upon our model being a good approximate description of the system (i.e., no multiyear limit cycles or chaotic attractors as the mapping between eigenvalues and actual stability is distorted in that case, [Certain \*et al.\*, 2018](#)). However, we already showed on a subset of this data that a fixed point in a MAR(1) model, perturbed by seasonality and abiotic variables, is an accurate description of the system ([Barraquand \*et al.\*, 2018](#)). Therefore, we are confident that the absence of complexity-resilience relationship found here is genuine. This absence of direct link between complexity and stability could be an actual feature of empirical systems, as shown previously by [Jacquet \*et al.\* \(2016\)](#) using a different technique, even though it does contradict theory based on random matrices, especially for competitive and/or mutualistic networks ([Allesina & Tang, 2012](#)). We also found that the percentage of mutualistic interactions, that is thought to affect the stability of a network ([Mougi & Kondoh, 2012](#); [Coyte \*et al.\*, 2015](#); [García-Callejas \*et al.\*, 2018](#)), does not have a major impact on our networks' resilience.

In addition to weighted connectance, indices at the network-level (e.g., linkage density) and

at the species or genus level (vulnerability and impact) approximate the average effects exerted  
 and sustained by any given taxa in the different study sites. While, at the network level, network  
 structure (either complexity measures or the percentage of mutualistic interactions) did not affect  
 stability, a relation emerged between self-regulation, necessary for coexistence, and genus-level  
 indices. We found that the more a genus is self-regulated, the more it tends to be vulnerable to  
 other genera impacts and the less it impacts other genera. We examined whether vulnerability  
 and impact could be affected by phylogenetic correlations; they were not as on Fig. 3 points were  
 not clustered according to genus, family or phylum. High self-regulation usually indicates large  
 niche differences with the rest of the community, and it makes therefore sense that a species/genus  
 whose needs strongly differ from the others only marginally impacts the resources of the other  
 coexisting species. This is what we expect under strong niche partitioning. Furthermore, a low  
 self-regulation was correlated with high average abundance, which echoes findings by Yenni *et al.*  
 (2017) who demonstrated that rare species usually show stronger self-regulation. This correlation  
 between relative rarity and self-regulation could also explain the lesser impact of high self-regulated  
 species/genus: a taxon which dominates the community composition can have a major effect on  
 the others, especially as they usually cover more space, while it is harder for rare, localised the least  
 common taxa to have large impacts. However, it was more difficult to explain the relationship  
 between self-regulation and vulnerability: a genus that is more self-regulated and rarer less common  
 was found here to be on average more vulnerable to other genera's increases in densities. Such  
 relation implies greater stability (*sensu* resilience, Ives *et al.* 2003, and also invariability, Arnoldi  
*et al.* 2019) for the network as a whole, because the taxa that are the most vulnerable to other  
 taxa's impacts are also those whose dynamics are intrinsically more buffered. By which mechanisms  
 this could happen is so far unclear and open to speculation. It could be just a "mass effect":  
 common taxa are in high enough numbers to deplete resources or change the environment in ways  
 that affects the rarer less common ones, but the reverse is not true. We caution, however, that the  
 relationships between vulnerability, impact and self-regulation that we evidenced are all relatively  
 weak: considerable stochasticity still dominates the distribution of interaction matrix coefficients.

## Ghosts of competition past and present

Overall, the dominance of niche differentiation in observational plankton studies – based on our analysis of the REPHY dataset and re-analysis of the MAR(1) literature – is similar to what has been recently found in plant community studies (Volkov *et al.*, 2007; Adler *et al.*, 2018b) or empirically parameterized food webs including horizontal diversity (Barabás *et al.*, 2017). Large niche differences might be due to the ghost of competition past, i.e., competition has occurred in the past, leading to strong selection and subsequent evolution, and then to progressive niche separation. In this scenario, species have evolved niches that allow them not to compete or to interact only weakly (very strong facilitative effects might be likewise destabilizing, Coyte *et al.*, 2015). The likely predator effects that we highlighted above could be comprised within such niche differentiation *sensu largo*: specialized predators can make strong conspecific density-dependence emerge (Bagchi *et al.*, 2014; Comita *et al.*, 2014), while switching generalists can also promote diversity (Vallina *et al.*, 2014). Both predators and resources have often symmetrical effects and can therefore contribute almost equally to such past niche differentiation (Chesson, 2018).

An intriguing new possibility, dubbed the “ghost of competition present” (Tuck *et al.*, 2018), suggests by contrast that spatial distributions in relation to abiotic factors might have a large impact on the interaction strengths inferred from temporal interaction models such as ours. Recent combinations of model fitting and removal experiments have shown that the model fitting usually underestimates the effect of competitors that are uncovered by removal experiments (Tuck *et al.*, 2018; Adler *et al.*, 2018a). This could occur for instance if species are spatially segregated (at a small scale) because each species only exists within a domain where it is relatively competitive (Pacala’s spatial segregation hypothesis Pacala & Levin 1997), while a focal species could spread out if competitors were removed. This means that a species can be limited by competitors, but act so as to minimize competition (a little like avoidance behaviour in animals) and maximize opportunities for positive interactions, which implies that competition is in effect hard to detect when all species are present. This would require spatial segregation between phytoplankton species at the scale of interactions, i.e., at the microscale. At the moment, it is known that turbulence generates



inhomogeneities at the microscale (Barton *et al.*, 2014; Breier *et al.*, 2018) but it is quite unclear how this affects multivariate spatial patterns of species distributions (*sensu* Bolker & Pacala 1999, or Murrell & Law 2003). Moreover, even if turbulence generates spatial structure with segregation between species, it is not quite clear that the “ghost of competition present” mechanism could work for plankton, because turbulence rather than organism movement dictates where the phytoplankton patches can or cannot appear.

## Acknowledgments

This study was only made possible by the dedication of all members of the REPHY program by Ifremer (REPHY, 2017), providing invaluable data through years of fieldwork. We are grateful to David Murrell for his careful reading and suggestions, and to Peter Adler for helpful exchanges. This study was supported by the French ANR through LabEx COTE (ANR-10-LABX-45).

**Supporting Information:** This article contains supporting information.

**Authors’ contributions:** CP and FB contributed equally to the project design. CP wrote the code for the analyses. FB and CP interpreted the results and wrote the manuscript.

**Data accessibility:** The REPHY dataset has already been published (REPHY, 2017) and all scripts for MAR and subsequent network analyses are available online in a GitHub repository (<https://github.com/CoraliePicoche/REPHY-littoral>). This repository will be made public upon acceptance and codes can be shared with referees should they wish to access them.

## References

- Abrams, P., Holt, R. & Roth, J. (1998). Apparent competition or apparent mutualism? Shared predation when populations cycle. *Ecology*, 79, 201–212. ([document](#))
- Adler, P., Ellner, S. & Levine, J. (2010). Coexistence of perennial plants: an embarrassment of niches. *Ecology Letters*, 13, 1019–1029. ([document](#))

- Adler, P., Kleinhesselink, A., Hooker, G., Taylor, J., Teller, B. & Ellner, S. (2018a). Weak interspecific interactions in a sagebrush steppe? Conflicting evidence from observations and experiments. *Ecology*, 99, 1621–1632. ([document](#))
- Adler, P., Smull, D., Beard, K., Choi, R., Furniss, T., Kulmatiski, A., Meiners, J., Tredennick, A. & Veblen, K. (2018b). Competition and coexistence in plant communities: intraspecific competition is stronger than interspecific competition. *Ecology Letters*, 21, 1319–1329. ([document](#))
- Allesina, S. & Tang, S. (2012). Stability criteria for complex ecosystems. *Nature*, 483, 205–208. ([document](#))
- Armstrong, R. & McGehee, R. (1976). Coexistence of species competing for shared resources. *Theoretical Population Biology*, 9, 317–328. ([document](#))
- Armstrong, R. & McGehee, R. (1980). Competitive exclusion. *The American Naturalist*, 115, 151–170. ([document](#))
- Arnoldi, J.F., Loreau, M. & Haegeman, B. (2019). The inherent multidimensionality of temporal variability: how common and rare species shape stability patterns. *Ecology Letters*, 22, 1557–1567. ([document](#))
- Bagchi, R., Gallery, R., Gripenberg, S., Gurr, S., Narayan, L., Addis, C., Freckleton, R. & Lewis, O. (2014). Pathogens and insect herbivores drive rainforest plant diversity and composition. *Nature*, 506, 85–88. ([document](#))
- Barabás, G., Michalska-Smith, M. & Allesina, S. (2017). Self-regulation and the stability of large ecological networks. *Nature Ecology & Evolution*, 1, 1870–1875. ([document](#))
- Barraquand, F., New, L., Redpath, S. & Matthiopoulos, J. (2015). Indirect effects of primary prey population dynamics on alternative prey. *Theoretical Population Biology*, 103, 44–59. ([document](#))
- Barraquand, F., Picoche, C., Maurer, D., Carassou, L. & Auby, I. (2018). Coastal phytoplankton community dynamics and coexistence driven by intragroup density-dependence, light and hydrodynamics. *Oikos*, 127, 1834–1852. ([document](#))

- Barton, A., Ward, B., Williams, R. & Follows, M. (2014). The impact of fine-scale turbulence on phytoplankton community structure: Phytoplankton and turbulence. *Limnology and Oceanography: Fluids and Environments*, 4, 34–49. ([document](#))
- Bersier, L.F., Banasek-Richter, C. & Cattin, M.F. (2002). Quantitative descriptors of food-web matrices. *Ecology*, 83, 2394. ([document](#))
- Bolker, B. & Pacala, S. (1999). Spatial moment equations for plant competition: understanding spatial strategies and the advantages of short dispersal. *The American Naturalist*, 153, 575–602. ([document](#))
- Breier, R., Lalescu, C., Waas, D., Wilczek, M. & Mazza, M. (2018). Emergence of phytoplankton patchiness at small scales in mild turbulence. *Proceedings of the National Academy of Sciences*, 115, 12112–12117. ([document](#))
- Brooker, R., Maestre, F., Callaway, R., Lortie, C., Cavieres, L., Kunstler, G., Liancourt, P., Tielbörger, K., Travis, J., Anthelme, F., Armas, C., Coll, L., Corcket, E., Delzon, S., Forey, E., Kikvidze, Z., Olofsson, J., Pugnaire, F., Quiroz, C., Saccone, P., Schiffers, K., Seifan, M., Touzard, B. & Michalet, R. (2008). Facilitation in plant communities: the past, the present, and the future. *Journal of Ecology*, 96, 18–34. ([document](#))
- Burson, A., Stomp, M., Greenwell, E., Grosse, J. & Huisman, J. (2018). Competition for nutrients and light: testing advances in resource competition with a natural phytoplankton community. *Ecology*, 99, 1108–1118. ([document](#))
- Cavieres, L. & Badano, E. (2009). Do facilitative interactions increase species richness at the entire community level? *Journal of Ecology*, 97, 1181–1191. ([document](#))
- Certain, G., Barraquand, F. & Gårdmark, A. (2018). How do MAR(1) models cope with hidden nonlinearities in ecological dynamics? *Methods in Ecology and Evolution*, 9, 1975–1995. ([document](#))
- Chamberlain, S., Bronstein, J. & Rudgers, J. (2014). How context dependent are species interactions? *Ecology Letters*, 17, 881–890. ([document](#))

- Chesson, P. (2018). Updates on mechanisms of maintenance of species diversity. *Journal of Ecology*, 106, 1773–1794. ([document](#))
- Chesson, P. & Huntly, N. (1997). The roles of harsh and fluctuating conditions in the dynamics of ecological communities. *The American Naturalist*, 150, 519–553. ([document](#))
- Comita, L., Queenborough, S., Murphy, S., Eck, J., Xu, K., Krishnadas, M., Beckman, N. & Zhu, Y. (2014). Testing predictions of the Janzen-Connell hypothesis: a meta-analysis of experimental evidence for distance- and density-dependent seed and seedling survival. *Journal of Ecology*, 102, 845–856. ([document](#))
- Coyte, K., Schluter, J. & Foster, K. (2015). The ecology of the microbiome: Networks, competition, and stability. *Science*, 350, 663–666. ([document](#))
- Cushing, J., Leverage, S., Chitnis, N. & Henson, S.M. (2004). Some discrete competition models and the competitive exclusion principle. *Journal of Difference Equations and Applications*, 10, 1139–1151. ([document](#))
- Descamps-Julien, B. & Gonzalez, A. (2005). Stable coexistence in a fluctuating environment: an experimental demonstration. *Ecology*, 86, 2815–2824. ([document](#))
- Felpeto, A., Roy, S. & Vasconcelos, V. (2018). Allelopathy prevents competitive exclusion and promotes phytoplankton biodiversity. *Oikos*, 127, 85–98. ([document](#))
- García-Callejas, D., Molowny-Horas, R. & Araújo, M. (2018). The effect of multiple biotic interaction types on species persistence. *Ecology*, 99, 2327–2337. ([document](#))
- García-Callejas, D., Molowny-Horas, R. & Araújo, M. (2018). Multiple interactions networks: towards more realistic descriptions of the web of life. *Oikos*, 127, 5–22. ([document](#))
- Griffiths, J., Hajdu, S., Downing, A., Hjerne, O., Larsson, U. & Winder, M. (2015). Phytoplankton community interactions and environmental sensitivity in coastal and offshore habitats. *Oikos*, 125, 1134–1143. ([document](#))

- Gross, K. (2008). Positive interactions among competitors can produce species-rich communities. *Ecology Letters*, 11, 929–936. ([document](#))
- Hampton, S., Holmes, E., Scheef, L., Scheuerell, M., Katz, S., Pendleton, D. & Ward, E. (2013). Quantifying effects of abiotic and biotic drivers on community dynamics with multivariate autoregressive (MAR) models. *Ecology*, 94, 2663–2669. ([document](#))
- Hampton, S., Izmet'Eva, L., Moore, M., Katz, S., Dennis, B. & Silow, E. (2008). Sixty years of environmental change in the world's largest freshwater lake - Lake Baikal, Siberia. *Global Change Biology*, 14, 1947–1958. ([document](#))
- Hampton, S., Scheuerell, M. & Schindler, D. (2006). Coalescence in the lake washington story: Interaction strengths in a planktonic food web. *Limnology and Oceanography*, 51, 2042–2051. ([document](#))
- Haydon, D. (1994). Pivotal assumptions determining the relationship between stability and complexity: an analytical synthesis of the stability-complexity debate. *The American Naturalist*, 144, 14–29. ([document](#))
- Hernández Fariñas, T., Bacher, C., Soudant, D., Belin, C. & Barillé, L. (2015). Assessing phytoplankton realized niches using a French national phytoplankton monitoring network. *Estuarine, Coastal and Shelf Science*, 159, 15–27. ([document](#))
- Holmes, E.E., Ward, E.J. & Scheuerell, M.D. (2014). Analysis of multivariate time-series using the MARSS package. *User guide: <http://cran.r-project.org/web/packages/MARSS/vignettes/UserGuide.pdf>*. ([document](#))
- Hubbell, S. (2001). *The unified neutral theory of biodiversity and biogeography*. Princeton University Press. ([document](#))
- Huber, V. & Gaedke, U. (2006). The role of predation for seasonal variability patterns among phytoplankton and ciliates. *Oikos*, 114, 265–276. ([document](#))


- Huisman, J. & Weissing, F. (2001). Biological conditions for oscillations and chaos generated by multispecies competition. *Ecology*, 82, 2682–2695. ([document](#))
- Hutchinson, G. (1961). The paradox of the plankton. *The American Naturalist*, 95, 137–145. ([document](#))
- Ives, A.R. & Carpenter, S.R. (2007). Stability and diversity of ecosystems. *Science*, 317, 58–62. ([document](#))
- Ives, A.R., Dennis, B., Cottingham, K.L. & Carpenter, S.R. (2003). Estimating community stability and ecological interactions from time-series data. *Ecological Monographs*, 73, 301–330. ([document](#))
- Jacquet, C., Moritz, C., Morissette, L., Legagneux, P., Massol, F., Archambault, P. & Gravel, D. (2016). No complexity-stability relationship in empirical ecosystems. *Nature Communications*, 7. ([document](#))
- James, A., Plank, M., Rossberg, A., Beecham, J., Emmerson, M. & Pitchford, J. (2015). Constructing random matrices to represent real ecosystems. *The American Naturalist*, 185, 680–692. ([document](#))
- Jamet, J.L., Boge, G., Richard, S., Geneys, C. & Jamet, D. (2001). The zooplankton community in bays of Toulon area (northwest Mediterranean Sea, France). *Hydrobiologia*, 557, 155–165. ([document](#))
- Kinlock, N.L. (2019). A Meta-analysis of Plant Interaction Networks Reveals Competitive Hierarchies as well as Facilitation and Intransitivity. *The American Naturalist*, 194, 640–653. ([document](#))
- Levine, J. & HilleRisLambers, J. (2009). The importance of niches for the maintenance of species diversity. *Nature*, 461, 254–257. ([document](#))
- Lévy, M., Franks, P. & Smith, K. (2018). The role of submesoscale currents in structuring marine ecosystems. *Nature Communications*, 9, 4758. ([document](#))
- Li, L. & Chesson, P. (2016). The effects of dynamical rates on species coexistence in a variable environment: the paradox of the plankton revisited. *The American Naturalist*, 188, E46–E58. ([document](#))

- McIntire, E. & Fajardo, A. (2014). Facilitation as a ubiquitous driver of biodiversity. *New Phytologist*, 201, 403–416. ([document](#))
- Modéran, J., Bouvais, P., David, V., Le Noc, S., Simon-Bouhet, B., Niquil, N., Miramand, P. & Fichet, D. (2010). Zooplankton community structure in a highly turbid environment (Charente estuary, France): Spatio-temporal patterns and environmental control. *Estuarine, Coastal and Shelf Science*, 88, 219–232. ([document](#))
- Mougi, A. & Kondoh, M. (2012). Diversity of interaction types and ecological community stability. *Science*, 337, 349–351. ([document](#))
- Murrell, D. & Law, R. (2003). Heteromyopia and the spatial coexistence of similar competitors. *Ecology Letters*, 6, 48–59. ([document](#))
- Mutshinda, C.M., O'Hara, R.B. & Woiod, I.P. (2009). What drives community dynamics? *Proceedings of the Royal Society B: Biological Sciences*, 276, 2923–2929. ([document](#))
- Narwani, A., Bentlage, B., Alexandrou, M., Fritschie, K., Delwiche, C., Oakley, T. & Cardinale, B. (2017). Ecological interactions and coexistence are predicted by gene expression similarity in freshwater green algae. *Journal of Ecology*, 105, 580–591. ([document](#))
- Ory, P., Hartmann, H., Jude, F., Dupuy, C., Del Amo, Y., Catala, P., Mornet, F., Huet, V., Jan, B., Vincent, D., Sautour, B. & Montanié, H. (2010). Pelagic food web patterns: do they modulate virus and nanoflagellate effects on picoplankton during the phytoplankton spring bloom? Effects of viruses and nanoflagellates on picoplankton. *Environmental Microbiology*, pp. 2755–2772. ([document](#))
- Pacala, S. & Levin, S. (1997). *Spatial ecology: the role of space in population dynamics and interspecific interactions*. Princeton University Press, Princeton, NJ. ([document](#))
- REPHY (2017). *REPHY dataset - French Observation and Monitoring program for Phytoplankton and Hydrology in coastal waters. 1987-2016 Metropolitan data*. <https://www.seanoe.org/data/00361/47248/>. ([document](#))


- Rosindell, J., Hubbell, S. & Etienne, R. (2011). The unified neutral theory of biodiversity and biogeography at age ten. *Trends in Ecology & Evolution*, 26, 340. ([document](#))
- de Ruiter, P. & Gaedke, U. (2017). Emergent facilitation promotes biological diversity in pelagic food webs. *Food Webs*, 10, 15–21. ([document](#))
- Scheef, L., Hampton, S. & Izmet'eva, L. (2013). Inferring plankton community structure from marine and freshwater long-term data using multivariate autoregressive models. *Limnology and Oceanography: Methods*, 11, 475–484. ([document](#))
- Tang, Y.Z., Koch, F. & Gobler, C.J. (2010). Most harmful algal bloom species are vitamin B1 and B12 auxotrophs. *Proceedings of the National Academy of Sciences*, 107, 20756–20761. ([document](#))
- Tilman, D., Kilham, S. & Kilham, P. (1982). Phytoplankton community ecology: the role of limiting nutrients. *Annual Review of Ecology and Systematics*, 13, 349–372. ([document](#))
- Titman, D. (1976). Ecological competition between algae: experimental confirmation of resource-based competition theory. *Science*, 192, 463–465. ([document](#))
- Tortajada, S., Niquil, N., Blanchet, H., Grami, B., Montanié, H., David, V., Glé, C., Saint-Béat, B., Johnson, G., Marquis, E., Del Amo, Y., Dubois, S., Vincent, D., Dupuy, C., Jude, F., Hartmann, H. & Sautour, B. (2012). Network analysis of the planktonic food web during the spring bloom in a semi enclosed lagoon (Arcachon, SW France). *Acta Oecologica*, 40, 40–50. ([document](#))
- Tuck, S., Porter, J., Rees, M. & Turnbull, L. (2018). Strong responses from weakly interacting species. *Ecology Letters*, 21, 1845–1852. ([document](#))
- Utermöhl, H. (1958). Zur Vervollkommnung der quantitativen Phytoplankton-Methodik. *Mitt. int. Ver. theor. angew. Limnol.*, 9. ([document](#))
- Vallina, S., Ward, B., Dutkiewicz, S. & Follows, M. (2014). Maximal feeding with active prey-switching: A kill-the-winner functional response and its effect on global diversity and biogeography. *Progress in Oceanography*, 120, 93–109. ([document](#))




- Venables, W.N. & Smith, D.M. (2013). the R Core Team. *An Introduction to R. Notes on R: A Programming Environment for Data Analysis and Graphics*. URL: [ran. r-project. org/doc/manuals/r-release/R-intro. pdf](http://ran.r-project.org/doc/manuals/r-release/R-intro.pdf). ([document](#))
- Violle, C., Nemergut, D., Pu, Z. & Jiang, L. (2011). Phylogenetic limiting similarity and competitive exclusion. *Ecology Letters*, 14, 782–787. ([document](#))
- Volkov, I., Banavar, J., Hubbell, S. & Maritan, A. (2003). Neutral theory and relative species abundance in ecology. *Nature*, 424, 1035–1037. ([document](#))
- Volkov, I., Banavar, J., Hubbell, S. & Maritan, A. (2007). Patterns of relative species abundance in rainforests and coral reefs. *Nature*, 450, 45–49. ([document](#))
- Yenni, G., Adler, P. & Ernest, S.K.M. (2017). Do persistent rare species experience stronger negative frequency dependence than common species? Asymmetric NFD and persistent rare species. *Global Ecology and Biogeography*, 26, 513–523. ([document](#))




**Fig 1. Interaction matrices estimated in 10 sites along the French coastline.** Four regions are distinguished: Brittany (a), Oléron (b), Arcachon (d) and the Mediterranean Sea (e). Only interactions within a clade (pennate and centric diatoms, dinoflagellates, other planktonic taxa) are allowed, as this is the best fitting interaction scenario (Fig S3). Taxon  $j$  (in columns) has an effect on taxon  $i$ 's growth rate (in rows) illustrated by proportional to the bar height. We present the interaction matrix minus the identity matrix ( $\mathbf{B} - \mathbf{I}$ ) because this compares unambiguously intra- and intergenera interactions. The scale for the coefficient values is given at the bottom left of panel a). 95% significance of coefficients is marked by asterisks (\*). The community composition is given in Table S2. The fraction of positive interactions in each matrix is given by points in c) while the dashed (resp., dotted) line represents the ratio of interactions remaining positive (resp., negative) for all sites of a given region.



**Fig 2. Relation between stability and complexity of the interaction networks.** The maximum modulus of the interaction matrix  $B$  eigenvalues indicates stability *sensu* resilience. Each color and shape corresponds to a given region. Metrics formula for weighted connectance and linkage density are given in the Supporting Information.



**Fig 3. Relation between vulnerability/impact and self-regulation.** Average vulnerability (effects of others on the focal taxon growth rate, a-b) and impact (effects of the focal taxon on others' growth rates, c-d), as well as self-regulation, are computed for untransformed (a-c) or absolute (b-d) values of the coefficients of the interaction matrix ( $\mathbf{B} - \mathbf{I}$ ) for the 10 study sites. Each color corresponds to a given region (Fig S1). Linear regressions are shown as black lines.



**Fig 4. Ratio of intra- to inter-group interaction strength in Multivariate AutoRegressive (MAR) models.** The ~~name of~~ reference for each study, corresponding to a code, is given in Table S3. Codes beginning with letters correspond to the present study (Ar: Arcachon; Ol: Oléron; Br: Brittany; Me: Mediterranean Sea). The symbol color and shape correspond to the sparsity of the interaction matrix (e.g., the proportion of null interactions in the matrix). Intergroup interactions were set to 0 when they were not specified in the articles (in most cases, authors removed non-significant interactions at the 95% threshold; Fig. S8 is the same figure taking into account only significant interactions)