

Figure 1: Map of the studied sites

| Code | Taxa  |
|------|---|
| AST  | Asterionella+Asterionellopsis+Asteroplanus            |
| CHA  | Chaetoceros   |
| CRY  | Cryptophytes  |
| DIT  | Ditylum   |
| EUG  | Euglenophytes   |
| GUI  | Guinardia   |
| GYM  | Gymnodinium+Gyrodinium                                |
| LEP  | Leptocylindrus  |
| NIT  | Nitzschia+Hantzschia                                  |
| PLE  | Pleurosigma+Gyrosigma                                 |
| PRO  | Prorocentrum  |
| PRP  | Protoperidinium+Archaeoperidinium+Peridinium          |
| PSE  | Pseudo-nitzschia                                      |
| RHI  | Rhizosolenia+Neocalyptrella                           |
| SCR  | Scrippsiella+Enciculifera+Pentapharsodinium+Bysmatrum |
| SKE  | Skeletonema   |
| THL  | Thalassionema+Lioloma                                 |
| THP  | Thalassiosira+Porosira                                |

Table 1: Name and composition of the phytoplanktonic groups used in the paper, based on<sup>1</sup>

| Name of site | Location                 | Region            | Number of points? <sup>1</sup> | Temperature             | Salin            |
|--------------|--------------------------|-------------------|--------------------------------|-------------------------|------------------|
| Men Er Roue  | 47°32'5" N / 3°5'37" W   | Brittany          | 503                            | 3.8-22.2 (14.4 +/- 3.7) | 20.1-38 (33.0)   |
| Loscolo      | 47°27'27" N / 2°32'18" W | Brittany          | 463                            | 5.7-22.4 (14.9 +/- 4.0) | 14.0-36.8 (32.0) |
| Croisic      | 47°18'0" N / 2°30'51" W  | Brittany          | 500                            | 4.8-28.9 (14.7 +/- 3.9) | 14.7-37.6 (31.0) |
| L'Eperon     | 46°16'13" N / 1°14'16" W | Oléron            | 460                            | 3.0-26.0 (15.3 +/- 4.8) | 13.0-36.6 (32.0) |
| Cornard      | 46°3'19" N / 1°7'50" W   | Oléron            | 491                            | 3.1-29.2 (15.6 +/- 4.8) | 19.0-38.1 (32.0) |
| Auger        | 45°47'59" N / 1°12'19" W | Oléron            | 524                            | 3.0-24.5 (15.4 +/- 4.4) | 23.9-36.0 (32.0) |
| Buoy7        | 44°32'32" N / 1°15'49" W | Arcachon          | 311                            | 7.2-23.9 (15.2 +/- 3.8) | 31.8-36.1 (34.0) |
| Teychan      | 44°40'25" N / 1°9'31" W  | Arcachon          | 494                            | 5.5-25.2 (15.5 +/- 4.6) | 20.6-35.8 (32.0) |
| Antoine      | 43°22'41" N / 4°50'45" E | Mediterranean Sea | 539                            | 4.6-30.0 (16.8 +/- 5.1) | 26.8-38.9 (32.0) |
| Lazaret      | 43°5'14" N / 5°54'23" E  | Mediterranean Sea | 512                            | 8.7-29.2 (17.4 +/- 4.2) | 21.6-39.6 (35.0) |

Table 2: Attempt of summary for our locations ; should we add the species for each region ?




Figure 2: Comparison of BIC with different interaction matrices, compared to the null model (diagonal interaction matrix, allowing only intragroup interactions), for four different sites separated by dashed lines (Brittany, Oléron, Arcachon and Mediterranean Sea) and 10 different subsites. Different interaction matrices may allow all interactions between taxa (unconstrained), only interactions within pennate diatoms, centric diatoms, dinoflagellates, or other phytoplanktonic taxa (pecten), only interactions within diatoms, dinoflagellates or other taxa (diatdin), or only interactions between taxa belonging to these different groups. As model structures (length of the times series taken into account) are different between sites and subsites, groups of bars should not be compared.

<sup>1</sup>From 1996, without linear interpolation




Figure 3: Interaction matrices estimated in Arcachon (a) and in the Mediterranean Sea (b). Only interactions between clades (pennate and centric diatoms, dinoflagellates, other planktonic taxa) are allowed. The figure should be read as taxon  $i$  having effect  $e_{ji}$  on taxon  $j$ . The scale for the coefficient values is given at the bottom left of panel a). 95% significance of coefficients was determined by bootstrapping and is marked by asterisks (\*). The identity matrix was subtracted to the interaction matrix ( $\mathbf{B}-\mathbf{I}$ ) in order to make effects on growth rates comparable. Composition of planktonic groups is given in Table 1.




Figure 4: Effect of abiotic variables (temperature, TEMP or salinity, SAL) on phytoplankton group in Brittany (a, b), Oléron (c, d), Arcachon (e, f) and in the Mediterranean Sea (g, h). Each color corresponds to a different site. Error bar corresponds to the 95% confidence interval around the estimated coefficient. All variables were normalized before estimation.




Figure 5: Variance of the coefficient in the interaction matrix (**B-I**), as a function of their mean, for 10 sites in 4 regions, with a model allowing interactions only within clads (see above). The mean-variance relation was either computed with raw values of intergroup interactions (a) or absolute values of the intergroup coefficients (b). Intragroup coefficients were not modified.

## References for the meta-analysis




Figure 6: Ratio of intra-to-intergroup interaction strength, when taking into account only the significant values (left), as a function of the density of the interaction matrix B-I (right). [This graph is not the final one but I'm wondering if we keep it]

| Code on the plot | Ref                                    | Dimension | Type of organisms             | System                     |           |
|------------------|--|-----------|-------------------------------|----------------------------|-----------|
| 1a               | [2], conditional least square estimate | 9         | Zooplankton                   | Lake                       | 107 point |
| 1b               | [2], total least square estimate       | 9         | Zooplankton                   | Lake                       |           |
| 2a               | [3]                                    | 2         | Phytoplankton                 | Lake                       | 10        |
| 2b               | [3]                                    | 3         | Zooplankton                   | Lake                       |           |
| 3a               | [4]                                    | 4         | Functional groups of plankton | Lake                       | 18        |
| 3b               | [4]                                    | 5         | Taxonomic groups of plankton  | Lake                       |           |
| 4a               | [5]                                    | 4         | Plankton                      | Lake                       |           |
| 4b               | [5]                                    | 4         | Plankton                      | Lake with high planktivory |           |
| 4c               | [5]                                    | 4         | Plankton                      | Lake with low planktivory  |           |
| 5a               | [6]                                    | 14        | Plankton                      | Lake                       |           |
| 5b               | [6]                                    | 14        | Plankton, growing season      | Lake                       |           |
| 6a               | [7]                                    | 13        | Plankton                      | Lake                       |           |
| 6b               | [7]                                    | 7         | Simpler web, plankton         | Lake                       |           |
| 7a               | [8]                                    | 10        | Ciliates                      | Lake                       |           |
| 7b               | [8]                                    | 10        | Phytoplankton                 | Lake                       |           |
| 8a               | [9]                                    | 3         | Insects                       | Terrestrial                |           |
| 9a               | [10]                                   | 2         | Lynx/Hare                     | Terrestrial                |           |
| 10a              | [11]                                   | 3         | Fish                          | Baltic Sea                 |           |
| 11a              | [12]                                   | 7         | Phytoplankton                 | Coastal site               |           |
| 11b              | [12]                                   | 7         | Phytoplankton                 | Offshore site              |           |
| 12a              | [13]                                   | 12        | Phytoplankton                 | Outside a bay              |           |
| 49412b           | [13]                                   | 12        | Phytoplankton                 | Inside a bay               |           |

Table 3: References used [TO COMPLETE]

## References

- [1] T. Hernández Fariñas, C. Bacher, D. Soudant, C. Belin, and L. Barillé. Assessing phytoplankton realized niches using a French national phytoplankton monitoring network. *Estuarine, coastal and shelf science*, 159:15–27, 2015.
- [2] A.R. Ives, S.R. Carpenter, and B. Dennis. Community interaction webs and zooplankton responses to planktivory manipulations. *Ecology*, 80(4):1405–1421, 1999.
- [3] J.L. Klug, J.M. Fischer, A.R. Ives, and B. Dennis. Compensatory dynamics in planktonic community responses to pH perturbations. *Ecology*, 81(2):387–398, 2000.
- [4] J.L. Klug and K.L. Cottingham. Interactions among environmental drivers: Community responses to changing nutrients and dissolved organic carbon. *Ecology*, 82(12):3390–3403, 2001.
- [5] A. R. Ives, B. Dennis, K. L. Cottingham, and S. R. Carpenter. Estimating community stability and ecological interactions from time-series data. *Ecological monographs*, 73(2):301–330, 2003.
- [6] Stephanie E. Hampton and Daniel E. Schindler. Empirical evaluation of observation scale effects in community time series. *Oikos*, 113(3):424–439, 2006.
- [7] Stephanie E. Hampton, Mark D. Scheuerell, and Daniel E. Schindler. Coalescence in the lake washington story: Interaction strengths in a planktonic food web. *Limnology and oceanography*, 51(5):2042–2051, 2006.
- [8] V. Huber and U. Gaedke. The role of predation for seasonal variability patterns among phytoplankton and ciliates. *Oikos*, 114(2):265–276, 2006.
- [9] K. Yamamura, M. Yokozawa, M. Nishimori, Y. Ueda, and T. Yokosuka. How to analyze long-term insect population dynamics under climate change: 50-year data of three insect pests in paddy fields. *Population ecology*, 48(1):31–48, 2006.
- [10] J.O. Vik, C.N. Brinch, S. Boutin, and N.C. Stenseth. Interlinking hare and lynx dynamics using a century’s worth of annual data. *Population ecology*, 50(3):267–274, 2008.
- [11] M. Lindegren, C. Möllmann, A. Nielsen, and N.C. Stenseth. Preventing the collapse of the Baltic cod stock through an ecosystem-based management approach. *Proceedings of the national academy of sciences*, 106(34):14722–14727, 2009.

- [12] J.R. Griffiths, S. Hajdu, A.S. Downing, O. Hjerne, U. Larsson, and M. Winder. Phytoplankton community interactions and environmental sensitivity in coastal and offshore habitats. *Oikos*, 125(8):1134–1143, 2015.
- [13] F. Barraquand, C. Picoche, D. Maurer, L. Carassou, and I. Auby. Coastal phytoplankton community dynamics and coexistence driven by intragroup density-dependence, light and hydrodynamics. *Oikos*, 127(12):1834–1852, 2018.